



## Physicochemical Properties and Bioanalytical Methods of Therapeutic Agents Utilized in the Management of Alzheimer's Disease

**Chika J Mbah\* and Chinwe M Onah**

*Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka, Nigeria*

**\*Corresponding Author:** Chika J Mbah, Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka, Nigeria.

**Received:** September 12, 2019; **Published:** September 13, 2019

**DOI:** 10.31080/ASPS.2019.03.0398

### Abstract

Alzheimer's disease (AD) is a neurodegenerative disease characterized by progressive deterioration in cognition, function and behavior. Patients therefore suffer from confusion, disorientation, impaired judgment, memory loss and have difficulty in expressing themselves. The objective of the study was to summarize literature reports on the physicochemical properties and bioanalytical methods of drugs used to treat Alzheimer's disease. Information was obtained from published works in scientific journals, official books as well the online search. The drugs currently used clinically to treat the disorder are donepezil, galantamine, memantine, rivastigmine and tacrine. Physicochemical properties of the drugs namely dissociation constants, melting points, partition coefficients and solubility are presented. Various analytical chromatographic methods (hyphenated and non-hyphenated) used to determine the drugs in biological fluids are discussed. Sample preparations were noted to involve extraction, centrifuging, addition of internal standard.

In general, hyphenated chromatographic methods (LC-ESI-MS/MS; LC-MS/MS; LC-APCI-MS/MS respectively) were observed to be the analytical methods of interest for the quantification of Alzheimer's drugs in biological matrices. In conclusion, the review of the literature has shown that accuracy, precision, specificity, sensitivity, rapidity might be responsible for the use of hyphenated chromatographic methods as the bioanalytical methods of choice.

**Keywords:** Alzheimer's Disease; Chromatographic Methods; Physicochemical Properties

### Abbreviations

AD: Alzheimer's Disease; APCI: Atmospheric Pressure Chemical Ionization; MRM: Multiple Reaction Monitoring; MS: Mass Spectrometry; ESI: Electrospray Ionization; NMDA: N-methyl-D-aspartate; RP: Reverse Phase; UV: Ultraviolet; SRM: Selected Reaction Monitoring; UPLC: Ultra-Performance Liquid Chromatography.

### Introduction

Neurodegenerative disease is a neurologic disorder characterized by neuronal loss and accumulation of insoluble extracellular or intracellular material in certain regions of the brain. The disorder which affects mostly the elderly is progressive and of unknown etiology. It could be of inherited form, but the occurrence could be

sporadic with genetic predisposition, aging and environmental factors as risk factors. Neurodegenerative disease could be:

- **Alzheimer's disease:** Most common cause of dementia, arising from neural injury occurring primarily in the cortex and hippocampus. It accounts for about 50% of clinically diagnosed dementia [1].
- **Amyotrophic lateral sclerosis:** The progressive weakness and muscle atrophy are as result of degeneration of bulbar, cortical and spinal neurons.
- **Huntington's disease:** A motor disorder caused by loss of a specific subset of striatal neurons and characterized by abnormal and excessive movements.
- **Parkinson's disease:** A disabling motor impairment disorder as a result of loss of nigrostriatal dopamine neurons.

The neurochemical changes in Alzheimer's disease form the basis for the symptomatic treatment with therapeutic agents such as donepezil, galantamine, memantine, rivastigmine and tacrine. The present study deals with Alzheimer's disease, focusing on the physicochemical properties and bioanalytical methods of therapeutic agents (drugs) employed in the treatment of the disease.

These agents are competitive and reversible cholinesterase inhibitors (AChEI), inhibiting the activity of enzyme cholinesterase and increasing the level of acetylcholine in brain. Furthermore, dysfunction of glutamatergic neurotransmission is involved in the etiology of the disease, hence the use of memantine, an N-methyl D-aspartate antagonist (NMDA) to treat moderate to severe form of the disease. Thus, these drugs are primarily used to improve motivation, anxiety level and confidence of Alzheimer's patients.

A number of analytical methods including capillary electrophoresis [2,3] and chromatographic methods have been used to determine these drugs in biological fluids however, chromatographic methods are the analytical methods mostly used. Biological fluids very often utilized are blood (whole blood, serum or plasma), urine, cerebrospinal fluid (CSF), bile and saliva.

In this context, we present the physicochemical properties as well as the bioanalytical methods of Alzheimer's drugs. They include:

Donepezil is chemically defined as 2,3-Dihydro-5,6-dimethoxy-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]1H-inden-1-one. It has

a molecular formula and molecular weight of  $C_{24}H_{29}NO_3$  and 379.49 g/mol respectively. The chemical structure is given in figure 1. It melts at 206-207 deg C. Donepezil is sparingly soluble in water, soluble in acetone, chloroform. The logarithm partition coefficient in octanol-water is 3.6 and has pKa value of 8.9.




Figure 1: Chemical structure of Donepezil.

Donepezil has been determined by numerous chromatographic methods, however only few will be described. They include;

#### Human plasma

- Lordachescu, *et al.* [4] hyphenated chromatographic system (LC-MS/MS, by MRM in positive mode) consisting of chiral RP-C<sub>18</sub> column, isocratic separation at room temperature using ammonium carbonate (pH 9.0) and acetonitrile as mobile phase at a flow rate of 0.3 ml/min, a total run time of about 8.0 min and a calibration range of 0.05-25 ng/ml. Internal standard was donepezil-d<sub>7</sub>.
- Khuroo, *et al.* [5] hyphenated chromatographic system (LC-ESI-MS/MS, by MRM in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at 35 deg. C using methanol and ammonium acetate (pH 6.2) as mobile phase at a flow rate of 0.4 ml/min, a total run time of about 9.0 min and a calibration range of 0.33-51 ng/ml. Internal standard was galantamine.
- Noetzli, *et al.* [6] hyphenated chromatographic system (UPLC-MS/MS, by MRM in positive mode) consisting of RP-C<sub>18</sub> column, gradient separation at room temperature using ammonium acetate (pH 9.3) and acetonitrile as mobile phase at a flow rate of 0.4 ml/min, a total run time of about 4.5 min and a calibration range of 1-300 ng/ml. Internal standard was galantamine radioisotope.
- Kim, *et al.* [7] hyphenated chromatographic system (LC-MS/MS, by MRM in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using ammonium acetate (pH 5) and acetonitrile as mobile phase at a flow rate of 0.2 ml/min, a total run time of about 5 min and a calibration range of 0.1-50 ng/ml. Internal standard was pioglitazone.

- Pilli, *et al.* [8] hyphenated chromatographic system (LC-MS/MS, by MRM in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using ammonium formate and acetonitrile as mobile phase at a flow rate of 0.6 ml/min, a total run time of about 3 min and a calibration range of 0.09-24.2 ng/ml. Internal standard was dipyridamole.

#### Human serum

Petrocheilou, *et al.* [9] non-hyphenated chromatographic system (UPLC-DAD) consisting of RP-C<sub>18</sub> column, gradient separation at room temperature using acetonitrile, methanol and buffer solution of sodium acetate/acetic acid, (0.2 M, pH 4.8) as mobile phase at a flow rate of 0.6 ml/min, a total run time of about 5 min. The same method was used to determine the drug in human cerebrospinal fluid and urine.

Galantamine, chemically is {[4aS-(4α,6β,8αR\*)]-4a,5,9,10,11,12-hexahydro-3-methoxy-11-methyl-6H-benzofuro-[3a,3,2ef]2benzazepin-6-ol}. It has a molecular formula and molecular weight of C<sub>17</sub>H<sub>21</sub>NO<sub>3</sub> and 287.35 g/mol respectively. The chemical structure is given in figure 2. It melts at 126-127 deg C. Galantamine is fairly soluble in hot water, freely soluble in alcohol, acetone, chloroform and less soluble in benzene and ether. The logarithm partition coefficient in octanol-water is 1.8 and has pKa value of 8.91, 14.81.




Figure 2: Chemical structure of Galantamine.

#### Galantamine has been determined by the following chromatographic methods

##### Human plasma

- Park, *et al.* [10] hyphenated chromatographic system (LC-ESI-MS/MS, in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using acetonitrile and ammonium acetate (0.01M) as mobile phase at a flow rate of 0.2 ml/min, a total run time of


about 2 min and a calibration range of 4-240 ng/ml. Internal standard was glimepride.

- Nirogi, *et al.* [11] hyphenated chromatographic system (LC-MS/MS, by MRM in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using 0.03% formic acid and acetonitrile as mobile phase at a flow rate of 1.5 ml/min, a total run time of about 3 min and a calibration range of 0.5-100 ng/ml. Internal standard was loratadine.
- Zhang, *et al.* [12] non-hyphenated chromatographic system (HPLC with fluorescence detector) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using acetonitrile, water and triethylamine (pH 7.0) as mobile phase at a flow rate of 1.0 ml/min, a total run time of about 15 min and a calibration range of 2-160 ng/ml. Internal standard was tramadol.
- Tencheva, *et al.* [13] non-hyphenated chromatographic system (HPLC with UV detector) consisting of RP-C<sub>8</sub> column, isocratic separation at room temperature using methanol, water and dibutylamine (pH 7.0) as mobile phase at a flow rate of 1.2 ml/min, a total run time of about 20 min and a detection limit of 0.05 µg/ml. Internal standard was codeine. The same method was used to determine the drug in human urine.
- Verhaeghe, *et al.* [14] hyphenated chromatographic system (LC-ESI-MS/MS, in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using acetonitrile and ammonium acetate (0.01M) as mobile phase at a flow rate of 1.5 ml/min, a total run time of about 3 min and a calibration range of 1-500 ng/ml. Internal standard was galantamine radioisotope.

##### Human serum

Claessens, *et al.* [15] non-hyphenated chromatographic system (HPLC with UV detector) consisting of normal phase silica column, isocratic separation at room temperature using 0.1% ethanol, dibutylamine, dichloromethane and n-hexane as mobile phase at a flow rate of 1.0 ml/min, a total run time of about 7 min and a calibration range of 10-100 ng/ml. Internal standard was phenacetin. The same method was used to determine the drug in human bile and urine.

Memantine defined chemically as 3,5-Dimethyladamantan-1-amine. It has a molecular formula and molecular weight of C<sub>12</sub>H<sub>21</sub>N and 179.3 g/mol respectively. The chemical structure is given in figure 3. It melts at 153-154 deg C. Memantine is soluble in water, freely soluble in alcohol and acetone. The logarithm partition coefficient in octanol-water is 3.28 and has pKa value of 10.7.




**Figure 3:** Chemical structure of Memantine.

### Memantine determination is mostly by the following chromatographic methods

#### Human plasma

- Konda., *et al.* [16] hyphenated chromatographic system (LC-ESI-MS/MS in positive mode) consisting of RP-C<sub>18</sub> column maintained at 40 deg C, isocratic separation at room temperature using acetonitrile and 0.1% formic acid as mobile phase at a flow rate of 0.6 ml/min, a total run time of about 4min and a calibration range of 50-50,000 pg/ml. Internal standard was mamantine-d<sub>6</sub>.
- Noetzli., *et al.* [17] hyphenated chromatographic system (LC-ESI-MS/MS in positive mode) consisting of RP-C<sub>18</sub> column, gradient separation at room temperature using ammonium acetate and acetonitrile as mobile phase at a flow rate of 0.8 ml/min, a total run time of about 15 min and a calibration range of 1-300 ng/ml. Internal standard was donepezil radioisotope.
- Zarghi., *et al.* [18] non- hyphenated chromatographic system (HPLC with fluorescence detector) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using acetonitrile and phosphate buffer (0.025M, pH 4.6) as mobile phase at a flow rate of 2.5 ml/min, a total run time of about 10 min and a calibration range of 2-80 ng/ml. Internal standard was amantadine.
- Pan., *et al.* [19] hyphenated chromatographic system (LC-ESI-MS/MS in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using methanol and 0.5% formic acid as mobile phase at a flow rate of 0.45 ml/min, a total run time of about 5 min and a calibration range of 0.1-25 ng/ml. Internal standard was procainamide HCl.
- Almeida., *et al.* [20] hyphenated chromatographic system (LC-ESI-MS/MS by SRM in positive mode) consisting of RP-C<sub>18</sub> column, isocratic separation at room temperature using methanol, water and formic acid as mobile phase at a flow rate of 0.15 ml/min, a total run time of about 3 min and a calibration range of 0.1-50 ng/ml. Internal standard was amantadine.

Rivastigmine, is chemically defined as [3-[(1S)-1-(dimethylamino)ethyl]phenyl] N-ethyl-N-methylcarbamate. It has a molecular formula and molecular weight of C<sub>14</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub> and 250.34 g/mol respectively. The chemical structure is given in figure 4. It melts at 121-123 deg C. Rivastigmine is sparingly soluble in water, very soluble in anhydrous ethanol and heptane. The logarithm partition coefficient in octanol-water is 2.3 and has pKa value of 8.85.




**Figure 4:** Chemical structure of Rivastigmine.

### Rivastigmine has been determined by the following chromatographic methods

#### Human plasma

- Amini and Ahmadiani [21] non-hyphenated chromatographic system (HPLC with UV detector) consisting of silica column maintained at 50 deg C, isocratic separation at room temperature using acetonitrile and sodium dihydrogen phosphate (pH 3.1) as mobile phase at a flow rate of 1.3 ml/min, a total run time of about 10 min and a calibration range of 0.5-16 ng/ml. Internal standard was donepezil
- Bhatt., *et al.* [22] hyphenated chromatographic system (LC-MS/MS by SRM in positive mode) consisting of RP-C<sub>8</sub> column maintained at 45 deg C, isocratic separation at room temperature using 0.1% formic acid and acetonitrile as mobile phase at a flow rate of 1 ml/min, a total run time of about 2 min and a calibration range of 0.2-20 ng/ml. Internal standard was zolpidem..
- Frankfort., *et al.* [23] hyphenated chromatographic system (LC-MS/MS by MRM in positive mode) consisting of RP-C<sub>18</sub> column, gradient separation at room temperature using ammonium hydroxide and methanol as mobile phase at a flow rate of 0.2 ml/min, a total run time of about 14 min and a calibration range of 0.25-25 ng/ml. There was no internal standard.
- Pommier and Frigola [24] hyphenated chromatographic system (LC-APCI-MS/MS by MRM in positive mode) consisting of RP-C<sub>8</sub> column maintained at 45 deg C, isocratic separation at room temperature using methanol and ammonium acetate (0.02 M) as mobile phase at a flow rate of 0.2 ml/min, a total run time of about 5 min and a calibration range of 0.2-30 ng/ml. Internal standard was rivastigmine d<sub>6</sub>.

Tacrine, is chemically defined as 1,2,3,4-Tetrahydroacridin-9-amine. It has a molecular formula and molecular weight of  $C_{13}H_{14}N_2$  and 198.26 g/mol respectively. The chemical structure is given in figure 5. It melts at 183-184 deg C. Tacrine is sparingly soluble in water; soluble in alcohol. The logarithm partition coefficient in octanol-water is 2.2 or 2.71 and has pKa value of 9.95.



**Figure 5:** Chemical structure of Tacrine.

### Tacrine has been determined by the following chromatographic methods

#### Human plasma

- Han., *et al.* [25] hyphenated chromatographic system (LC-MS/MS by MRM in positive mode) consisting of RP  $C_{18}$  maintained at 30 degree C, isocratic separation at room temperature using ammonium acetate (10 mM), 1% formic acid and acetonitrile as mobile phase at a flow rate of 1 ml/min, a total run time of about 3 min and a calibration range of 0.01-10 ng/ml. Internal standard was tramadol.
- Hansen., *et al.* [26] non-hyphenated chromatographic system (HPLC with fluorescence detector) consisting of RP  $C_8$ , isocratic separation at room temperature using ammonium acetate (0.2 M) and acetonitrile as mobile phase at a flow rate of 1.3 ml/min, a total run time of about 40 min and a calibration range of 6-60 nM. Internal standard was 1,2,3,4-Tetrahydro-9-acridanone. The method was also used to determine the drug in human urine.
- Ekman., *et al.* [27] non-hyphenated chromatographic system (HPLC with UV detector) consisting of RP  $C_{18}$  maintained, isocratic separation at room temperature using phosphate buffer and acetonitrile as mobile phase at a flow rate of 1.1 ml/min, a total run time of about 14 min and limit of detection of 0.3ng/ml. There was no internal standard.

#### Human serum

Forsyth., *et al.* [28] non-hyphenated chromatographic system (HPLC with fluorescence detector) consisting of RP  $C_{18}$ , isocratic separation at room temperature using methanol, water, 1% triethylamine (pH 5) as mobile phase at a flow rate of 1.5 ml/min, a total run time of about 8 min and a calibration range of 1-20 ng/ml. Internal standard was 1,2,3,4-Tetrahydro-9-acridanone.

The study has shown that in the chromatographic analyses, the biological fluids containing the drugs would be treated to free the drugs from interferences prior to injecting into the chromatograph. Treatment may involve cloud-point extraction, liquid-liquid extraction, microextraction, protein precipitation, solid-phase extraction. Due to the fact that biological fluids contain proteins in significant amounts, protein precipitation is very often carried out using solvents such as acetonitrile and/or methanol prior to other treatments.

### Conclusion

Alzheimer's therapeutic agents (drugs) are cholinesterase inhibitors and N-methyl-D-aspartate (NMDA) receptor antagonists and act by increasing the concentration and duration of action of acetylcholine or reducing the glutamatergic overstimulation in brain. The physicochemical data have shown that these drugs do not have good aqueous solubility but possess very favourable partition coefficients. Accurate determination of these drugs in biological fluids, are very vital because it assists clinicians to make better decision with regards to dosage selection and dosage regimens respectively. Amongst the hyphenated analytical methods, LC-MS/MS or LC-ESI-MS/MS method has proven to be the bioanalytical method of choice. Only few non-hyphenated bioanalytical methods have been used to determine Alzheimer's disease drugs. Finally, the study has shown that most of these bioanalytical methods have adequate accuracy, precision, specificity, sensitivity and rapidity to allow the estimation of Alzheimer's drugs from whole blood, plasma, serum, urine and other biological matrices.

### Bibliography

1. Areosa SA and Sherriff F. "Memantine for dementia". *The Cochrane Database of Systematic Reviews* 3 (2003).
2. Nicolaou IN and Kapnissi-Christodoulou CP. "Simultaneous determination of nine acetylcholinesterase inhibitors using micellar electrokinetic chromatography". *Journal Chromatographic Science* 49 (2011): 265-271.



3. Nicolaou I and Kapnissi-Christodoulou CP. "Development of a capillary electrophoresis-mass spectrometry method for the determination of rivastigmine in human plasma- optimization of the limits of detection and quantitation". *Method Development by Use of Capillary Electrophoresis and Applications in Pharmaceutical, Biological and Natural Samples*, INTECH (2012).
4. Iordachescu A., et al. "LC-MS-MS method for the simultaneous determination of donepezil enantiomers in plasma". *Chromatographia* 75 (2012): 857-866.
5. Khuroo AH., et al. "ESI-MS/ MS stability-indicating bioanalytical method development and validation for simultaneous estimation of donepezil, 5-desmethyl donepezil and 6-desmethyl donepezil in human plasma". *Biomedical Chromatography* 26 (2012): 636-649.
6. Noetzli M., et al. "Simultaneous determination of antidementia drugs in human plasma: procedure transfer from HPLC-MS to UPLC-MS/MS". *Journal of Pharmaceutical and Biomedical Analysis* 64-65 (2012): 16-25.
7. Kim KA., et al. "Pharmacokinetic comparison of orally disintegrating and conventional donepezil formulations in healthy Korean male subjects: a single-dose, randomized, open-label, 2sequence, 2-period crossover study". *Clinical Therapeutics* 33 (2011): 965-972.
8. Pilli NR., et al. "A rapid and sensitive LC-MS/MS method for quantification of donepezil and its active metabolite, 6-O-desmethyl donepezil in human plasma and its pharmacokinetic application". *Biomedical Chromatography* 25 (2011): 943-951.
9. Petrocheilou M., et al. "A simple and direct HPLC-DAD method for the simultaneous determination of galantamine, donepezil and rivastigmine in cerebrospinal fluid, blood serum and urine". *Journal Applied Bioanalysis* 3 (2017): 59-69.
10. Park YS., et al. "Quantification of galantamine in human plasma by validated liquid chromatography-tandem mass spectrometry using glimepiride as an internal standard: application to bioavailability studies in 32 healthy Korean subjects". *Journal of Chromatographic Science* 50 (2012): 803-809.
11. Nirogi RVS., et al. "Quantitative determination of galantamine in human plasma by sensitive liquid chromatography-tandem mass spectrometry using loratadine as an internal standard". *Journal Chromatographic Science* 45 (2007): 97-103.
12. Zhang LJ., et al. "Pharmacokinetics and bioequivalence studies of galantamine hydrobromide dispersible tablet in healthy male Chinese volunteers". *Drug Development and Industrial Pharmacy* 33 (2007): 335-340.
13. Tencheva J., et al. "Reversed-phase liquid chromatography for the determination of galanthamine and its metabolites in human plasma and urine". *Journal Chromatography* 421(1987): 396-400.
14. Verhaeghe T., et al. "Development and validation of a liquid chromatographic-tandem mass spectrometric method for the determination of galantamine in human heparinised plasma". *Journal of Chromatography B* 789 (2003): 337-346.
15. Claessens HA., et al. "High performance liquid chromatographic determination of galanthamine, a long acting anticholinesterase drug, in serum, urine and bile". *Journal Chromatography* 275 (1983): 345-353.
16. Konda RK., et al. "Bioanalytical method development and validation of memantine in human plasma by high performance liquid chromatography with tandem mass spectrometry: application to bioequivalence study". *Journal Analytical Methods in Chemistry* 2012 (2012): 101249.
17. Noetzli M., et al. "Simultaneous determination of antidementia drugs in human plasma for therapeutic drug monitoring". *Therapeutic Drug Monitoring* 33 (2011): 227-238.
18. Zarghi A., et al. "Sensitive and rapid HPLC method for determination of memantine in human plasma using OPA derivatization and fluorescence detection: application to pharmacokinetic studies". *Scientia Pharmaceutica* 78 (2010): 847-856.
19. Pan RN., et al. "Determination of memantine in human plasma by LC-MS-MS. Application to a pharmacokinetic study". *Chromatographia* 270 (2009): 783-788.
20. Almeida AA., et al. "Determination of memantine in human plasma by liquid chromatography electrospray tandem mass spectrometry: application to a bioequivalence study". *Journal of Chromatography B* 848 (2007): 311-316.
21. Amini H and Ahmadiani A. "High performance liquid chromatographic determination of rivastigmine in human plasma for application in pharmacokinetic studies". *Journal Pharmaceutical Research* 9 (2010): 115-121.

22. Bhatt J., *et al.* "A rapid and sensitive liquid chromatography tandem mass spectrometry (LC-MS/MS) method for the estimation of rivastigmine in human plasma". *Journal Chromatography B* 852 (2007): 115-121.
23. Frankfort SV, *et al.* "A simple and sensitive assay for the quantitative analysis of rivastigmine and its metabolite NAP 226-90 in human EDTA plasma using coupled liquid chromatography and tandem mass spectrometry". *Rapid Communications in Mass Spectrometry* 20 (2006): 3330-3336.
24. Pommier F and Frigola R. "Quantitative determination of rivastigmine and its major metabolite in human plasma by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry". *Journal of Chromatography B* 784 (2003): 301-313.
25. Han J., *et al.* "Quantitation of the tacrine analogue octahydroaminoacridine in human plasma by liquid chromatography-tandem mass spectrometry". *Journal Pharmaceutical Biomedical Analysis* 50 (2009): 171-174.
26. Hansen LL., *et al.* "Determination of tacrine and its metabolites in human plasma and urine by high performance liquid chromatography and fluorescence detection". *Journal of Chromatography B* 712 (1998): 183-191.
27. Ekman L., *et al.* "Determination of tacrine and its 1-hydroxy metabolite in plasma using column liquid chromatography with ultraviolet detection". *Journal Chromatography* 494 (1989): 397-402.
28. Forsyth DR., *et al.* "Determination of tacrine hydrochloride in human serum by chloroform extraction, reversed phase high performance liquid chromatography and fluorimetric detection". *Journal Chromatography* 433 (1988): 352-358.

**Volume 3 Issue 10 October 2019**

**© All rights are reserved by Chika J Mbah and Chinwe M Onah.**