



Assessment of Prevalence and Level of Knowledge of Polycystic Ovarian Syndrome (PCOS) Among Nursing College students at AIIMS, Raipur

Vahitha*

Department of Women's Health, India

*Corresponding Author: Vahitha, Department of Women's Health, India.

Received: November 21, 2023

Published: February 06, 2024

© All rights are reserved by Vahitha*.

Abstract

Introduction and Background: Polycystic ovarian syndrome (PCOS) is the most common endocrine disorder among girls. It has a wide range of symptoms such as irregular cycles, infertility, obesity, hirsutism and acanthosis nigran. Many factors influence PCOS such as genetics, environment, lifestyle changes, sedentary life and diet. The present study was conducted to assess the prevalence and level of knowledge of Polycystic Ovarian Syndrome (PCOS) among college girls and find out the association on level of knowledge with selected demographic variables.

Materials and Methods: This descriptive cross-sectional study was done among Nursing students in a tertiary care hospital and teaching institute in Raipur, India. Students who were willing to participate were included in the study. Informed consent was obtained from all the participants of the study. Institute Research Review committee and Ethical clearance from Institutional Ethics Committee (IEC) was obtained. A total of 180 participants were included and used convenient sampling method to collect the data. The participants were assured that participation was voluntary, and confidentiality would be maintained. The questionnaires were given after describing the purpose of the study and it was pretested semi-structured questionnaire was devised and validated. The first part of the questionnaire covered the socio-demographic details. The second part of the questionnaire included the questions about their level of knowledge and clinical symptoms about polycystic ovarian syndrome. After obtaining the data, it was entered in excel spread sheet and analyzed using SPSS software version 16, statistical analysis were done by using Chi-square test.

Results: The study revealed that majority of the students 162 (90%) had poor knowledge (<11) and 18 (10%) had adequate level of knowledge (≥ 12) respectively and majority of the students 129 (71.7%) had suspected and 51 (28.3%) had diagnosed in clinical evaluation respectively.

Conclusion: Thus, this study helps to know about the prevalence and level of knowledge and possibility of suspecting symptoms among Nursing students and also to diagnose it early, so that lifestyle changes and appropriate actions can be taken.

Keywords: PCOS; Nursing Students

Introduction

Adolescence is a dynamic phase of rapid growth and development during this stage physical, physiological and behavioural changes occur in our body [1]. They constitute more than 1.2 billion worldwide, and about 21% of the Indian population [2]. Adolescent girls and young women aged 15-24 years constitute approximately 880 million of the world's population [3]. Adolescence is divided into four periods early adolescence includes ages 10-12 years, middle adolescence ages 13-15 years, late adolescence ages 16-18 years, and young adults ages 19-25 years [4]. Menarche symbolizes the onset of sexual development and is characterized by the onset of the first menstrual bleeding.

The average age at menarche is 13.8 years [3]. Puberty is considered a sequence of events that happens in an individual where physical changes occur, resulting in physical characteristics devel-

opment and capacity to reproduce. These physical changes are regulated by hormones that are produced by the pituitary gland such as luteinizing hormone and follicle-stimulating hormone. In the early stage of puberty, levels of luteinizing hormone and follicle-stimulating hormone increase, stimulating the production of sex hormones. The increased levels of sex hormones (primarily estrogen) result in physical changes, including maturation of the breasts, ovaries, uterus, and vagina [4].

Normally, these changes occur sequentially during puberty, resulting in sexual maturity [5]. PCOS is a very complex syndrome, with typical imbalances of hormones and metabolic factors. Lifestyle modification focusing on diet and exercise behavioral modification is preferred as the first-line treatment for PCOS. Several studies have shown, weight loss of 5-10% of total body weight in overweight women with PCOS with the help of balanced nutrition

and exercise training which can lead to a reduction of central fat deposition, reduced circulating insulin and androgen levels, improved insulin sensitivity and restoring ovulation, improving menstrual cycles and decrease cardiovascular disease risk factors [6].

In one of study done in Bhopal 2017, the estimated prevalence of PCOS is 9.1% in the population among the age group of 15-21 years [1]. Women with PCOS are at an increased risk for infertility, preeclampsia, early pregnancy loss, and endometrial cancer. Moreover, because of the association of PCOS with insulin resistance, evidence suggests that women with PCOS are at an increased risk for developing type-2 diabetes, dyslipidemia, hypertension, and heart disease [7].

In a study done in Maharashtra shows 58.3% of the student confirmed with PCOS have a positive history of diabetes mellitus [3]. Certain normal physiologic changes that occur during adolescence can minimal symptoms of PCOS, including oligomenorrhoea, acne, and polycystic ovaries. Physical evaluation for persistent oligomenorrhoea is necessary to determine early signs of PCOS, especially when the symptoms persists 2 years beyond menarche need to be follow up [4]. One study done in Mangalore showing , there 76% of the participate were with average knowledge and 10.7% with good knowledge regarding polycystic ovarian syndrome [7]. Since early detection and management is key to alleviation of PCOS symptoms intended to generate awareness so that students could seek medical help [8]. The present study was conducted to assess the prevalence and level of knowledge of Polycystic Ovarian Syndrome (PCOS) among colleges girls and find out the association on level of knowledge with selected demographic variables [8].

Methodology

- **Research design:** The research design adopted for the study was descriptive cross sectional design.
- **Study setting:** This study was conducted in a tertiary care hospital and teaching institute, Raipur where the students are studying in a undergraduate nursing programme.
- **Population and Sampling methods:** The sampling population for the study includes college’s girls in Nursing College, Raipur and they are available during the course of data collection. Convenient sampling method was used to collect the data. Students were included as girls aged 18 to 22 years and those who have attained menarche. Girls who are taking treatment for PCOS and having other co morbidities were excluded from the study.

Sample size calculated based on estimation of single proportion, considering an expected prevalence of PCOS as 8.34% among girls aged 18-22 years, Gupta., *et al.* with an absolute precision of 5% and 95% confidence level, the sample size is 180.

Study tool and scoring

A self-developed questionnaire was used for which both content and face validation was done by the experts of Faculty of Nursing. The questionnaire consists of three domains: First part is concerned with demographics of the respondents. Second part of questionnaire contained 20 questions about PCOS Knowledge. The third part of questionnaire consists of clinical evaluation regarding the prevalence of PCOS. There were total 20 knowledge questions and scoring will be done by making cutoff value of 11. The score ≤ 11 will be considered as poor knowledge while score ≥ 12 will be considered as adequate knowledge. For clinical evaluation, a total of 12 sign and symptoms were given and students having 4-8 symptoms were considered as suspected while students having more than 8 symptoms were considered as diagnosed.

Data collection procedure

After obtaining permission from Institute Research Review Committee and Institute Ethics Committee, written consent was obtained from participants. A formal administrative permission was obtained from the Principal and Dean (Academics) in AIIMS Raipur The information was collected by using self administered questionnaires. The result was evaluated based on response received from participants in the questionnaire. The questionnaire was distributed among female students and responses were collected. After collecting filled questionnaires, education about PCOS was provided through brochure (written material) and short lecture was delivered to students.

Ethical aspects

Formal approval was obtained from Institute Research Review Committee, Institute Ethics Committee and formal administrative permission was obtained from the Principal and Dean (Academics) in AIIMS Raipur.

Data analysis

Categorical variable such as mother’s occupation, father’s occupation, religion, residence History of PCOS, and family history will be expressed as frequency and percentage. The continuous variable such as age, height, weight , age of menarche, hip and waist ratio, level of knowledge will be expressed as mean with standard derivation or median with inter quartile range according to the assumption of normality. The association of risk factor of PCOS with socio –demographic variables will be assessed using Chi- square test or Fisher’s exact test. All statistical analysis will be carried out in SPSS version 19 and p – value < 0.05 is considered statistically significant.

Results

The present study was conducted among the Nursing students in a tertiary teaching institute, Raipur. A total of 180 participants

were included in this study. Table 1 shows frequency and percentage wise distribution of level of knowledge of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur. Majority of the students 162 (90%) had poor knowledge and 18 (10%) had adequate level of knowledge respectively.

Level of knowledge	Frequency (n)	Percentage (%)
Poor knowledge	162	90
Adequate knowledge	18	10
Total	180	100

Table 1: Frequency and percentage wise distribution of level of knowledge of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur (N = 180).

Table 2 shows frequency and percentage wise distribution of Clinical evaluation of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur. Majority of the students 129 (71.7%) had suspected and 51(28.3%) had diagnosed in Clinical evaluation respectively.

Clinical evaluation of Polycystic Ovarian Syndrome (PCOS)	Frequency (n)	Percentage (%)
Suspected	129	71.7
Diagnosed	51	28.3
Total	180	100

Table 2: Frequency and percentage wise distribution of Clinical evaluation of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur (N = 180).

Table 3 shows that Correlation between the knowledge and Clinical evaluation of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur indicates the positive correlation and shows the results r- value is (0.226), p-value is (0.002) are significant.

Correlation	Mean	Standard deviation	'r' value	'p' value	Correlation
Knowledge	13.70	1.891	0.226	0.002	Positive
Clinical evaluation	6.37	2.055		**S	

Table 3: Correlation between the knowledge and Clinical evaluation of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur (N = 180).

*-p < 0.001 highly significant. NS-Non significant.

Discussion

Adolescence is a unique period where we can observe many change from puberty to adulthood. It is a period of changes in different levels such as in physical, psychological, social interaction with others and emotion adjustment with changing life style. After puberty, every individual undergoes physical, sexual, emotional,




Figure 1

psychological changes which cause imbalance due to changes in hormones level. These changes play an important role to understand the health risks which may be associated with PCOS or others genetic factors which can indirectly affect the health.

PCOS is a common health issue among teenagers and young women due to sedentary life style, lack of exercise. It has affected 5% to 10% young girls in their reproductive phase. One of the complications we can see is infertility. There is no cure for PCOS but it is treatable and manageable with life style modification.

Majority of the students (92) are 19 years of age (86.7%), studying second year 52(28.9), most of their parents are self employed 109(60.6%), attained menarche at the age of 13 years 71(39.9%), regarding BMI, most of them were in normal 116(64.6%) , waist hip ratio was high in 133(73.9%). The study also found that majority of the students, 85 (47.2%) who were not doing any extracurricular activity. The present study also found that Correlation between the knowledge and Clinical evaluation of Polycystic Ovarian Syndrome (PCOS) among Nursing College students at AIIMS, Raipur indicates the positive correlation and shows the results r- value is (0.226), p-value is (0.002) are significant. Majority of the students 129(71.7%) had suspected and 51(28.3%) had diagnosed in Clinical evaluation respectively.

The study conducted by Krithika S., *et al.* shows the similarity among college students. An observational study conducted among 300 adolescent girls aged between 14-18years and the data was collected by questionnaire and modified Ferriman - Gallwey scal. Mean, t- test and chi- square test were used. The study findings showed that 96.3% were not aware of PCOS .The study revealed the prevalence rate of PCOS is about 12.3% .The mean score regarding level of knowledge before and after the structured teaching programme was 13.25 and 23.16 respectively.

Another study, Mihika Aggarwal., *et al.* (2019) conducted a cross sectional study on prevalence of PCOS and risk factors associated

with it among medical students. The aim of the study is to find the prevalence of PCOS and its risk factors associated with it among women, where 456 medical, dental, physiotherapy students in the age group of 17-24 years took part in this study. A self-administered questionnaire was prepared on the basis of the Cronin, *et al.* questionnaire was applied to collect data and Pearson chi-square and mean test were used. Among those with PCOS, the mean age was 21.18 years. From this study prevalence was 21.05%, girls are aware of PCOS, 22.22% of the subjects were at high risk and 77.77% were at low risk for PCOS.

Mahesh Gupta, *et al.* (2018) conducted a cross sectional study of polycystic ovarian syndrome among young women in Bhopal, Central India, to find the prevalence of PCOS. A structured self-administrative questionnaire along with interview technique was applied to collect the data and total 500 college girls were included in the study. Mean and chi-square are used to calculate the result. From this study prevalence was 8.2% and 21.6% girls are aware of PCOS. Among all the risk factors, BMI ≥ 25 (P value < 0.0001) and waist hip ratio ≥ 0.85 . Lack of awareness was found among majority of girls (78.4).

Conclusion

This present study was conducted to find out prevalence and level of knowledge among nursing students in a tertiary teaching institute, Raipur. It reveals that, to bring more awareness among the Nursing students and modify the lifestyle changes to prevent the occurrence of the symptoms. Hence, it is necessary for us to decrease its incidence and complications. Early diagnosis of PCOS and its prompt treatment will help the girls to improve quality of life.

Acknowledgement

The authors of the study would like to thank the institution and members who facilitated and helped us to carry out the study. We would also like to thank all the study participants who participated in this study.

Bibliography

1. Sreepoorna Pramodh. "Exploration of Lifestyle Choices, Reproductive Health Knowledge, and Polycystic Ovary Syndrome (PCOS) Awareness Among Female Emirati University Students". *International Journal of Women's Health* 12 (2020): 927-938.

2. Mahesh G., *et al.* "A cross sectional study of polycystic ovarian syndrome among young women in Bhopal, Central India". *International Journal of Community Medicine and Public Health* 5 (2018): 95-100.

3. Mihika A., *et al.* "A cross sectional study on prevalence of PCOS and risk factors associated with it among medical students". *IJOGR* 6 (2019): 522-526.

4. Gupta M., *et al.* "A study to assess the prevalence of polycystic ovarian disease among girls aged 15-21 years from selected schools and colleges in Bhopal City". *Indian Journal of Youth and Adolescent Health* 4 (2017): 2-5.

5. Kaewnin J., *et al.* "Prevalence of polycystic ovary syndrome in Thai university adolescents". *Gynecological Endocrinology* 34 (2017): 476-480.

6. Sunanda B and Sabitha Nayak. "A Study to Assess the Knowledge Regarding PCOS (Polycystic Ovarian Syndrome) among Nursing Students at NUINS". *NUJHS* 6 (2016): 24-26.

7. Kalavathi DB and Amrita NS. "A descriptive study of polycystic ovarian syndrome in adolescent girls among a tertiary care hospital of Bangalore". *International Journal of Applied and Basic Medical Research* 4 (2015): 453-457.

8. Kirthika SV., *et al.* "Effect of structured awareness programme on polycystic ovarian syndrome (PCOS) among adolescent girls research". *Journal of Pharmacy Technology* 12 (2019): 6097-6100.

9. Varughese AK and Tauro VG. "Effectiveness of structured teaching programme (STP) on knowledge regarding polycystic ovarian syndrome (PCOS) among adolescent girls, Kollam". *International Journal of Nursing Education and Research* 6 (2018): 360-362.

10. Karkar M., *et al.* "A study to assess the knowledge regarding polycystic ovarian syndrome among undergraduate students in selected colleges of Pune city". *The Pharma Innovation Journal* 8 (2019): 192-194.

11. Fernandez RC., *et al.* "Sleep disturbances in women with polycystic ovary syndrome: Prevalence, pathophysiology, impact and management strategies". *Nature and Science of Sleep* 10 (2018): 45-64.

12. Bharathi RV., *et al.* "An epidemiological survey: Effect of predisposing factors for PCOS in Indian urban and rural population". *Middle East Fertility Society Journal* 22 (2019): 313-316.

13. Nidhi R., *et al.* "Prevalence of polycystic ovarian syndrome in Indian adolescents". *Journal of Pediatric and Adolescent Gynecology* 24.4 (2011): 223-227.

14. Nimo Biam B P. "Effectiveness of Self Instructional Module on Knowledge Regarding Polycystic Ovary Syndrome among Engineering Students". *International Journal of Novel Research in Healthcare and Nursing* 2.3 (2015): 66-69.

15. Rahman S., *et al.* "Study of the Effect of Food Habit, Lifestyle and Daily Trip on Physical and Mental Status of Subjects at Islamic University in Kushtia, Bangladesh". *Open Journal of Statistics* 2.2 (2012): 219.