



Role of Endophytes in Insect Control

Anjitha George*

Scientist, ICAR-Central Citrus Research Institute, Nagpur, India

*Corresponding Author: Anjitha George, Scientist, ICAR-Central Citrus Research Institute, Nagpur, India.

Received: September 28, 2017; Published: October 03, 2017

Plants have a complex immune system that defends them against herbivores and microbial pathogens but that also regulates the interactions with mutualistic organisms (e.g. mycorrhizal fungi and plant growth-promoting rhizobacteria). Endophytes are such microorganisms that live within plants for at least a part of their life cycle without causing any visible manifestation of disease [1]. The presence of endophytes is symptomless, unobtrusive, and established entirely inside the living host plant tissues [2]. During this association, none of the interacting partners is discernibly harmed, and the individual benefits depend on both the interacting partners. It is also believed that endophytes have important roles in plant protection, acting against herbivores, insects and pathogens of the host and may also increase plant resistance to pathogens and biotic and abiotic stresses [3,4].

The most frequently encountered endophytes are fungi [5], and currently, to our knowledge, all reported endophytes are fungi or bacteria (including actinomycetes). The existence of fungi inside the organs of asymptomatic plants has been known since the end of the 19th century [6] and the term “endophyte” was first proposed in 1866 [7]. It has been estimated that there may be as many as one million different endophytic fungal taxa, thus endophytes may be hyper diverse [8]. Most of the taxonomic groups contain entomopathogenic genera, such as *Metarhizium*, *Beauveria*, *Lecanicillium*, *Nomuraea*, *Entomophthora*, and *Neozygites* to name a few [9].

Bacterial endophytes colonize an ecological niche similar to that of phytopathogens, which makes them suitable as biocontrol agents [10,11]. Endophytic bacteria have been isolated from both monocotyledonous and dicotyledonous plants, ranging from woody tree species, such as oak and pear to herbaceous crop plants such as sugar beet and maize. Many endophytes are members of common soil bacterial genera, such as *Pseudomonas*, *Burkholderia* and *Bacillus* [12]. These genera are well known for their diverse range of secondary metabolic products including antibiotics, anti-cancer compounds, volatile organic compounds, antifungal, antiviral, insecticidal and immunosuppressant agents.

Scope of Endophytes as Biocontrol Agent

Use of entomopathogenic microorganisms or those that inhibit/antagonise other microorganisms pathogenic to plants is an alternative that may contribute to reduce or eliminate the use of chemical products in agriculture. Fungal biocontrol agents are promis-

ing because they act by contact and do not require ingestion, they can be mass-produced very easily and are quite host specific. Much has been published on the effects of endophytes on insect herbivores have concentrated on turf and agronomic grasses infested with endophytic clavicarpitacean fungi (Ascomycota: Hypocreales: Clavicarpitaceae), which systemically infect mostly grasses in the Poaceae, Juncaceae, and Cyperaceae. Endophyte-infested grasses have also shown high resistance to foliar-feeding insects and have been attributed to having biologically active alkaloids in infested grasses which alter the life cycle of the insect (antibiosis effect) [13,14]. Commonly cited example is that of *Neotyphodium*-infested perennial ryegrass (*Lolium perenne* L.) and tall fescue (*Festuca arundinacea* Schreb.) which has shown to have negative effects on over 40 insect species in six orders [15]. Clement, et al. [16] reported different effects on two aphids (bird-cherry oat aphid, *Rhopalosiphum padi* (L.) and rose grass aphid, *Metopolophium dirhodum* (Walker)) and the wheat stem sawfly (*Mayetiola destructor* (Say)) exposed to different wild barleys infected with *Neotyphodium*. *Beauveria bassiana* (Balsamo) Vuillemin has been reported as an endophyte in maize [17-19], potato, cotton, cocklebur, and jimson weed [20], tomato [21,22], in sorghum, chilli, sunflower and beans [23], on the cocoa relative *Theobroma gileri* [24], in the bark of *Carpinus caroliniana* Walter [25], in seeds and needles of *Pinus monticola* Dougl. ex. D. Don [26], in opium poppy [27], on date palm [28], in bananas [29], and in coffee [30]. In addition, *Lecanicillium lecanii* (Zimm.) has been reported in sorghum, cotton, wheat, chickpea, pigeon pea, mango and banana [23]; *Paeecilomyces* sp. in *Musa acuminata* [31] and in rice [32].

Endophytic bacteria have been found in virtually every plant studied, where they colonize the internal tissues of their host plant and can form a range of different relationships including symbiotic, mutualistic, commensalistic and trophobiotic. Endophytic bacteria can promote plant growth and yield and can act as biocontrol agents. Bacterial endophytes colonize an ecological niche similar to that of phytopathogens, which makes them suitable as biocontrol agents [10,11]. Bacteria, especially the genus *Bacillus*, have significant participation among the commercialized biological control products. Up to 50% of these products are bacterial formulations, from various species of *Bacillus*. The identification of different action mechanisms can lead to the combination of isolates to control a broad spectrum of pests [33].

Akello, *et al.* [34] studied the effect of endophytic *B. bassiana* in banana plants on the banana weevil, *Cosmopolites sordidus* (Germar) and reported dead mycosed insects in the rhizome of *B. bassiana*-inoculated plants, suggesting a direct mode of action through mycosis infection of the banana weevils by the fungus. Endophytic colonization of banana by *B. bassiana* significantly reduced the survivorship of grubs of banana weevil resulting in 42 - 87 per cent reduction in plant damage Date palm pulp of the endophytically colonized seedlings was used in a laboratory diet for the grubs of the red palm weevil (RPW). The mortality of grubs during 14 days achieved 80.3 per cent, under laboratory conditions [35]. Zhou, *et al.* [36] reported that endophytic fungus, *Chaetomium globosum* Kunze inhibited root-knot nematode, *Meloidogyne incognita* (Kofoid & White) infection and reduced female reproduction below ground.

Future Prospects

Exploitation of endophyte-plant interactions can result in the promotion of plant health and can play a significant role in low-input sustainable agriculture applications for both food and non-food crops. An understanding of the mechanisms enabling these endophytic bacteria to interact with plants will be essential to fully achieve the biotechnological potential of efficient plant-bacterial partnerships for a range of applications. One promising area of research for future studies is developing endophytes (and rhizobacteria) to promote the sustainable production of biomass and bioenergy crops in conjunction with phytoremediation of soil contamination. The role of microbial endophytes protecting plants against insects has been well documented in various countries. However, the research is based mainly on endophytes in few host plants. Apart from isolated studies, very recently, efforts are being directed to recognize the role of endophytes in insect control. This is a broad field of investigation that is almost entirely open to new findings. The future use of biological control in combination with endophytes along with commercial pesticides applied to the seed or seedling could lead to synergistic effects on one or multiple pests and disease causing agents. IPM involving endophytes reduces costs and environmental impact, while allowing the biological agent to build up momentum for insect pest control.

Bibliography

- Bacon CW and JF White. "Microbial endophytes". Marcel Dekker, New York (2000).
- Kusari S, *et al.* "Metabolomics of endophytic fungi producing associated plant secondary metabolites: progress, challenges and opportunities". U. Roessner (Ed.), Metabolomics, In-Tech, Rijeka, Croatia (2012): 241-266.
- Ahlholm JU, *et al.* "Environmental conditions and host genotype direct genetic diversity of *Venturia ditricha*, a fungal endophyte of birch trees". *Evolution* 56.8 (2002): 1566-1573.
- Kogel KH, *et al.* "Endophyte or parasite--what decides?" *Current Opinion in Plant Biology* 9.4 (2006): 358-363.
- Staniek A, *et al.* "Endophytes: exploiting biodiversity for the improvement of natural product-based drug discovery". *Journal of Plant Interactions* 3.2 (2008): 75-93.
- Guerin P. "Sur la presence d'un champignon dans l'ivraie". *Journal of Botanique* 12 (1898): 230-238.
- de Bary A. "Morphologie and Physiologie der Pilze, Flechten, and Myxomyceten". Hofmeister's Handbook of Physiological Botany, Volume II (Leipzig, Germany: Engelmann) (1866).
- Strobel G and Daisy B. "Bioprospecting for Microbial Endophytes and Their Natural Products". *Microbiology and Molecular Biology Reviews* 67.4 (2003): 491-502.
- Deshpande MV. "Mycopesticide production by fermentation: Potential and challenges". *Critical Reviews in Microbiology* 25.3 (1999): 229-243.
- Berg G, *et al.* "Endophytic and ecdophytic potato associated bacterial communities differ in structure and antagonistic function against plant pathogenic fungi". *FEMS Microbiology Ecology* 51 (2005): 215-229.
- Berg T, *et al.* "PCR-based detection of *Xanthomonas campestris* pathovars in Brassica seed". *Plant Pathology* 54.3 (2005): 416-427.
- Lodewyckx C, *et al.* "Isolation, characterization and identification of bacteria associated to the zinc hyperaccumulator *Thlaspi caerulescens* subsp. *Calaminaria*". *International Journal of Phytoremediation* 4.2 (2002): 101-115.
- Clay K. "Fungal endophytes of plants: biological and chemical diversity". *Natural Toxins* 1.3 (1992): 147-149.
- Leuchtman A. "Systematics, distribution, and host specificity of grass endophytes". *Natural Toxins* 1.3 (1992): 150-162.
- Clement SL, *et al.* "Controlling sap-sucking insect pests with recombinant endophytes expressing plant lectin". *Nature Proceedings* (1994).
- Clement SL, *et al.* "Detrimental and neutral effects of wild barley-*Neotyphodium* fungal endophyte associations on insect survival". *Entomologia Experimentalis et Applicata* 114.2 (2005): 119-125.
- Cherry AJ, *et al.* "Pathogen incidence and their potential as microbial control agents in IPM of maize stem borers in West Africa". *BioControl* 44.3 (1999): 301-327.
- Selosse MA, *et al.* "Symbiotic microorganisms, a key for ecological success and protection of plants". *Comptes rendus Biologies* 327.7 (2004): 639-648.
- Arnold AE and Lewis LC. "Ecology and evolution of fungal endophytes, and their roles against insects". In: Vega, F.E., Blackwell, M. (Eds.), *Insect-Fungal Associations: Ecology and Evolution*. Oxford University Press, New York (2005): 74-96.
- Jones KD. "Aspects of the biology and biological control of the European corn borer in North Carolina". Ph.D. thesis, Department of Entomology, North Carolina State University (1994).

21. Leckie BM. "Effects of *Beauveria bassiana* mycelia and metabolites incorporated into synthetic diet and fed to larval *Helicoverpa zea*, and detection of endophytic *Beauveria bassiana* in tomato plants using PCR and ITS". M.S. thesis, Department of Entomology, The University of Tennessee (2002).
22. Ownley BH., *et al.* "Beauveria bassiana, a dual purpose bio-control organism, with activity against insect pests and plant pathogens". In: Lartey, R.T., Cesar, A.J. (Eds.), *Emerging Concepts in Plant Health Management*. Research Signpost, India (2004): 255-269.
23. Kambrekar DN., *et al.* "Isolation, identification and pathogenicity of microbial endophytes in different plants in northern Karnataka". *Proceeding of 9th Annual KSTA Conference on Science, Technology and Innovations in the 21st Century* (2016): 66.
24. Evans HC., *et al.* "Endophytes and mycoparasites associated with an indigenous forest tree, *Theobroma gileri*, in Ecuador and a preliminary assessment of their potential as biocontrol agents of cocoa diseases". *Mycological Progress* 2.2 (2003): 149-160.
25. Bills GF and Polishook JD. "Microfungi from *Carpinus caroliniana*". *Canadian Journal of Botany* 69.7 (1991): 1477-1482.
26. Ganley RJ and Newcombe G. "Fungal endophytes in seeds and needles of *Pinus monticola*". *Mycological Research* 110.3 (2005): 318-327.
27. Quesada-Moraga EA., *et al.* "Laboratory evaluation of entomopathogenic fungi *Baeuveria bassiana* and *Metarhizium anisopliae* against pupal and adults of *Ceratitis capitata* (Dipter: Tephritidae)". *Journal of Economic Entomology* 99.6 (2006): 1955-1966.
28. Go´mez-Vidal S., *et al.* "Endophytic colonization of date palm (*Phoenix dactylifera* L.) leaves by entomopathogenic fungi". *Micron* 37.7 (2006): 624-632.
29. Akello JT., *et al.* "*Beauveria bassiana* (Balsamo) Vuillemin as an endophyte in tissue culture banana (*Musa* spp.)". *Journal of Invertebrate Pathology* 96.1 (2007): 34-42.
30. Posada F., *et al.* "Inoculation of coffee plants with the fungal entomopathogen *Beauveria bassiana* (Ascomycota: Hypocreales)". *Mycological Research* 111.6 (2007): 749-758.
31. Cao LX., *et al.* "Endophytic fungi from *Musa acuminata* leaves and roots in South China". *Journal of Microbiology and Biotechnology* 18.2 (2002): 169-171.
32. Tian XL., *et al.* "Study on the communities of entophytic fungi and endophytic actinomycetes from rice and their antipathogenic activities in vitro". *World Journal of Microbiology and Biotechnology* 20.3 (2004): 303-309.
33. Lutz MP., *et al.* "Signaling between bacterial and fungal bio-control agent in a strain mixture". *FEMS Microbiology Ecology* 48.3 (2004): 447-455.
34. Akello J., *et al.* "Effect of endophytic *Beauveria bassiana* on populations of the banana weevil, *Cosmopolites sordidus*, and their damage in tissue cultured banana plants". *Entomologia Experimentalis et Applicata* 129.2 (2008): 157-165.
35. Arab YA and Deeb HM. "The use of endophyte *Beauveria bassiana* for bio-protection of date palm seedlings against red palm weevil and *Rhizoctonia* root-rot disease". *Scientific Journal of King Faisal University (Basic and Applied Sciences)* 13.2 (2012): 91-101.
36. Zhou W., *et al.* "The fungal endophyte *Chaetomium globosum* negatively affects both above and below-ground herbivores in cotton". *FEMS Microbiology and Ecology* 92.10 (2016): 1-22.

Volume 1 Issue 4 October 2017

© All rights are reserved by Anjitha George.