



Emission of N₂O Gas from Arable Lands: Relation with Soil pH

Muhammad Shaaban*

College of Resources and Environment, Huazhong Agricultural University, Wuhan, China

***Corresponding Author:** Muhammad Shaaban, College of Resources and Environment, Huazhong Agricultural University, Wuhan, China.

Received: January 10, 2018; **Published:** January 11, 2018

Climate change is a threat to the sustainable development, and changes in the climate have been noticed since the early 1950s. Atmospheric greenhouse gas (GHG) concentrations have increased significantly in the last three decades. Soils and fossil fuels are the main sources of GHGs in the atmosphere and nitrous oxide (N₂O) is one of the most important greenhouse gases since it has long lifetime (about 114 yrs), high global warming potential and ability to destroy ozone (O₃) layer.

Agricultural activities, including the extensive use of organic and inorganic nitrogen (N) fertilizers, have contributed to the increased emissions of N₂O to the atmosphere. Application of N fertilizers to soils provides surplus ammonium (NH₄⁺-N) and nitrate (NO₃⁻-N) which can undergo transformation via microbial nitrification and denitrification processes leading to N₂O emissions. The excessive use of N fertilizers also generates H⁺ concentrations creating soil acidification. Annually, about 20 to 33 kmol H⁺ ha⁻¹ is generated in intensive cropping areas through the excessive use of N-based fertilizers. Agricultural soils are becoming acidic worldwide in intensive farming systems due to high application rates of N fertilizers. Approximately, 30% of ice-free land worldwide is under the influence of acidification. Soil acidification is a major problem for crop growth and different techniques are recommended to counteract the problem of soil acidification, including acid tolerant crop varieties, balanced and appropriate use of fertilizers, and the application of lime and dolomite.

Application of lime is recommended in agricultural soils for amelioration of acidity. Increasing soil pH following lime application can accelerate the nitrification process and reduces the chance of NH₄⁺-N uptake by plants since many plants prefer NH₄⁺-N to NO₃⁻-N. Applications of lime can also decrease N₂O emissions through a variety of mechanisms in soil. Recently, it has been illustrated that N₂O emissions are decreased at higher pH levels by enhancement of N₂O reductase enzyme following lime application. Application of lime in acidic soils is presenting tremendous results of decreasing N₂O emissions from arable lands of low pH and attracting scientists and researchers for its advancement at global level in order to minimize the adverse effects of climate change and global warming [1-4].

Bibliography

1. Guo J., *et al.* "Significant acidification in major Chinese croplands". *Science* 327.5968 (2010): 1008-1010.
2. Qu Z., *et al.* "Excessive use of nitrogen in Chinese agriculture results in high N₂O/(N₂O+ N₂) product ratio of denitrification, primarily due to acidification of the soils". *Global Change Biology* 20.5 (2013): 1685-1698.
3. Shaaban M., *et al.* "Nitrous oxide emission from two acidic soils as affected by dolomite application". *Soil Research* 52 (2014): 841-848.
4. Shaaban M., *et al.* "Effects of dicyandiamide and dolomite application on N₂O emission from an acidic soil". *Environmental Science and Pollution Research* 23.7 (2016): 6334-6342.

Volume 2 Issue 2 February 2018

© All rights are reserved by Muhammad Shaaban.