



Effects of Groundnut Shell, Rice Husk and Rice Straw on the Productivity of Maize (*Zea mays* L.) and Soil Fertility in the Guinea Savannah Zone of Ghana

Israel K Dzomeku* and Osman Illiasu

Department of Agronomy, University for Development Studies, Tamale, Ghana

*Corresponding Author: Israel K Dzomeku, Associate Professor and Dean, Department of Agronomy, University for Development Studies, Tamale, Ghana. E-mail: idzomeku@uds.edu.gh

Received: December 30, 2017; Published: February 09, 2018

Abstract

A field experiment was conducted at Nyankpala, near Tamale during the 2014 cropping season to investigate the effects of indigenous organic materials (groundnut shell, rice husk and rice straw) on yield components and yield of maize. The study was a $3 \times 3 \times 3$ factorial experiment consisting of the three organic materials at three levels (2.5, 5 and 7.5 t ha⁻¹ on dry matter basis) and three NPK rate (zero control, 45-30-30 kg/ha and 90-60-60 kg/ha) laid out in a randomized complete block design with four replications. Best grain yield of 4781 kg/ha was obtained with 7.5 t/ha groundnut shell plus 90-60-60 kg NPK/ha but 5 t/ha of groundnut shell plus 90-60-60 kg NPK/ha, 7.5 t/ha groundnut shell plus 45-30-30 kg NPK/ha, 7.5 t/ha of rice husk or rice straw plus 90-60-60 kg NPK/ha gave equal yields. Longest cob was obtained with 5 to 7.5 t/ha groundnut shell (18.9 cm) or 7.5 t/ha rice husk (17.9 cm) or 7.5 t/ha rice straw (17.9 cm), each plus at least 45-30-30 kg NPK/ha. However, sole application of 5 to 7.5 t/ha groundnut shell notably determined cob weight (116.4 g/cob), 100 seed weight (24.9g) and stover weight (4822 kg/ha). Similarly, sole application of NPK at 45-30-30 kg/ha was adequate to enhance cob length (16.8 cm), cob weight (127.4 g/cob) and stover weight (4514 kg/ha). Grain yield correlated positively with stover weight ($r = 0.851^{**}$) and cob length ($r = 0.601^{**}$).

Keywords: Soil Fertility; Organic Matter; Groundnut Shell; Rice Husk; Rice Straw

Introduction

Maize (*Zea mays* L.) is one of the major important cereal crops that provide staple food to a large number of the rapidly growing human population in the world especially sub-Saharan Africa. It is a rich source of food for human population, fodder for animals (especially ruminants), and feed for poultry and provides raw material for industrial use. Maize has a carbohydrate content of about 71%, but low in protein. It is processed into a wide range of foods and beverages and consumed as breakfast, main meals, or snacks.

Despite the importance of maize as a food security crop, many factors militate against attainment of its grain yield potential in Ghana. Low soil fertility has been identified as a major factor reducing crop yields in Ghana [1]. A quantum of literature is available on a similar cereal crop rice, to prove low inherent soil fertility as a major cause of poor crop productivity [2,3]. Nyalemegbe., *et al.* [4] attributed the low inherent soil fertility partly to low levels of soil organic matter and observed that rice yields in continuously cropped fields declined with time, even with the application of mineral fertilizers.

Issaka., *et al.* [5] reported abundance of indigenous organic materials available in Ghana that could be used to improve soil fertility. JIRCAS [6] also reported on the quantity, quality and distribution of various organic materials in Ghana. According to this

report, rice straw, cow dung and human excreta were evaluated in Northern region as chiefly effective materials applied at 3 t/ha, could improve soil fertility and soil physical properties and its buffer capacity. Samy., *et al.* [7] observed noted good potential for organic based farming with a combination of organic fertilizers to attain maximum yields. There is growing interest of plant materials grown in situ to provide litter fall, root decay, green manure incorporation, crop residue returns and animal excreta and its subsequent decomposition to enhance soil fertility [8]. Research to date has predominantly focused on inorganic fertilizers and organic materials from animal waste, solid waste and compost, with least attention on indigenous raw materials. The use of indigenous organic materials (such as rice straw, rice husk, and grounded groundnut shell) as soil amendments for agricultural practices is limited in Ghana. The objectives of the study were to determine the best promising indigenous organic materials for maize production in Northern Ghana using rice straw, rice husk and groundnut shell with and without inorganic NPK fertilizer on yield components and grain yield of maize.

Materials and Methods

Study Site Description

The experiment was conducted at the University for Development Studies at Nyankpala, near Tamale, in the Guinea savannah zone of Ghana, at latitude 9° 25'14"N, longitude 0° 58' 42"W and

at an altitude of 183 m above sea level [9]. The area experiences unimodal rainfall with an annual mean rainfall of 1000 to 1022 mm. Temperature distribution is fairly uniform with mean monthly minimum of 21.9°C and a maximum of 34.1°C. It has a minimum relative humidity of 46% and maximum of 76.8%. The soil of the study site is a typical upland soil, developed from iron stone gravel and ferruginized ironstone brash [10]. The soil is classified as a Haplic Lixisol [11] and locally referred to as the Tingoli series [12].

Experimental Design and Treatments

The experiment was a 3 × 3 × 3 factorial experiment made up of four indigenous organic materials (groundnut shell, rice husk and rice straw) at three levels (2.5, 5 and 7.5 t ha⁻¹ dry matter basis) and three NPK rate (0-0-0 kg NPK ha⁻¹, 45-30-30 kg NPK ha⁻¹ and 90-60-60 kg NPK ha⁻¹), laid in a randomized complete block design with four (4) replications. Plots measured 5m × 5m with 1m and 2m alleys between plots replications respectively.

Agronomic Practices

The experimental field was tractor ploughed and two weeks after harrowed and laid-out after which the organic materials were applied and manually incorporated into the soil with a hoe. The experimental field was left undisturbed for 21 days before planting. Planting of ‘Wang Data’ maize variety was done by sowing two seeds per hill at a spacing of 80 cm x 45 cm. NPK 15-15-15 was applied at 0-0-0 kg NPK ha⁻¹, 45-30-30 kg NPK ha⁻¹ and 90-60-60 kg NPK ha⁻¹. Atrazine (80 WP) was applied seven days after planting (DAP) and supplemented with hoe weeding at 35 DAP and 56 DAP.

Soil Analysis

Before planting, three soil samples were randomly collected at 0 - 20 cm soil depth from each replicate by using an auger and the samples bulked together to get one composite sample to determine the initial soil physio-chemical properties (Table 1). After harvesting soil samples were collected in the same manner, as before planting, but replicates of each treatment were mixed with each other and analysis was done with samples for each treatment (Table 2), with the results being prepared for a second paper.

Table 1: Initial Soil Analysis, July, 2014.

Initial soil	pH	% OC	% N	Mg/kg P	Mg/kg K	Mg/kg Ca	Mg/kg Mg
	5.5	0.12	0.01	3.6	51.8	64.7	27.9

Table 2: Soil Chemical Analysis after Harvesting, November, 2014.

Treatment	Soil pH	% OC	% N	Mg/Kg P	Mg/Kg K	Mg/Kg Ca	Mg/Kg Mg
2.5 t/ha Groundnut shells	5.5	0.8	0.1	8.9	98.5	64.7	47.3
2.5 t/ha Groundnut shells + 45-30-30 kg/ha NPK	5.4	1.7	0.2	12.8	61.3	186.8	65.6
2.5 t/ha Groundnut shells + 90-60-60 kg/ha NPK	5.0	0.8	0.07	8.4	52.7	122.6	47.8
5 t/ha Groundnut shells	5.1	2.9	0.3	8.7	55.0	178.9	72.5
5 t/ha Groundnut shells + 45-30-30 kg/ha NPK	4.5	3.0	0.3	10.6	58.6	164.7	88.7
5 t/ha Groundnut shells + 90-60-60 kg/ha NPK	4.8	3.2	0.3	14.9	59.4	188.6	97.9
7.5 t/ha Groundnut shells	5.1	2.8	0.3	8.0	151.8	245.8	87.6
7.5 t/ha Groundnut shells + 45-30-30 kg/ha NPK	5.1	2.8	0.3	6.8	114.3	214.4	71.9
7.5 t/ha Groundnut shells + 90-60-60 kg/ha NPK	5.0	3.0	0.3	8.7	198.5	267.0	92.7
2.5 t/ha Rice Husk	4.7	0.8	0.04	3.8	74.8	98.7	58.4
2.5 t/ha Rice Husk + 45-30-30 kg/ha NPK	4.6	1.0	0.1	5.5	74.0	136.9	64.3
2.5 t/ha Rice Husk + 90-60-60 kg/ha NPK	5.1	0.8	0.1	5.8	75.0	118.5	72.2
5 t/ha Rice Husk	4.7	0.8	0.1	3.9	72.3	83.2	52.1
5 t/ha Rice Husk + 45-30-30 kg/ha NPK	4.8	1.0	0.1	5.7	68.8	96.9	81.2
5 t/ha Rice Husk + 90-60-60 kg/ha NPK	5.1	0.8	0.1	4.9	84.4	114.8	52.5
7.5 t/ha Rice Husk	4.6	0.8	0.1	5.7	76.6	92.7	57.2
7.5 t/ha Rice Husk + 45-30-30 kg/ha NPK	4.9	0.8	0.1	5.5	86.4	124.2	48.8
7.5 t/ha Rice Husk + 90-60-60 kg/ha NPK	5.1	1.3	0.2	7.8	92.6	132.1	82.5
2.5 t/ha Rice Straw	4.9	1.9	0.2	4.5	64.4	98.3	58.6
2.5 t/ha Rice Straw + 45-30-30 kg/ha NPK	4.7	2.8	0.3	9.0	86.5	196.8	64.9
2.5 t/ha Rice Straw + 90-60-60 kg/ha NPK	4.9	2.4	0.3	8.7	73.9	175.7	74.4
5 t/ha Rice Straw	5.0	1.7	0.3	6.5	56.5	118.3	58.8
5 t/ha Rice Straw + 45-30-30 kg/ha NPK	4.9	3.3	0.3	11.6	78.6	226.4	84.2
5 t/ha Rice Straw + 90-60-60 kg/ha NPK	4.9	3.4	0.4	12.8	98.9	218.8	78.7
7.5 t/ha Rice Straw	5.1	1.9	0.3	10.0	64.3	114.3	57.0
7.5 t/ha Rice Straw + 45-30-30 kg/ha NPK	4.8	3.4	0.3	12.7	76.9	198.7	77.5
7.5 t/ha Rice Straw + 90-60-60 kg/ha NPK	4.9	3.3	0.4	11.9	112.7	228.3	65.0

Data Collection

Five plants were randomly selected per plot and tagged for the measurement of growth and yield characteristics such as plant height, leaf area index, days to 50% flowering and height of cob attachment, cob weight, cob length, stover weight, 100 seed weight and grain yield.

Statistical Analysis

The data were subjected to analyses of variance using GenStat statistical package. Treatment means were separated using Least Significant Difference at 5% significant level. Correlation and regression analysis were also examined.

Results and Discussions

Days to 50% Flowering

Organic materials supplemented with inorganic NPK fertilizer enhanced early flowering in maize with 2.5 to 7.5 t/ha of groundnut shell or rice husk or rice straw supplemented with at least 45-30-30 kg/ha NPK took the shortest days of 47 to 50 days to 50% flowering (Figure 1). Timely availability of nutrients mainly nitrogen from the organic source and NPK fertilizer could have provided adequate availability of the required crop growth conditions which positively supported the physiological functions of the crop to early flowering as reported by Khan, *et al* [13].




Figure 1: Effects of organic material by NPK fertilizer on days to 50% flowering of maize. Bars represent SEMs.

Cob Length

Combination of organic materials and NPK amended soil significantly ($p < 0.01$) increased cob length. Entries of 5 to 7.5 t/ha groundnut shell or rice husk or rice straw plus at least 45-30-30 kg/ha NPK maximise cob length (Figure 2). Lengthy cobs supported by the synergy of organic and NPK amended entries could be attributed to high growth rate attained by the crop due to timely availability of adequate nutrients from both sources of soil fertility amendments with consequential increased dry matter accumulation. In a previous study, Uzoma, *et al.* [14] reported combined application of organic and inorganic fertilizers positively affected maize ear characteristics and ascribed it to incorporation of organic material that improved soil physical properties and the increase in mineralization as a result of the addition of synthetic fertilizers. Bilalis, *et al.* [15] observed that sustainable yield index and agro-

nomical efficiency was more stable under combined organic and inorganic fertilization in maize production.

Cob Weight

Cob weight as influenced by organic materials was in the range of 118 to 127 g/cob with 5 to 7.5 t/ha groundnut shell supporting the highest weight followed by 7.5 t/ha rice straw similar to 5-7.5 t/ha of rice husk (Figure 3). Inorganic fertilizer application increased cob weight in the range of 127 to 135 g/cob maximised at 90-60-60 kg/ha NPK but similar to 45-30-30 kg/ha NPK (Figure 4). The maximum maize cob weight observed with organic matter could be attributed to improved nutrient availability to crop through enhanced organic matter decomposition and mineralization processes and directly from applied N which improved maize root and shoot development, as previously reported [15,16].




Figure 2: Effects of organic material by NPK fertilizer on cob length of maize. Bars represent SEMs.




Figure 3: Effects of the organic materials on cob weight of maize. Bars represent SEMs.




Figure 4: Effects of NPK fertilizer on cob weight of maize. Bars represent SEMs.

Grain Yield

Integration of organic material and NPK fertilizer significantly ($p < 0.001$) increased the parameter with maximum grain yield of maize (4781 kg/ha) at 7.5 t/ha groundnut shell plus 90-60-60 kg/ha NPK (Figure 5). However, four other entries of 5 t/ha of groundnut shell plus 90-60-60 kg/ha NPK, 7.5 t/ha groundnut shell plus 45-30-30 kg/ha, 7.5 t/ha of either rice husk or rice straw plus 90-60-60 kg/ha NPK gave similar grain yields. Combined organic and nitrogen fertilization promoted maize yield in comparison to the organic materials alone and agreed with findings of Sadeghi and Bahrani [17] who observed optimum crop growth with the highest crop residues and nitrogen. The results could be attributed to the overall improvement in soil chemical, physical and biological properties [18]. The observations confirm findings of Asai, et al. [19] who observed that integrated nitrogen strategies convincingly enhance maize yield. Continuous application of smaller amount of manure plus fertilizer increased grain yield and sustained soil fertility in the sub-humid zone of Ghana and was noted to reduce nutrient mining and environmental degradation [16].

100 Seed Weight

Hundred seed weight varied due to the effects of applied organic materials ($p < 0.001$) and mineral nitrogen ($p < 0.05$), with application of 7.5 t/ha groundnut shells supporting highest size of 27.7g (Figure 6). Whilst 90-60-60 kg NPK/ha optimised seed weight (28.6g), 45-30-30 kg NPK/ha gave similar size (Figure 7). Khan, et al. [13] observed that lower nitrogen level in the soil resulted in lighter seeds. Similar to our results maize yield and yield components was reported to increase with organic matter application [20].




Figure 5: The synergistic effect of organic materials and NPK fertilizer on grain yield of maize after harvesting during the 2014 cropping season. Bars represent SEMs.




Figure 6: Effects of the organic materials on 100 seed weight of maize. Bars represent SEMs.




Figure 7: Effects of NPK fertilizer on 100 seed weight of maize. Bars represent SEMs.

Stover Weight

Stover weight was both highly significantly determined by effects of organic materials and mineral nitrogen. The highest stover weight obtained with the applications of 5 t/ha - 7.5 t/ha groundnut shell and 7.5 t/ha rice straw (Figure 8) could be due to continues slow release and adequate availability of crop nutrients from the organic materials buried in the soil, which were less subjected to leaching losses. NPK fertilizer application at 90-60-60 kg/ha maximized stover weight of 4792 kg/ha but was similar to 45-30-30 kg NPK/ha (Figure 9). It was reported organic amendments positively increased crop growth and net assimilation rates with consequential high maize productivity [14].




Figure 8: Effects of the organic materials on stover weight of maize. Bars represent SEMs.




Figure 9: Effects of NPK fertilizer on stover weight of maize. Bars represent SEMs.

Correlation Analysis

Stover weight highly correlated ($r = 0.851$) with grain yield followed by cob length ($r = 0.601$). This indicates that grain yield at harvest was determined by the length of the cob and stover weight (Table 3).

Table 3: Correlation between cob length, cob weight, stover Weight, 100 seed weight and grain yield.

	Grain Yield	Cob Length	Cob Weight	100 Seed Weight	Stover Weight
Grain Yield	1.000				
Cob Length	0.601	1.000			
Cob Weight	0.313	0.173	1.000		
100 Seed Weight	0.499	0.091	0.323	1.000	
Stover Weight	0.851	0.733	0.149	0.369	1.000

Soil Analysis after Harvesting

Soil analysis after harvesting showed treatments had affected the chemical properties of the soil over the initial. The soil amendments slightly affected the pH of the soil. The soil amendments also positively increased the organic carbon content and the major plant nutrient elements (nitrogen, phosphorus, potassium, calcium and magnesium) of the soil (Table 2). Masulili, *et al.* [21] reported on the improvement of soil properties with organic soil amendments. Saha, *et al.* [22] reported that rice straw incorporation increased soil organic matter content. Notably, Tualar, *et al.* [23] observed that the interaction between the application of organic fertilizers and inorganic fertilizers (nitrogen, phosphorus and potassium) affected organic carbon content and soil cation exchange capacity.

Conclusion

The results showed integrated use of organic amendments of groundnut shells, rice husk and rice straw with NPK fertilizer at 45-30-30 kg/ha increased maize grain and yield components.

Grain yield of maize was maximised by 7.5 t/ha groundnut shell plus 90-60-60 kg/ha NPK, being 4781 kg/ha. However, the lower amount of organic matter required in 5 t/ha of groundnut shell plus 90-60-60 kg/ha NPK, and lower input of NPK in 7.5 t/ha groundnut shell plus 45-30-30 kg/ha which equalled the maximum could be preferred by farmers who can only afford lower quantities of organic material or NPK. Application of 7.5 t/ha of either rice husk or rice straw plus 90-60-60 kg/ha NPK also performed similar to the maximum in grain yield. The application of 2.5 to 7.5 t/ha of groundnut shell or rice husk or rice straw plus at least 45-30-30 kg NPK/ha enhanced days to 50% flowering and cob length. Generally 5 to 7.5 t/ha of groundnut shell alone were outstanding in enhancing cob weight, 100 seed weight and stover weight. The application of NPK fertilizer alone at 45-30-30 kg NPK/ha promoted cob weight, 100 seed weight and stover weight. Overall implication showed combination of organic matter and inorganic NPK fertilizer could play a more significant role in enhancing and replenishing soil nutrients and sustained maize production.

Bibliography

1. Issaka RN, *et al.* "Characterization of Indigenous Resources in Ghana: Locally Available Soil Amendments and Fertilizers for Improving and Sustaining Rice Production". CSIR-SRI/CR/RNI/2010/01 (2011).
2. Senayah JK, *et al.* "Characteristics of Major Lowland Rice-Growing Soils in the Guinea Savannah Voltaian Basin of Ghana". *Agriculture and Food Science Journal of Ghana* 7 (2008): 445-458.
3. Abe S., *et al.* "Soil Fertility Potential for Rice Production in West African Lowlands". *Japan Agricultural Research Quarterly* 44.4 (2010): 343-355.
4. Nyalemegbe KK, *et al.* "Integrated Organic-Inorganic Fertilizer Management for Rice Production on the Vertisols of the Accra Plains of Ghana, West Africa". *West African Journal of Applied Ecology* 16.1 (2009): 23-31.
5. Issaka RN, *et al.* "Indigenous Fertilizing Materials to Enhance Soil Productivity". In Ghana. Dr. Joann Whalen (Ed); Soil Fertility Improvement and Integrated Nutrient Management – A Global Perspective (2012).
6. JIRCAS. "The Study of Improvement of Soil Fertility with Use of Indigenous Resources in Rice Systems of Sub-Sahara Africa". Business Report (2010).

7. Samy J., *et al.* "Organic Rice Farming System (Studies on the Effect of Organic Matter on Rice Yield, Soil Properties and Environment)". A Research Project of Perez-Guerrero Trust Fund (PGTF) for Economic and Technical Cooperation among Developing Countries, Members of the Group of 77. Strategic Environment and Natural Resources Research and Development Institute (NARDI) (1997).
8. Sanchez PA., *et al.* "Replenishing Soil Fertility in Africa". Sanchez, *et al.* (eds) Proceeding of an International Symposium cosponsored by Division A-6 (International Agronomy) and S-4 (Soil Fertility and Plant Nutrition) and the International Center for Research in Agroforestry, held at the 88th Annual Meeting of the American Society of Agronomy and Soil Science Society of America, Indianapolis, Indiana, SSSA Special Publication No. 51 (1996).
9. NAES. "Nyankpala Agricultural Experimental Station". Nyankpala, Ghana. Annual Report (1992).
10. Adu SV. "Report on the Detailed Soil Survey of the Central Agricultural Station, Nyankpala". Soil Research Institute, Kumasi, Ghana (1957).
11. FAO/UNESCO. "Soil Map of the World: Revised Legend". FAO, Rome (1997): 119.
12. Serno G and Van de Weg RF. "Preliminary Assessment of the (Available) Existing Soil Information of Nyankpala Agricultural Experimental Station, Tamale, Ghana". Stiboka, Wageningen, the Netherlands (1985).
13. Khan HZ., *et al.* "Effect of Rate and Source of Organic Material on the Production Potential of Spring Maize (*Zea mays* L.)". *Pakistan Journal of Agricultural Science* 45.1 (2008): 40-43.
14. Uzoma KC., *et al.* "Effect of Cow Manure Biochar on Maize Productivity under Sandy Soil Condition". *Soil Use and Management* 27.2 (2011): 205-212.
15. Bilalis Dimitrios., *et al.* "Combined Organic/Inorganic Fertilization Enhances Soil Quality and Increased Yield, Photosynthesis and Sustainability of Sweet Maize Crop". *Australian Journal of Crop Science* 4.9 (2010): 722-729.
16. Abunyewa AA., *et al.* "Integrated Manure and Fertilizer Use, Maize Production and Sustainable Soil Fertility in Sub Humid Zone of West Africa". *Journal of Agronomy* 6.2 (2007): 302-309.
17. Sadeghi H and Bahrani MJ. "Effects of Crop Residue a Nitrogen Rates on Yield and Yield Components of Two Dry land Wheat (*Triticum aestivum* L.) Cultivars". *Plant Production Science* 12.4 (2009): 497-502.
18. Sahoo D., *et al.* "Effect of Twenty-Five Years of Fertilizer Application on Productivity of Rice-Rice System". Swarup A., Reddy DD and Prasad RN. (Eds.) In: Long-term Soil Fertility Management through Integrated Plant Nutrient Suly. Indian Institute of Soil Science Bhopa, India (1998): 229-237.
19. Asai H., *et al.* "Biochar Amendment Techniques for Upland Rice Production in Northern Lagos 1. Soil Physical Properties, Leaf SPAD and Grain Yield". *Field Crops Research* 111 (2009): 81-84.
20. Steiner C., *et al.* "Long Term Effects of Manure, Charcoal and Mineral Fertilization on Crop Production and Fertility on a Highly Weathered Central Amazonian Upland Soil". *Journal of Plant and Soil* 292.1-2 (2007): 275-290.
21. Masulili A., *et al.* "Rice Husk Biochar for Rice Based Croing System in Acid Soil 1. The Characteristics of Rice Husk Biochar and Its Influence on the Properties of Acid Sulfate Soils and Rice Growth in West Kalimantan, Indonesia". *Journal of Agriculture Science* 2.1 (2010): 39-47.
22. Saha PK., *et al.* "Long-Term Integrated Nutrient Management for Rice-Based Croing Pattern: Effect on Growth, Yield, Nutrient Uptake, Nutrient Balance Sheet and Soil Fertility". *Communications in Soil Science and Plant Analysis* 38.5-6 (2007): 579-610.
23. Tualar Simarmata., *et al.* "Alcation of Straw Compost and Biofertilizers to Remediate the Soils Health and to Increase the Productivity of Paddy Rice in Indonesia". Conference on International Research on Food Security, Natural Resource Management and Rural Development organised by: Georg-August Universität Göttingen and University of Kassel-Witzenhausen. Tropentag, Göttingen, Germany (2012).

Volume 2 Issue 3 March 2018

© All rights are reserved by Israel K Dzomeku and Osman Illiasu.