



Linking Climate Change with Global Phenology

Sanjay Kumar^{1*} and Neha Chopra²

¹Research Officer, Department of Botany, D.S.B. Campus, Kumaun University, Nainital, India

²Research Scholar, Department of Botany, D.S.B. Campus, Kumaun University, Nainital, India

*Corresponding Author: Sanjay Kumar, Research Officer, Department of Botany, D.S.B. Campus, Kumaun University, Nainital, India.

Received: January 31, 2018; Published: March 22, 2018

Abstract

At present, phenological studies achieved a prominent position in monitoring and predicting the timing of recurrent life cycle events. Phenology also provide a better understanding to manage and conserve environment. This article provides a useful information how phenology, a multidisciplinary science plays an important role in conservational biology. We tried to explain changes in plant phenology caused by rising temperature, their impacts on species diversity and plant-animal interactions, and how conservation efforts could be improving in relation to plant resource organization. We listed the effects of phenological shifts and examine how phenological research can contribute to estimate and manage the land-use change and other natural and anthropogenic disturbances, such as fire, exotic and invasive species. Various instruments used for monitoring phenological activities at species-rich area are also discussed here.

Keywords: Phenology; Conservation; Pollinator; Frugivorous; Land-Use Pattern

Introduction

Phenology explores how the plants and animals will respond according to seasonal changes. In varying environmental conditions, the effect of rising temperature on plant phenophase [1] and animal [2] is extensively studied to know the variation in ecosystem dynamics resulting from a mismatch in the timing of interdependent phenophases [3]. Climate also affects the timing of flower colour and carbon flux of atmosphere [4]. Phenology is the best indicator of climate change as it is direct link between temperature and plant and animal development. The average global temperature has risen by 0.85°C which is also be proven by several number of phenological studies. This average global temperature is predicted to continue to rise and is predicted to continue to rise [5].

The phenology of many organisms is also affected by anthropogenic effects. Several studies showed [6] that the rising temperature change the species phenology earlier by several days to weeks. These studies are supported by experimental studies which link rising temperature to phenological shifts [7]. However, despite of rising temperature, plant phenology is also affected by day length, rainfall, nutrient availability and timing of snow melt [8] which are influenced by human activity. Abiotic ecosystem processes viz., soil moisture and nutrient pool influenced by plant species diversity [9]. These experiments were conducted on a smaller scale so the magnitude of nitrogen deposition and changing precipitation pat-

terns was untested. Nitrogen deposition and changed precipitation regimes have also changed flowering time [10]. Rather than temperature, disturbances in species composition, abundance, or diversity also affect phenology which are influenced by human activities [11]. The role of biotic interactions on leafing, flowering and other phenological events is critical to understand.

Long-term phenological records of Northern Hemisphere shows a close relationship between rising temperature and earlier onset of leafing and flowering [1,12]. The conservation and management of natural systems can be improve with greater knowledge of the causes regulating and controlling plant cycles and differences across species, populations and communities [13]. Various vegetation monitoring techniques viz., digital photographs, satellite-derived phenology [4,14,15] can be applied worldwide to know the temporal shifts in vegetation.

Phenology helps in biotic conservation

Relation between leafing and herbivory

Study of leafing phenology, directly linked to ecosystem processes is very useful for conservation [13], understanding plant-water relations, primary productivity in terrestrial ecosystems, as well as gas exchange rates, biogeochemical cycling, dynamics of carbon sequestration [13] and defining the length of growing seasons and seasonal patterns of photosynthesis at local to global

scales [15]. So leaf phenology provides key information for ecosystem process models that forecast responses to land-use change, atmospheric chemistry, and climate [15]. Any shifts on onset and end of growing season due to climate change may have consequences on net primary production. Premature leaf senescence in deciduous forests may be induced by increased temperature and frequent drought, affecting the efficiency of nutrient resorption and length of growing seasons and impacts carbon uptake and ecosystem nutrient cycling [16], and therefore management practices [17]. Climate change, land-use change or use of insecticides shifts herbivorous insect phenology which in turn threatens plant population viability leading to increases in herbivore damage [18]. To avoid damage from insects, plants adopt many phenological strategies such as producing large, synchronous pulses of leaves to satiate herbivores and the timing of leafing peaks to the season with the lowest insect densities [19].

In agriculture, mismatch timing between highest insect density and their natural enemies due to climate change also decreases the effectiveness of biocontrol measures [20]. So it is necessary to have a detailed knowledge of phenological dynamics of folivorous animals and their host/target plants to conserve and manage both herbivores and plant populations, and when designing pest control programmes in natural and agricultural ecosystems [15].

Relation between flowering and pollinator

In tropics, delimited flowering seasonality during springtime, typical of temperate and boreal ecosystems is generally absent. In tropics, animal-pollinated species has been estimated at 94% [21] where flowers with their aromatic compounds attract females providing food for pollination services enabled strong and diverse adaptations in flower visitors, maintaining rich assemblages of highly specialized floral foragers, such as bees and humming birds.

Relation between fruiting and frugivorous

Frugivorous animals, affects seed dispersal and germination effectiveness [22] critically rely on fruits for their diet, population size, social behaviour, reproduction, and movements - depend on fruit abundance and seasonality [23]. Neotropical plant species contains high percentage of fruits which dispersed by frugivorous animals [24] whose population controlled by low fruit production or changes in fruit supply over time according to their nutritional content, morphology and colour [25], with consequences for their conservation and management [26]. Lower fruit availability affects vertebrate frugivores. Hunting for valuable animals also affects dispersal mechanism which in turn also affects seedling establishment [27]. This is especially serious for large-seeded plant species who are dependent on these animals for their seed dispersal [28] e.g. defaunation of large-gaped frugivorous birds has been singled out as the main cause of rapid evolutionary change in palm seed size [27].

To conserve tropical communities, it is necessary to understand interconnection between accessibility of resources for primary consumers and seasonal variations in climate [29].

Shifts in phenological activity and species interaction

Effects of temperature on plant phenology is not same for all species. It changes the duration of reproductive phase and timing of phenological activity at community level [28,30]. Any fluctuation in phenological activity in some plant species can also affect other plants through competition and/or facilitation for pollinators and seed dispersers [31]. Some studies show that phenology plays a crucial role in the stability and diversity of mutualistic communities [32] which is important for the conservation and management of plant-pollination interactions and mutualistic networks [33]. In this context, one potential threat from climate change is the temporal uncoupling of mutualistic species interactions [30,33]. Fluctuations in global temperature cause early/delay in flowering and fruiting time which in turn affects the primary consumer and seed dispersers results low recruitment and resource scarcity [33]. Ultimately, both the ability of animal partners to forage on changing host plants as well as the maintenance of viable services for host plants from these mutualistic partners will influence the severity of potential effects of phenological mismatches and the conservation of mutualistic networks [33].

Changes in phenology, climatic conditions and land-use pattern

Divisions and edge effects

Land-use changes result in habitat loss, causes declines in plant and animal diversity [34]. This activity produces variations in light, temperature and humidity conditions of that habitat. These variations affect phenological activity, plant-animal interactions and ecological services [35]. An increase in flowering and fruiting activity in natural habitats with increased sunlight, is reported in several studies [36]. However, in fragmented areas and those subjected to edge effects, this higher production in reproductive plant parts does not always favour the reproductive success and recruitment of native species from the original plant community [36]. This may be caused by new environmental conditions of fragmented area with loss of pollinators and seed dispersers [35].

Effects of fire on phenology

Fire determines vegetation physiognomy and species diversity [37]. It stimulates flowering and fruiting [38], seed germination [39], increases fruit production [40], and/or phenological cycle by shifting the flowering/fruiting date [40], but may also depress the availability of large-seeded fruits (Barlow and Peres, 2006). Regarding benefits, fire also reduces flowering and fruiting by destroying buds, flowers and fruits, affecting species that reproduce during the fire season [41] and/or favour invasive species [42].

Phenological study of fire-prone area helps to improve management practices and restoration strategies. High fire frequency results low flower and fruit production [41] results in low regeneration [43].

Phenological patterns and species interactions

The investigation of how native, exotic and invasive species cooperate could profit by recognizing plant phenology as a key attribute impacting their associations [44]. Invasive species cause decline in native species or local extinctions because they change the structure and function of native ecosystems [45]. The native and exotic species which are closely related compete for pollinators and seed dispersers, change fruit quality, quantity, dispersal of seed and so community structure and ecosystem functioning [46]. Exotic species can leaf, bloom and fruit at that time when natives were not and get benefits from a priority effect [44]. Compared to natives, exotic species have also longer leaf and fruit period which provide the advantages over natives by providing more adaptability to environmental changes with implications for management and conservation [44]. When disturbance promote biomass growth, native species act as invasive species; e.g. native liana hyper abundance resulting from increased temperature and CO₂ availability associated with global atmospheric change [47]. The trees which are liana supported therefore modified by light competition which affects vegetative and reproductive phase [48]. During scarcity of flowering trees, native lianas provide flower resources to pollinators [49]. Such area of should take into account for forest conservation and management.

Tools for observing phenological activities

Phenological databases

The phenological studies and observation in modern era not fit in current conservation strategies. Observers organized phenology results like a seed collection calendar [50]. So its required to add additional objectives like estimate relationship between fruiting phenology and seed germination, dormancy [51], and storage behaviour [52] in seasonal habitats for choosing better adaptable species, to make methods for breaking dormancy, and seed preservation. From herbaria we can reconstruct past historical patterns of plant phenology [53] which provide data on reproductive pattern of any species whose phenoligical information was is not available [54]. Now-a-days digital photographs were also used for the same purpose. Dendrochronology is also very useful to understand effect of climate change on tree growth rings and cambial activity [55]. The results are used to develop growth models and to understanding how climatic conditions affects growth and cambial activity. This information used to make better management and conservation plan [56].

Phenological monitoring

With increasing importance of climate change and global warming, phenological monitoring techniques are also developing. These techniques help in predicting upcoming effects of climate change

and address applied environmental and conservation issues [57]. Beside traditional phenological observations, this information will promote scientists to use alternative methods like remote sensing based phenology from regional to global scales [58] and deployment of in situ digital cameras for continuous monitoring of multiple simultaneous sites, referred to as near-surface remote phenology [4]. In species rich tropical areas, sampling can be expensive and labour-intensive so alternative techniques can be used for phenological studies [14]. Pehno cams can be useful tools in such area which detect changes in leafing events according to changes in the red, green and blue (RGB) channels [15]. Orbital remote sensing is also another useful which records biophysical and biochemical vegetation parameters [58]. Remote sensing detects seasonal vegetation changes over a large range of spatial and temporal scales, and have been incorporated into conservation practices [59,60].

Conclusions and Recommendations for Future Research

Now it is necessary to use new technology and tools for observing phenology, because phenology enters in a new phase which need all these techniques for long term data collection. As already discussed that new predicting models are developing to study phenological shift at different scales in current climatic scenario, these mechanisms will help us to improve models capabilities. More advance studies are required to improve conservation and management policies at species and community level. We can also address the effects of rising temperature on local and global scale with geographical variations.

Bibliography

1. Menzel A., *et al.* "European phenological response to climate change matches the warming pattern". *Global Change Biology* 12.10 (2006): 1969-1976.
2. Newson SE., *et al.* "Long-term changes in the migration phenology of UK breeding birds detected by large-scale citizen science recording schemes". *Ibis* 158.3 (2016): 481-495.
3. Visser ME and Hollerman LJM. "Warmer springs disrupt the synchrony of oak and winter moth phenology". *Proceedings of the Royal Society B: Biological Sciences* 268.1464 (2001): 289-294.
4. Richardson A D., *et al.* "Climate change, phenology, and phenological control of vegetation feedbacks to the climate system". *Agricultural and Forest Meteorology* 169 (2013): 156-173.
5. IPCC "Climate Change 2014: Impacts, Adaptation, and Vulnerability". Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2014).
6. Wolkovich EM., *et al.* "Warming experiments under predict plant phenological responses to climate change". *Nature* 485.7399 (2012): 494-497.

7. Miller-Rushing AJ, et al. "The effects of phenological mismatches on demography". *Philos Trans R Soc Lond B Biological Science* 365.1555 (2010): 3177-3186.
8. Cleland EE, et al. "Shifting plant phenology in response to global change". *Trends in Ecology and Evolution* 22.7 (2007): 357-365.
9. Cardinale BJ, et al. "The functional role of producer diversity in ecosystems". *American Journal of Botany* 98.3 (2011): 572-592.
10. Hoover SER, et al. "Warming, CO₂, and nitrogen deposition interactively affect a plant-pollinator mutualism". *Ecology Letters* 15.3 (2012): 227-234.
11. Tylianakis JM, et al. "Global change and species interactions in terrestrial ecosystems". *Ecology Letters* 11.12 (2008): 1351-1363.
12. Schwartz M D, et al. "Onset of spring starting earlier across the northern hemisphere". *Global Change Biology* 12.2 (2006): 343-351.
13. Polgar C A and Primack R B. "Leaf-out phenology of temperate woody plants: from trees to ecosystems". *New Phytologist* 191.4 (2011): 926-941.
14. Alberton B, et al. "Using phenological cameras to track the green up in a cerrado savanna and its on-the-ground validation". *Ecological Informatics* 19 (2014): 62-70.
15. Morissette J T, et al. "Tracking the rhythm of the seasons in the face of global change: phenological research in the 21st century". *Frontiers in Ecology and the Environment* 7.5 (2009): 253-260.
16. Estiarte M and Peñuelas J. "Alteration of the phenology of leaf senescence and fall in winter deciduous species by climate change: effects on nutrient proficiency". *Global Change Biology* 21.3 (2015): 1005-1017.
17. Eriksson O, et al. "Historic hay cutting dates from Sweden 1873-1951 and their implications for conservation management of species-rich meadows". *Biological Conservation* 184 (2015): 100-107.
18. van Asch M and Visser ME. "Phenology of forest caterpillars and their host trees: the importance of synchrony". *Annual Review of Entomology* 52 (2007): 37-55.
19. Lamarre GPA, et al. "Leaf synchrony and insect herbivory among tropical tree habitat specialists". *Plant Ecology* 215.2 (2014): 209-220.
20. Thomson LJ, et al. "Predicting the effects of climate change on natural enemies of agricultural pests". *Biology Control* 52.3 (2010): 296-306.
21. Ollerton J, et al. "How many flowering plants are pollinated by animals?" *Oikos* 120.3 (2011): 321-326.
22. Schupp E W, et al. "Seed dispersal effectiveness revisited: a conceptual review". *New Phytologist* 188.2 (2010): 333-353.
23. Hanya G and Chapman CA. "Linking feeding ecology and population abundance: a review of food resource limitation on primates". *Ecological Research* 28.2 (2013): 183-190.
24. Hawes J E and Peres C A. "Ecological correlates of trophic status and frugivory in neotropical primates". *Oikos* 123.3 (2014): 365-377.
25. Camargo MGG, et al. "Fruit color and contrast in seasonal habitats- a case study from a cerrado savanna". *Oikos* 122.9 (2013): 1335-1342.
26. Kannan R and James DA. "Fruiting phenology and the conservation of the great pied hornbill (*Buceros bicornis*) in the western Ghats of southern India". *Biotropica* 31.1 (1999): 167-177.
27. Galetti M and Dirzo R. "Ecological and evolutionary consequences of living in a defaunated world". *Biological Conservation* 163 (2013): 1-6.
28. Dirzo R, et al. "Defaunation in the Anthropocene". *Science* 345.6195 (2014): 401-406.
29. Wright SJ and Calderon O. "Seasonal, El Niño and longer-term changes in flower and seed production in a moist tropical forest". *Ecology Letters* 9.1 (2006): 35-44.
30. Hoye TT, et al. "Shorter flowering seasons and declining abundance of flower visitors in a warmer Arctic". *Nature Climate Change* 3 (2013): 759-763.
31. Burkle LA and Alarcon R. "The future of plant-pollinator diversity: understanding interaction networks across time, space, and global change". *American Journal of Botany* 98.3 (2011): 528-538.
32. Thébaud E and Fontaine C. "Stability of ecological communities and the architecture of mutualistic and trophic networks". *Science* 329.5993 (2010): 853-856.
33. Memmott J, et al. "Global warming and the disruption of plant-pollinator interactions". *Ecology Letters* 10.8 (2007): 710-717.
34. Laurance WF. "Theory meets reality: how habitat fragmentation research has transcended island biogeographic theory". *Biological Conservation* 141.7 (2008): 1731-1744.
35. Hagen M, et al. "Biodiversity, species interactions and ecological networks in a fragmented world". *Advances in Ecological Research* 46 (2012): 89-210.
36. Athayde EA and Morellato LPC. "Anthropogenic edges, isolation and the flowering time and fruit set of *Anadenanthera peregrina*, a cerrado savanna tree". *International Journal of Biometeorology* 58.4 (2014): 443-454.
37. Carvalho GH and Batalha MA. "The drivers of woody species richness and density in a Neotropical savannah". *Biology Letters* 9 (2013): 20130412.
38. Pausas J G, et al. "Plant functional traits in relation to fire in crown-fire ecosystems". *Ecology* 85.4 (2004): 1085-1100.

39. Williams PR, *et al.* "Germinable soil seed banks in a tropical savanna: seasonal dynamics and effects of fire". *Austral Ecology* 30.1 (2005) 79-90.
40. Paritsis J, *et al.* "Vegetation disturbance by fire affects plant reproductive phenology in a shrubland community in north-western Patagonia, Argentina". *New Zealand Journal of Ecology* 30.3 (2006): 387-395.
41. Alvarado ST, *et al.* "Fire and the reproductive phenology of endangered Madagascar sclerophyllous woodlands". *South African Journal of Botany* 94 (2014): 79-87.
42. D'Antonio CM. "Fire, plant invasions, and global changes". In: Mooney HA, Hobbs RJ (Eds.), *Invasive Species in a Changing World*. Island Press, Washington, DC (2000): 65-93.
43. Alvarado ST, *et al.* "Achieving sustainable conservation in Madagascar: The case of the newly established Ibity Mountain Protected Area". *Tropical Conservation Science* 8.2 (2015): 367-395.
44. Wolkovich EM and Cleland EE. "The phenology of plant invasions: a community ecology perspective". *Frontiers in Ecology and the Environment* 9.5 (2011): 287-294.
45. Vilà M., *et al.* "Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems". *Ecology Letters* 14.7 (2011): 702-708.
46. Morales CL and Traveset A. "A meta-analysis of impacts of alien vs. native plants on pollinator visitation and reproductive success of co-flowering native plants". *Ecology Letters* 12.7 (2009): 716-728.
47. Schnitzer S A, *et al.* "Lianas in gaps reduce carbon accumulation in a tropical forest". *Ecology* 95.11 (2014) 3008-3017.
48. Avalos G., *et al.* "Colonization strategies of two liana species in a tropical dry forest canopy". *Biotropica* 39.3 (2007): 393-399.
49. Morellato LPC and Leitão-Filho H. "Reproductive phenology of climbers in a Southeastern Brazilian forest". *Biotropica* 28.2 (1996): 180-191.
50. Packard S., *et al.* "The Tallgrass Restoration Handbook: For Prairies, Savannas and Woodlands". 2nd New edition. Island Press, Washington DC (2005).
51. Yang W., *et al.* "Dispersal and germination syndromes of tree seeds in a seasonal evergreen monsoon rainforest on Hainan Island, China". *Seed Science Research* 23 (2013): 41-55.
52. Pritchard H W, *et al.* "Ecological correlates of seed desiccation tolerance in tropical African dryland trees". *American Journal of Botany* 91.6 (2004): 863-870.
53. Hart R, *et al.* "Herbarium specimens show contrasting phenological responses to Himalayan climate". *Proceedings of the National Academy of Sciences of the United States of America* 111.29 (2014): 10615-10619.
54. Rawal D S., *et al.* "Herbarium records identify sensitivity of flowering phenology of eucalypts to climate: implications for species response to climate change". *Austral Ecology* 40.2 (2015): 117-125.
55. Schweingruber F H. "Dendrochronology- an extremely exact measuring tool for the study of environmental and human history". *Naturwissenschaften* 83 (1996): 370-377.
56. Brienens RJW., *et al.* "Attaining the canopy in dry and moist tropical forests: strong differences in tree growth trajectories reflect variation in growing conditions". *Oecologia* 163.2 (2010): 485-496.
57. Miller-Rushing A J and Weltzin J. "Phenology as a tool to link ecology and sustainable decision making in a dynamic environment". *New Phytologist* 184.4 (2009): 743-745.
58. Reed BC., *et al.* "Remote sensing phenology: status and the way forward". In: Schwartz, M.D. (Ed.), *Phenology: An Integrative Environmental Science*. Springer, Netherlands (2013): 91-113.
59. Nagendra H., *et al.* "Remote sensing for conservation monitoring: assessing protected areas, habitat extent, habitat condition, species diversity, and threats". *Ecological Indicators* 33 (2013) 45-59.
60. Vázquez D P, *et al.* "Uniting pattern and process in plant-animal mutualistic networks: a review". *Annals of Botany* 103.9 (2009): 1445-1457.

Volume 2 Issue 4 April 2018

© All rights are reserved by Sanjay Kumar and Neha Chopra.