



Pesticide Residues in Strawberries

Ewa Szpyrka*

Department of Analytical Chemistry, Faculty of Biotechnology, University of Rzeszow, Rzeszow, Poland

*Corresponding Author: Ewa Szpyrka, Department of Analytical Chemistry, Faculty of Biotechnology, University of Rzeszow, Rzeszow, Poland.

Received: January 23, 2018; Published: March 23, 2018

Strawberries (*Fragaria ×ananassa* Duchesne) is important fruit of family *Rosaceae* Juss., grown throughout the world. It is rich in vitamin C, sugar, organic acids, anthocyanin, phosphorus, iron, other minerals, vitamins, etc. and its flavour is characterized as fruity, sweet and tart. It is utilized for the production of purees, juice concentrate, juice, jams, preserves and rose red wine. The major strawberry producing countries are USA, Spain, Japan, Poland, Korea and Russian Federation [1].

The protection of strawberries against diseases, pests and weeds is mainly based on the chemical method. In planning protection programs, it is necessary to conduct monitoring in individual phenological phases, which will allow to assess the severity of diseases, and in the case of pests - also to determine the thresholds of danger. The identification of pests is also very important, including the use of familiarity with the symptoms of their feeding [2]. There are 131 plant protection products registered in Poland to control strawberry plantations. Almost half of them are fungicides [3].

The result of chemical protection of strawberry plant are pesticide residues in fruits. According to literature data strawberry fruits contain a lot of pesticide residues, even over 60% of samples contain one or multiple substances, but their concentrations usually do not exceeded their Maximum Residues Levels [4-6]. Pesticide residues most often found in strawberry samples are fungicides: azoxystrobin, boscalid, cyprodinil, difenoconazole, dithiocarbamates, fludioxonil, folpet, pyrimethanil, tebuconazole, pyraclostrobin, tetraconazole, thiophanate-methyl and fenhexamid and insecticide - chlorpyrifos [5,6]. Although strawberries contain a lot of pesticide residues, dietary exposure associated with consumption of these fruits is assessed as low [7,8].

Bibliography

1. Sharma S., *et al.* "An overview on Strawberry [*Fragaria ×ananassa* (Weston) Duchesne ex Rozier] wine production technology, composition, maturation and quality evaluation". *Natural Product Radiance* 8.4 (2009): 356-365.
2. Labanowska BH., *et al.* "Methodology of integrated protection of strawberries". Instytut Ogrodnictwa, Skierniewice and Ministerstwo Rolnictwa i Rozwoju Wsi (2017): 44.
3. The engine of plant protection products. Ministry of Agriculture and Rural Development.
4. "The 2015 European Union report on pesticide residues in food European Food Safety Authority". *EFSA Journal* 15.4 (2017): 4791.
5. Sójka M., *et al.* "Pesticide residue levels in strawberry processing by-products that are rich in ellagitannins and an assessment of their dietary risk to consumers". *NFS Journal* 1 (2015): 31-37.
6. Matyaszek A., *et al.* "Pozostałości środków ochrony roślin w owocach truskawek pochodzących z Polski południowo-wschodniej w latach 2014–2015". *Bromatologia i Chemia Toksykologiczna* 49.2 (2016): 210-216.
7. Winter CK and Katz JM. "Dietary Exposure to Pesticide Residues from Commodities Alleged to Contain the Highest Contamination Levels". *Journal of Toxicology* (2011): 589674.
8. Quijano L., *et al.* "Chronic cumulative risk assessment of the exposure to organophosphorus, carbamate and pyrethroid and pyrethrin pesticides through fruit and vegetables consumption in the region of Valencia (Spain)". *Food and Chemical Toxicology* 89 (2016): 39-46.

Volume 2 Issue 4 April 2018

© All rights are reserved by Ewa Szpyrka.