



Intestinal Integrity of Broilers Exposed to Heat Stress

Arda Sözcü*

Ödemiş Vocational Highschool, Ege University, Turkey

*Corresponding Author: Arda Sözcü, Ödemiş Vocational Highschool, Ege University, Turkey.

Received: October 22, 2018; Published: December 06, 2018

Abstract

Commercial broilers are increasingly being more susceptible to environmental stress factors, due to rapid growth. Due to global warming, heat stress has become one of the most important affecting factors of broiler performance. This review focused on the intestinal integrity of broilers exposed to heat stress during growing period.

Keywords: Broiler; Heat Stress; Intestinal Integrity; Feed Efficiency

Introduction

During last decades, significant developments have been made in broiler performance including an increment in growth rate and feed efficiency, by genetic selection [1]. However, during the genetic selection processes, some vital organs, such as cardiovascular system, respiratory system and skeletal system, could not develop as parallel to heavier body weight of broilers [2]. This imbalance between the major systems and body weight causes some physiological reactions, health problems and finally economical losses, especially under extreme environmental conditions, for example heat stress due to hot climate conditions [3].

Physiological responses to heat stress

Broilers exposed to high environmental temperatures exhibit some physiological reactions to alleviate with heat stress. There are some mechanisms to maintain the body temperature of the bird within a normal ranges, by respiratory-evaporative [4-6] and evaporative cutaneous mechanism [7,8], sensible heat loss via radiation, convection [3] and conductance [9]. During long term heat stress conditions, acid-base balance is exposed to a disturbance and cause a respiratory alkalosis [10]. On the other hand, some behavioural changes are observed in birds as fast panting, moving away from others, lifting their wings, increasing water consumption and reducing the feed consumption. These changes cause detrimental effects for performance and productivity of broilers, and finally results with an increment in mortality [11].

Heat stress causes some serious economical losses by decline in feed intake, but also in digestibility of nutrients [12]. A peripheral vasodilation could develop associated with extreme heat production, then blood flow and subsequently functionality of intestine show a decline [13,14]. In a study performed by Souza, *et al.* [12] reported that broilers that were exposed to longterm heat stress conditions increased heat production with a value of 35.5%, whereas decreased metabolic energy retention and efficiency [by 20.9% and 32.% respectively], nitrogen retention and efficiency [by 50.4% and 33.1%, respectively], compared to the control broilers.

Heat stress and intestinal integrity

Under heat stress conditions, some morphologic and physiologic changes are observed in the gastrointestinal tract on account of functionality and integrity of intestinal epithelium [15-17]. Such as: change in intestinal microflora [18], disruption in blood flow of gastrointestinal tract [19] and deterioration in intestinal morphological traits, such as villus height, crypt depth, villus width, ratio between villus height and crypt depth and villus absorptive surface area [20,21]. Deng, *et al.* [22] found a decline in villus height and ratio between villus height and crypt depth under heat stress. In another study performed by Burkholder, *et al.* [21], it was reported that villus height decreased to 18.8% in birds exposed to 30°C for 24 h, compared to the control birds raised at 23°C.

It is well known that intestinal mucosa has an important barrier role to protect the organism against pathogenic bacterias and also for digestion and absorption of nutrients [23-25]. The integrity of the intestinal barrier is composed of enterocytes, mucus and immune cells, for example macrophages [26-28]. When the integrity of intestinal barrier under heat stress conditions impair [29,30], intestinal permeability and subsequently intestinal inflammation could show an increment [27,31]. Stress conditions cause a decline in protective functionality of intestinal epithelium and increase susceptibility of birds against some diseases, for example *Salmonella* spp. infections [17,21,31].

As a conclusion, when considering adverse effects of heat stress, some solutions and strategies to alleviate of the health problems and economical losses in broiler production, could be developed with regard to nutritional and management interventions.

Bibliography

- Havenstein GB., *et al.* "Growth, livability, and feed conversion of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets". *Poultry Science* 82 (2003a): 1500-1508.
- Havenstein GB., *et al.* "Carcass composition and yield of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets". *Poultry Science* 82.10 (2003b): 1509-1518.
- Yahav S. "Alleviating heat stress in domestic fowl: different strategies". *World's Poultry Science Journal* 65 (2005): 712-739.
- Richards S. "Evaporative water loss in domestic fowls and its partition in relation to ambient temperature". *The Journal of Agricultural Science* 87 (1976): 527-532.
- Seymour SR. "Convective heat transfer in the respiratory systems of panting animals Comparative". *Biochemistry and Physiology* 35.1 (1972): 119-127.
- Marder J and Z Arad. "Panting and acid base regulations in heat stressed birds". *Comparative Biochemistry and Physiology* 94A (1989): 395-400.
- Webster MD and JR King. "Temperature and humidity dynamics of cutaneous and respiratory evaporation in pigeons". *Columba livia Journal of Comparative Physiology* 157.2 (1987): 253-260.
- Ophir E., *et al.* "Cutaneous blood flow in the pigeon *Columba livia*: its possible relevance to cutaneous water evaporation". *Journal of Experimental Biology* 205.17 (2002): 2627-2636.
- Wolfenson D., *et al.* "Evaporative cooling of ventral regions of the skin in heat stressed laying hens Poult". *Science Journal* 80.7 (2001): 958-964.
- Borges SA., *et al.* "Acid-base balance in broilers". *World's Poultry Science Journal* 63.01 (2007): 73-81.
- Syafwan S., *et al.* "Heat stress and feeding strategies in meat-type chickens". *World's Poultry Science Journal* 67.4 (2011):853-673.
- Souza LFA., *et al.* "How heat stress continuous or cyclical interferes with nutrient digestibility, energy and nitrogen balances and performance in broilers". *Livestock Science* 192 (2016): 39-43.
- Yamauchi K. "Review on chicken intestinal villus histological alterations related with intestinal function". *The Journal of Poultry Science* 39.4 (2002): 229-242.
- Attia YA and Hassan SS. "The effect of increasing dietary protein concentration with or without increasing energy levels on improving tolerance of broiler chickens to heat stress". *European Poultry Science* 81 (2017).
- Meddings JB and Swain MG. "Environmental stress-induced gastrointestinal permeability is mediated by endogenous glucocorticoids in the rat". *Gastroenterology* 119.4 (2000): 1019-1028.
- Soderholm JD and Perdue MH. "Stress and gastrointestinal tract. II. Stress and intestinal barrier function". *American Journal of Physiology Gastrointestinal and Liver Physiology* 280.1 (2001): 7-13.
- Quinteiro-Filho WM., *et al.* "Heat stress impairs performance parameters, induces intestinal injury, and decreases macrophage activity in broiler chickens". *Poultry Science* 89.9 (2010): 1905-1914.
- Suzuki K., *et al.* "Effect of crowding and heat stress on intestinal flora, body weight gain, and feed efficiency of growing rats

- and chicks". *Nippon Juigaku Zasshi* 45.3 (1983): 331-338.
19. Wolfenson D. "The effect of acclimatization on blood flow and its distribution in normothermic and hyperthermic domestic fowl". *Comparative Biochemistry and Physiology* 85.4 (1986): 739-742.
 20. Mitchell MA and Carlisle AJ. "The effects of chronic exposure to elevated environmental temperature on intestinal morphology and nutrient absorption in the domestic fowl (*Gallus domesticus*)". *Comparative Biochemistry and Physiology* 101.1 (1992): 137-142.
 21. Burkholder KM., et al. "Influence of stressors on normal intestinal microbiota, intestinal morphology, and susceptibility to *Salmonella enteritidis* colonization in broilers". *Poultry Science* 87.9 (2008): 1734-1741.
 22. Deng W., et al. "The probiotic *Bacillus licheniformis* ameliorates heat stress-induced impairment of egg production, gut morphology, and intestinal mucosal immunity in laying hens". *Poultry Science* 91.3 (2012): 575-582.
 23. Fagarasan S. "Intestinal IgA synthesis: a primitive form of adaptive immunity that regulates microbial communities in the gut". *Current Topics in Microbiology and Immunology* 308 (2006): 137-153.
 24. Keita AV and Soderholm JD. "The intestinal barrier and its regulation by neuroimmune factors". *Neurogastroenterology and Motility* 22.7 (2010): 718-733.
 25. Alhenaky A., et al. "The effect of heat stress on intestinal integrity and *Salmonella* invasion in broiler birds". *Journal of Thermal Biology* 70(B) (2017): 9-14.
 26. Singleton KD and Wischmeyer PE. "Oral glutamine enhances heat shock protein expression and improves survival following hyperthermia". *Shock* 25 (2006): 295-299.
 27. Chappell L., et al. "The immunobiology of avian systemic salmonellosis". *Veterinary Immunology and Immunopathology* 128.1-3 (2009): 53-59.
 28. Lambert GP. "Stress-induced gastrointestinal barrier dysfunction and its inflammatory effects". *Journal of Animal Science* 87.14 (2009): 101-108.
 29. Niu ZY., et al. "Effects of different levels of vitamin E on growth performance and immune responses of broilers under heat stress". *Poultry Science* 88.10 (2009): 2101-2107.
 30. Song J., et al. "Effect of a probiotic mixture on intestinal microflora, morphology, and barrier integrity of broilers subjected to heat stress". *Poultry Science* 93.3 (2014): 581-588.
 31. Quinteiro-Filho WM., et al. "Acute heat stress impairs performance parameters and induces mild intestinal enteritis in broiler chickens: role of acute HPA axis activation". *Journal of Animal Science* 90.6 (2012): 1986-1994.

Volume 3 Issue 1 January 2019

© All rights are reserved by Arda Sözcü.