



Endophytic Fungi for Plant Growth Promotion and Adaptation under Abiotic Stress Conditions

Ajar Nath Yadav*

Department of Biotechnology, Akal College of Agriculture, Eternal University, Baru Sahib, India

*Corresponding Author: Ajar Nath Yadav, Department of Biotechnology, Akal College of Agriculture, Eternal University, Baru Sahib, India.

Received: December 03, 2018; Published: December 14, 2018

The endophytic fungi are colonizing the interior of the plant parts. Among diverse microbial groups, endophytic fungi may play a biological role with their host plant for different attributes. Endophytic fungal interactions with plants confer protection against to abiotic stresses of temperature, salinity, pH and water deficient. The plant infected with endophytic fungi results in significant increase in biomass, improved commercial plant production and thus are useful in agro-forestry and biotechnological agricultural applications. The signalling molecules such as flavonoids, isoflavonoids, and phenolic are released from the roots of plant attracting the fungus from rhizosphere to colonize inside the plant as an endophyte. Fungal endophytes colonize the tissues of host; transmission of endophytes follows two routes vertical transmission of endophytic fungi take place from mother plant to offspring, via seeds, often referred to as "true endophytes". Horizontal transmission occurs via soil or air-borne spores. To date, endophytes in forbs infect the leaves of their hosts through horizontal transmission.

Plant- microbe interaction resulted in adaptation, plant growth promotion, uptake of micronutrient and production of different groups of secondary metabolites and bioactive compounds with potential applications in agriculture, medicine, industry allied sectors [1-4]. Due to protection of plant against biotic and abiotic stresses the beneficial endophytic fungi are considered as an eco-friendly bioresources. The members of different phylum such as Ascomycota, Basidiomycota, Mucoromycota and Oomycota have been reported as plant growth promoters and protectants under the natural normal as well as abiotic stress conditions [5]. Among the fungal groups, arbuscular mycorrhizal (AM) fungi are known to promote activities which can improve agricultural developments. In exchange for the AM fungi providing all of these nutrients, the plant in turn provides the mycorrhizae with carbon and other nutrients [6-10]. A large number of endophytic fungal species belonging to different genera including *Acremonium*, *Alternaria*, *Aspergillus*, *Berkleasium*, *Chaetomium*, *Cladosporium*, *Claviceps*, *Collectotrichum*, *Cryptococcus*, *Curvularia*, *Fusarium*, *Geomyces*,

Glomus, *Leptospora*, *Metarhizium*, *Microdochium*, *Neotyphodium*, *Ophiognomonina*, *Paecilomyces*, *Penicillium*, *Phaeomonella*, *Phyllosticta*, *Piriformospora*, *Rhizoctonia*, *Rhizopus*, *Rhodotorula*, *Talaromyces*, *Trichoderma*, *Walleimia* and *Xylaria* have been isolated from different host plants [1,6,10,11].

Endophytic fungi are agriculturally important as they can enhance plant growth; improve plant nutrition through different direct and indirect PGP attributes including solubilization of phosphorus, potassium and zinc; production of phytohormones (Indole acetic acids, gibberellic acids and cytokinin), Fe-chelating compounds, hydrolytic enzymes, hydrogen cyanide, ammonia [6,10,12-15]. In endophytic relationships, growth promoting fungi reside within the apoplastic spaces in the host plants. Sustainable agriculture requires the use of strategies to increase or maintain the current rate of food production while reducing damage to the environment and human health. The use of microbial plant growth promoters is an alternative to conventional agricultural technologies. Endophytes start their journey as rhizospheric microbes or soil microbiomes by different mechanisms. Plants deal with a series of adverse environmental and biotic stresses like drought, cold, hyper salinity, or pathogenesis. Fungal endophytes help plants to adapt under the abiotic stress of temperature, pH, salinity and drought via indirect mechanisms e.g. through induced systemic resistance (ISR), bioremediation, biocontrol. The fungi present inside tissue have capability to protect plants from pathogens and insects.

The fungal endophytes have significant roles for living organisms and human health due to production of wide range of bioactive compounds. The endophytic fungi along with other plant growth promoting microbes as single or in consortium could potentially be used as future commercial bioinoculants and biocontrol agents for crops growing under the normal as well as abiotic stress conditions for sustainable agriculture and environments. The significance of enzymes and other bioactive compounds produced by fungal endo-

phytes may be used for bioremediation of environmental pollutants such as polychlorinated hydrocarbons, and polyaromatic hydrocarbons for sustainable environments. For exploring the endophytes of biotechnological importance isolation of endophytes by laboratory techniques and scale up are the important steps. Recently, interest has been generated for the research of endophytic microbiomes in different plants for secondary metabolites for potential applications in industry, agriculture, pharmaceuticals and allied sectors may be by advancements in genetic engineering, metagenomics, meta-transcriptomics, microbial fermentation, proteomics, and drug design techniques intelligent productivity of some potential candidates for discovering new drugs can be increased. Future research on endophytic fungi are of biotechnological interest due to their potential of being used as genetic vectors, biological control agents, source of secondary metabolites, antimicrobial agent, antitumor compounds, antibiotics, immunosuppressant, production of natural antioxidant, antiviral compounds, insecticidal products, and antidiabetic agents.

Acknowledgement

The authors are grateful to Prof. Harcharan Singh Dhaliwal, Vice Chancellor, Eternal University, Baru Sahib, Himachal Pradesh, India for providing infra-structural facilities and constant encouragement.

Competing Interests

The authors declare no conflict of interest.

Bibliography

1. Yadav AN., et al. "Microbiome in Crops: Diversity, distribution and potential role in crops improvements, in Crop Improvement through Microbial Biotechnology, R Prasad, SS Gill, and N Tuteja, Editors". *Elsevier: USA* (2018): 305-332.
2. Biswas S., et al., "Study on the activity and diversity of bacteria in a New Gangetic alluvial soil (Eutrocept) under rice-wheat-jute cropping system". *Journal of Environmental Biology* 39.3 (2018): 379-386.
3. Yadav AN., et al. "Psychrotrophic Microbiomes: Molecular Diversity and Beneficial Role in Plant Growth Promotion and Soil Health, in Microorganisms for Green Revolution-Volume 2: Microbes for Sustainable Agro-ecosystem, DG Panpatte, et al., Editors". *Springer: Singapore* (2018): 197-240.
4. Yadav N., et al. "Biodiversity and biotechnological applications of novel plant growth promoting methylotrophs". *Journal of Applied Biotechnology and Bioengineering* 5.6 (2018): 342-344.
5. Yadav AN., et al. "Beneficial microbiomes: Biodiversity and potential biotechnological applications for sustainable agriculture and human health". *Journal of Applied Biology and Biotechnology* 5.6 (2017): 1-13.
6. Suman A., et al. "Endophytic Microbes in Crops: Diversity and Beneficial impact for Sustainable Agriculture, in Microbial Inoculants in Sustainable Agricultural Productivity, Research Perspectives, D Singh, P Abhilash, and R Prabha, Editors". *Springer-Verlag: India* (2016): 117-143.
7. Finlay RD. "Ecological aspects of mycorrhizal symbiosis: with special emphasis on the functional diversity of interactions involving the extraradical mycelium". *Journal of Experimental Botany* 59.5 (2008): 1115-1126.
8. Bonfante P., et al. "Mechanisms underlying beneficial plant-fungus interactions in mycorrhizal symbiosis". *Nature Communications* 1 (2010): 48.
9. Mathur N., et al. "Mycorrhiza: A Potent Bioinoculant for Sustainable Agriculture in Microbial Technology "The Emerging Era" H Pathak and A Sharma, Editors". *Lap Lambert Academic Publishing Ag and Co. Kg: Dudweiler Landstr, Germany* (2011): 230-245.
10. Verma P., et al. "Beneficial Plant-Microbes Interactions: Biodiversity of Microbes from Diverse Extreme Environments and Its Impact for Crop Improvement, in Plant-Microbe Interactions in Agro-Ecological Perspectives: Volume 2: Microbial Interactions and Agro-Ecological Impacts, DP Singh, HB Singh, and R Prabha, Editors". *Springer Singapore: Singapore* (2017): 543-580.
11. Yadav AN., et al. "Biodiversity of the Genus Penicillium in Different Habitats, in New and Future Developments in Microbial Biotechnology and Bioengineering, Penicillium System Properties and Applications, VK Gupta and S Rodriguez-Couto, Editors". *Elsevier: Amsterdam* (2018): 3-18.
12. Saxena AK., et al. "Microbial diversity of extreme regions: An unseen heritage and wealth". *Indian Journal of Plant Genetic Resources* 29.3 (2016): 246-248.
13. Yadav AN., et al. "Plant microbiomes and its beneficial multifunctional plant growth promoting attributes". *International Journal of Environmental Sciences and Natural Resources* 3.1 (2017): 1-8.

14. Yadav AN. "Biodiversity and Biotechnological Applications of Host-Specific Endophytic Fungi for Sustainable Agriculture and Allied Sectors". *Acta Scientific Microbiology* 1.5 (2018): 01-05.
15. Yadav AN., *et al.* "Stress-Adaptive Microbes for Plant Growth Promotion and Alleviation of Drought Stress in Plants". *Acta Scientific Agriculture* 2.6 (2018): 85-88.

Volume 3 Issue 1 January 2019

© All rights are reserved by Ajar Nath Yadav.