



False Smut (*Ustilaginoidea Virens* (Cke.) Tak.): A Serious Major Constraint in Rice (*Oryza Sativa* L.) Production

H Mahmud*

Additional Deputy Director (Crop), Department of Agricultural Extension, Kushtia, Bangladesh

*Corresponding Author: H Mahmud, Additional Deputy Director (Crop), Department of Agricultural Extension, Kushtia, Bangladesh.

Received: December 03, 2018; Published: December 14, 2018

The rice false smut is a worldwide disease and already changed from a minor disease to a major disease in many rice-growing countries in Asia since 1970. It has been found in many countries, viz; China, India, Japan, Italy, Australia, Philippines, Brazil and Myanmar [1, 2]. Rice false smut disease was also found to occur severely in BRRI dhan49 during Aman season with remarkable yield losses in Bangladesh. With the change of weather condition, large application of nitrogen fertilizer and large-scale planting of hybrid rice, the rice false smut has become more and more serious. Outbreaks of this disease often occur in cold weather and reduce the grain quality and yield [3], and late sowing and application of high doses of Nitrogen also favours the development of disease [4].

The highest disease incidence (61.20%) and yield loss (14.18%) were reported [5], and yield losses caused by RFS (rice false smut) disease is attributed to both smut balls as well as chaffiness, reduction in grain weight and infertility of the spikelet near the smut balls [6]. The false smut balls have toxin including ergot alkaloid toxin that can cause rumination stopping in cows, suppress the tubulin of mammals and cause necroses of liver, kidney, and bladder tissues in mice [7, 8].

The false smut (chlamydospores) contains mycotoxins (ustiloxins) that are toxic to animals and contaminate rice seeds and grains [4]. The chlamydospores survive in winter in the soil and become a primary source of infection of the rice plants [3]. The chlamydospores germinate on coleoptile epidermal cells of rice seedlings, and infected hyphae invade intercellular spaces and reach at the meristematic tissues of rice plants [3]. In addition, the scientists [9] confirmed in greenhouse tests that rice seedlings grown from infested seeds were infected by *U. virens*. However, confusion remains about the disease cycle and infection process in rice false smut [10]. Moreover, false smut not only threatens rice production in yield and quality but also produces toxins that are dangerous to human health and livestock.

Significant effect in controlling severity of false smut was observed with Potent 250 EC (Propiconazole 0.1%) as reported [11]. Researchers [12] reported that two sprays of 50% propiconazole EC at 300 g a.i. ha⁻¹ and of 10% difenoconazole GR at 225 g a.i. ha⁻¹, exhibited the best control of rice false smut. Treated seed with trifloxystrobin (Trilex 2000), and propiconazole plus trifloxystrobin (Stratego) were also effective in reducing false smut with foliar application at the heading stage [13]. Efficacy of indore isolates of *Trichoderma* spp. against *U. virens* showed significant antagonistic potential [14].

The integrated use of fungicides and good agricultural practices appear to be effective for management of false smut disease of rice.

Bibliography

1. SH Ou. "Rice Diseases". M. Commonwealth Mycological Institute (1972): 288-295.
2. DS Dodan and R Singh. "False smut of rice: present status". *Agriculture Research* 17.4 (1996): 227-240.
3. T Ashizawa., et al. "Quantification of the rice false smut pathogen from soil in Japan using real-time PCR". *European Journal of Plant Pathology* 128 (2010): 221-232.
4. M Ahonsi., et al. "Effect of variety and sowing date on false smut incidence in upland rice in Edo State, Nigeria". *IRRI Notes* 25 (2000): 14.
5. A Singh., et al. "Occurrence of rice false smut in Bundelkhand region". *International Journal Plant Protection* 5.2 (2012): 283-285.
6. R Rani. "Variability in *Ustilaginoidea virens* (Cke.) Tak. causing false smut of rice and identification of resistance sources". Department of Plant pathology, Punjab Agricultural University, Ludhiana-141004, India (2014).

7. KI Nakamura., *et al.* "Lupinosis"-like lesions in mice caused by ustiloxin, produced by *Ustilagoidea virens*". *A morphological study. Natural Toxins* 2 (1994): 22-28.
8. Y Li., *et al.* "Ustiloxins new antimitotic cyclic peptides: Interaction with porcine brain tubulin". *Biochemical Pharmacology* 149 (1995): 1367-1372.
9. M Ditmore., *et al.* "Infection of plants of selected rice cultivars by the false smut fungus, *Ustilagoidea virens*, in Arkansas" *Pest Management: Diseases. B. R. Wells Rice Research Studies, AAES Research Series* 550 (2006): 132-138.
10. Y Tang., *et al.* "Elucidation of the infection process of *Ustilagoidea virens* (teleomorph: *Villosiclava virens*) in rice spikelets". *Plant Pathology* 62 (2013): 1-8.
11. H Mahmud and I Hossain. "Comparative efficacy of BAU-Bio-fungicide and synthetic fungicides in management of diseases of rice (*Oryza sativa* L.) for quality seed production". *Brazilian Journal of Botany* 40.2 (2017): 389-397.
12. Y Chen., *et al.* "Frequency distribution of sensitivity of *Ustilagoidea virens* to four EBI fungicides, prochloraz, difenoconazole, propiconazole and tebuconazole, and their efficacy in controlling rice false smut in Anhui Province of China". *Phytoparasitica* 41 (2013): 277-284.
13. X Zhou. "Field evaluation of fungicides for management of rice false smut and kernel smut". Texas A&M University System, Agri Life Research, Beaumont, TX, USA (2012).
14. A Kumar., *et al.* "Influence of *Trichoderma* spp. against *Ustilagoidea virens* inciting false smut of rice". *Environment and Ecology* 32.1 (2014): 163-168.

Volume 3 Issue 1 January 2019

© All rights are reserved by H Mahmud.