



## Comparison Between Value of Indian Major Carps and Air Breathing Carps in Open Inland Fishery Sector

Swapna Ghosh<sup>1</sup> and Pratik Majumdar<sup>2\*</sup>

<sup>1</sup>Department of Biotechnology, Neotia Institute of Technology Management and Science, Sarisha, West Bengal, India

<sup>2</sup>ICAR-Central Inland Fisheries Research Institute, Biotechnology Laboratory Barrackpore, Kolkata, West Bengal, India

**\*Corresponding Author:** Pratik Majumdar, ICAR-Central Inland Fisheries Research Institute, Biotechnology Laboratory Barrackpore, Kolkata, West Bengal, India.

**Received:** July 20, 2020

**Published:** August 28, 2020

© All rights are reserved by **Swapna Ghosh and Pratik Majumdar.**

### Abstract

In India, 75% population are live in rural area and agriculture fish farming are major occupation of them who are live in village. The total fish production of 10.07 million metric tonnes in India presently has nearly 65% contribution from the inland sector and nearly the same from culture fisheries. Actually Fish constitutes a major source of protein in our diet. Fish apart from being important in human diet, its fatty acid are currently under intense scientific investigation because of numerous health benefits attributed by them. The amount of protein in fish muscle is usually somewhere between 15-20%, but values lower as 15% or as high as 28% are occasionally met with in some species. IMC are demandable and easily acceptable for consumers, it can fulfil population demand for their nutritional value in everyday life and air breathing fish are not easily available and acceptable by all classes of consumer. Among IMC Rohu (*Labeo rohita*) achieve the 1<sup>st</sup> position as it can fulfil population demand for their nutritional value at a lower cost for the above reasons. Even induced breeding become more popular in IMC where as the anthropogenic activity is tough for air breathing fishes. Their seeds collected from the natural environment, but due to exposure of chemicals in land and water the population become enough less day to day. So we have to save them from the list of endangered species for their nutritious value as compare to protein and lipid.

**Keywords:** Indian Major Carps (IMC); Air Breathing Fish; Protein; Lipid; Human Diet; Population Demand; Scientific Investigation; Induced Breeding; Nutritious

### Introduction

Fish constitutes a major source of protein in our diet and its growth depends on number of factors such as food, space, temperature, salinity, season and physical activity [1]. Fish have a low caloric value and low fat but high protein content. Studies on the chemical composition of fish have been carried out on marine fish, but analyses carried out on fresh water fish are rare. Fish can

remove from malnutrition of people easily. The live weight of Majority of fish usually consists of about 70-80% of water, 20-30% of protein, 2-12% of lipid [2]. The reason for superior food conversion efficiency of fish is due to assimilation of protein rich diet because of their lower dietary energy requirements. Fish helps to reduce blood pressure by small amount but improve blood clotting regulation. All proteins are chains of chemical units. They are

linked together to make one long molecule. These units, of which there are about 20 types, are called amino acids. Among all proteins some are essential in the human diet for the maintenance of good health. Lysine and methionine are two essential amino acids, present in high concentrations in fish proteins. Fish protein gives a good combination of amino acids which is highly active to people’s nutritional requirements. The main body constituents of the fish are including water, lipid, ash, protein. Carbohydrates and non- protein compounds are also important constituents but are present in small amount and are usually ignored during analysis [2,3]. The study objective is comparison between value of Indian major carps and air breathing carps in open inland fishery sector.

**Materials and Methods**

**Study area**

Fresh fish samples of *Labeo rohita* (rohu), *Anabas testudineus* (koi), *Heteropneustes fossilis* (singhi) were purchased from whole seller fish market (Gate Bazar) at Midnapore Town, Paschim Midnapore.

**Methods of data collection**

We had visited the market and collected the data from the sellers from March to May 2017 about the different types of fish. By consulting the buyers also, our objectives matched with them that air breathing fishes are how much costlier and demandable also.

**Study design**

The fish samples were immediately packed in autoclaved poly bag and brought into the laboratory. Fishes are washed with distilled water, measured the length and weighted by digital weight machine. They are all approximately 100 gm.

SL NO.	Name of fish	Length
1.	Rohu	20 cm
2	Singhi	23.8 cm
3.	Koi	16.2 cm

**Table 1:** Length of Fishes.

**Frequency of analysis -biochemical estimation**

Protein can be estimated from the fish flesh according to the methods of Lowry et al. [4] using Folin -Phenol reagent. 200 mg of muscle tissue was taken from fish muscle and homogenized in 5.0

ml of 0.1(N) NaCl. Then 2.0 ml of 5% PCA was added to the above mixture. Centrifuged the homogenate at 10,000 rpm for 10 minutes and then supernatant was discarded. Again washed with 2.0 ml of 5% PCA. 5.0 ml of diethyl ether was added to the pellet and again centrifuged at 10,000 rpm for 10 minutes and then decant the supernatant. The pellet was dissolved in 2.0 ml of 0.1(N) NaOH. This mixture used for protein determination.

Pipette out 0.2ml of above mixture in a test tube. Then made up the volume to 0.5 ml with 0.1(N) NaOH/ distilled water. Blank were prepared contains 0.5ml of distilled water instead of sample mixture. 5.0 ml of copper reagent was added to each of the test tube including blank. The mixture was properly mixed by cyclomixture and allowed to stand for 10 minutes. Then 0.5 ml of Folin Phenol reagent was added to each of the test tube and incubated at room temperature for 30 minutes. Blue colour developed after 30 minutes was read in Spectrophotometer at 660nm. Standard curve was prepared earlier using BSA as a protein source and protein content was calculated with the help of this standard curve. The values were expressed as mg/g wet weight of tissue and the concentrations were expressed as percentage. This value can be obtained from the graphical method of Lawry standard curve.

Lipid from the fish muscle was estimated by the method of Folch et al. [5] using chloroform methanol (1:1) mixture. Then lipid content was calculated by the following formula. Lipid content in the muscle (g/kg) = (F-I) x 100/X g/g% Where, F= final weight of watch glass, I= initial weight of watch glass, x= g muscle sample taken for estimation.

**Results and Discussion**

The information of protein and lipid are important for the fish used as food by the consumers. This also facilitates the selection of most appropriate species having higher protein contents and condition for human consumption. The body composition is used as an indicator to assess the nutritional status and condition of fish.

	Fish samples	Protein (g/g%) (avg)	Lipid (g/g%) (Avg)
1.	Singhi (stinging cat fish)	18.6± 1.3	1.4 ± 0.2
2.	Koi (climbing perch)	16.9± 0.3	4.4± 0.8
3.	Rohu	15.49± 0.4	2.9 ± 0.1

**Table 2:** Protein Lipid average SD value.

### Future Aspects

The average value of protein variation was 16.99% which is near to the findings of [6,7] the protein content is 18-20% [8]. The finding results of protein are most of the same value with the above table 2. The variation of lipid ranges from 1.5% to 5.2%. In general average value of lipid is 7.35% which is supported by the study of [6,7].

Air breathing fishes can survive in waste water bodies. In case of aerobic hypoxia it can take oxygen from air by their air breathing organ. No need of supplementary food in shallow water bodies. Cat fish (Magur, Singhi, Koi) plays as an important role in everyday life as a nutritious food for medical treatment. But in case of artificial breeding the breeding rate and survival rate of fry is too low. We are trying for better improvement through artificial breeding.

### Conclusion

Indian major carps are highly valuable, demandable and nutritional food for the population. According to cost IMC is better than breathing fishes. For this cause we should be increase the production of IMC, so that maximum population will be eligible to buy this type of fishes in future. If it's possible so we can make a new invention for all.

### Bibliography

1. Weatherly AH and Gill HS. "The biology of fish growth". Academic Press, London, UK (1987).
2. Love RM. "The chemical biology of fishes". Academic Press 1 London, U.K (1970).
3. Cui Y and Wootton RJ. "Bioenergetics of growth of cyprinids, Phoxinus, the effect of the ration and temperature on growth rate and efficiency". *Journal of Fish Biology* 33(1988): 763-773.
4. Lowry OH., et al. "Protein measurement with the Folin phenol reagent". *Journal of Biological Chemistry* 193 (1951): 265-275.
5. Folch J., et al. "A simple method of the isolation and purification of total lipid from animal tissue". *Journal of Biological Chemistry* 226 (1957): 497-509.
6. Pradhan SC., et al. "Biochemical studies of muscle". *International Journal of Current Research* 4.12 (2012): 228-234.

7. Mahboob S., et al. "Proximate composition of wild and farmed *Labeo rohita* (Rohu), Pakistan". *Journal of Zoology* 36 (2004): 39-43.
8. Jindal M., et al. "Effect of feeding dietary protein sources on daily excretion in *Channa punctatus* for sustainable aquaculture". *Advances in Aquatic Ecology* 21 (2008).

#### Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: [www.actascientific.com/](http://www.actascientific.com/)

Submit Article: [www.actascientific.com/submission.php](http://www.actascientific.com/submission.php)

Email us: [editor@actascientific.com](mailto:editor@actascientific.com)

Contact us: +91 9182824667