

Effect of Some *Rhizobium* Strains on Fenugreek Growth and Biological Control of Sclerotinia Stem Rot of Fenugreek Caused by *Sclerotinia trifoliorum*

Hemissi Imen, Hachana Amira*, Sonia Mejri and Arfaoui Hanen

Agronomic Sciences and Techniques Laboratory, National Institute of Agricultural Research of Tunisia (INRAT), Carthage University, Tunisia

*Corresponding Author: Hachana Amira, Agronomic Sciences and Techniques Laboratory, National Institute of Agricultural Research of Tunisia (INRAT), Carthage University, Tunisia.

Received: November 23, 2020

Published: April 16, 2021

© All rights are reserved by Hemissi Imen, et al.

Abstract

Fenugreek is an annual leguminous crop grown for hay and grains in Tunisia. It is also considered as a valuable rotation crop with cereals. Sclerotinia rot was observed in production fields since 2010. The aim of this work is to study the antagonistic activity of different *Rhizobium* strains against *Sclerotinia trifoliorum* in dual culture *in vitro* and under greenhouse conditions. Among the 32 strains tested, 26 isolates had effective control on *Sclerotinia trifoliorum in vitro*. In order to study the biological control mechanisms, the *Rhizobium* strains ability to produce volatile compounds and to solubilize phosphate were investigated. The results showed that 18 strains were able to solubilize phosphorus and 19 strains produced volatile compounds. In pot trials, the percentage of fenugreek plants inoculated with different rhizobia showed significant reduce sclerotinia stem rot symptoms compared to the uninoculated plants. Among these rhizobiums, the strain Soli proved efficient against the pathogen *in vitro* and in pot experiments. The results promise the use of rhizobia for protection of fenugreek against *Sclerotinia trifoliorum*.

Keywords: Antagonism; Biocontrol; *Rhizobium*; *Sclerotinia trifoliorum*; *Trigonella foenum graecum*

Introduction

Fenugreek (*Trigonella foenum graecum*) is an annual plant belongs to the family *Leguminosae*. It is the famous spices in human food. The seeds and green leaves of fenugreek are used in food as well as in medicinal application that is the old practice of human history.

An increasing interest is given to this crop in Tunisia. The fenugreek is grown in rotation with cereals and used mainly for hay and grains. Many investigations have demonstrated that fenugreek seeds of Tunisian cultivars are a rich source of protein, fat, fiber and carbohydrates and suggested that it should be considered in the food and feed consumption [1,2]. Other research has indicated the medicinal benefits of this legume [3,4]. Haouala, et al. [5] and Omezzine, et al. [6] have reported the antifungal, insecticidal and allelopathic potential of fenugreek extracts from aerial parts of a Tunisian cultivar.

Fenugreek is subjected to attack by number of diseases. Among these diseases damping-off, root-rot caused by *Rhizoctonia solani* Kuhan, *Fusarium solani* Mart and *Macrophomia phaseolina* [7]. Recently, Gargouri, et al. [8] have reported the occurrence of sclerotinia stem rot of fenugreek caused by *Sclerotinia trifoliorum* and *S. sclerotiorum* in Tunisia.

Management of these diseases is achieved mainly by the use of resistant cultivars and fungicide application. But frequent application of fungicides to the soil has not only caused environmental hazards like water and soil pollution but also destruction of non-target beneficial microorganisms in soil. Recently, biocontrol approaches have been initiated by using antagonistic microorganisms to combat these diseases.

Biological control agents for plant diseases are currently being examined as alternatives to synthetic pesticides due to their perceived level of safety and minimal environmental impacts. Strains of several bacterial species such as *Bacillus*, *Pseudomonas* and recently the *Rhizobium* group were isolated and found to effectively control various soil-borne plant pathogenic fungi under greenhouse and field conditions. As compared to the other biocontrol agents, Rhizobia offer the great advantage of symbiotic nitrogen fixation by association with legumes [9]. Rhizobia have several mechanisms of action that allow them to control pathogens. These mechanisms include competition for iron by production of siderophores [10,11], competition for nutrients [12], production of antibiotics [13,14], promotion of plant growth, in terms of better shoot height, root length, dry weight and root nodulation [15,16], and induction of plant defence mechanisms [17]. The aim of the present work was to select *Rhizobium* isolates with antagonistic activity against *sclerotinia* stem rot of fenugreek caused by *Sclerotinia trifoliorum*. Isolates were purified from fenugreek nodules and their effects on fungal growth and disease development were assessed *in vitro*, in dual cultures, as well as *in vivo* under greenhouse conditions.

Materials and Methods

Plant material

The seeds of fenugreek variety 'Ryhana' used in this work were provided by the Legume Program, Field Crop Laboratory, Institut National de la Recherche Agronomique de Tunisie (INRAT).

Bacterial isolates

Rhizobium isolates were obtained from nodules of 50 days old fenugreek plants using the crushed nodule method [18]. All isolates were purified and tested for their ability to form nodules on fenugreek as previously described [19]. Thirty two isolates were collected from different localities (Table 1). These strains were grown at 28°C [18] on a yeast extract mannitol medium containing 0.08% yeast extract (w/v) and 1% mannitol (w/v). Stocks of strains were prepared on yeast extract-mannitol agar and kept at -70°C (under 30% of glycerol) for long-term storage and at 4°C as source cultures. A culture was repeated every 6 months to have stocks of younger generations.

N°	<i>Rhizobium</i> strains	Localities	N°	<i>Rhizobium</i> strains	Localities
1	Mat 1/6	Mateur	17	Ari 2	Ariana
2	Mat 1/3	Mateur	18	Tak	Takelsa
3	Mat 1/10	Mateur	19	Jbn	Jbeniana
4	Mat 3/16	Mateur	20	Uti	Utique
5	Mat 3/26	Mateur	21	Mat M	Mateur
6	Mat 3/17	Mateur	22	Elj	Eljem
7	Mat 3/22	Mateur	23	INRAT	Ariana
8	Mat 3/15	Mateur	24	Soli	Soliman
9	Mat 1/11	Mateur	25	Morg2	Morneg
10	Mat 1/1	Mateur	26	Bej	Beja
11	Mat 1/9	Mateur	27	Sad	Saida
12	Mat 3/28	Mateur	28	Sbit	Sbeitla
13	Mat 3/25	Mateur	29	Mok	Mokenine
14	Morg1	Morneg	30	Morg3	Morneg
15	MenT	Menzel Temim	31	Han	Hancha
16	Ari 1	Ariana	32	Mid	Mida

Table 1: Rhizobium strains collected from different localities used in experiments.

Fungal isolate

Sclerotinia trifoliorum was originally isolated from fenugreek in naturally infested field in northern Tunisia, according to the method described [11]. A fungal culture of the pathogen was cultured on PDA in 9-cm-diameter Petri dishes at 25 °C in the dark for 10 d for the production of *sclerotia*.

Antagonism test in dual culture

The *in vitro* inhibition of mycelial growth of *Sclerotinia trifoliorum* by the bacterial isolates was tested using the dual culture technique as described by Paulitz., *et al.* [20] and Landa., *et al.* [21]. Three 50 µl drops from the 10⁸ CFU ml⁻¹ bacterial suspension were equidistantly placed on the margins of PDA plates and incubated at 25°C for 24 h. A 4-mm agar disc from fresh PDA cultures of *Sclerotinia trifoliorum* was placed at the center of the PDA plate for each

bacterial isolate and incubated at 28°C for 7 days. The radius of the fungal colony towards and away from the bacterial colony was measured. The percentage growth inhibition was calculated using the following formula:

$$\% \text{ inhibition} = (R-r)/r \times 100$$

Where r is the radius of the fungal colony towards the bacterial colony and R is the maximum radius of the fungal colony away from the bacterial colony.

Eluciation of antagonistic traits : Volatile antifungal compounds

The production of volatile antifungal compounds by the *Rhizobium* isolates was assayed by a sealed plate method as described by Fiddman and Rossall [22]. From 72-h-old cultures of *Rhizobium* strains in yeast extract mannitol (YEM) liquid media [18], 200µl was spread on YEMA medium in a Petri dish. After incubation at 28°C for 24 h, a second Petri dish containing PDA was inoculated with a plug (Φ: 6 mm) of the test fungus in the center of the plate, inverted and placed over the bacterial culture. Each two plates, containing pathogen and bacteria, were sealed together with parafilm and incubated at 25°C. This ensured that both organisms were growing in the same atmosphere though physically separated. As a control, a Petri dish containing YEMA medium without bacteria was placed under the PDA medium inoculated with the fungal pathogen. Fungal growth was measured as increases in the radial length after 5 days. Each test was replicated three times

Phosphate solubilization efficiency

Bacterial isolates were then employed for phosphate solubilization by streaking them on the Pikovskaya's agar medium [23] and incubated for 7 days at 28±2 °C. The phosphate solubilization was expressed as positive and negative depending on the halo zone formation. The size of the clear zone around the colonies showing phosphate solubilization was noted. The P solubilizing ability of bacteria can be assessed in terms of the solubilization index (SI). As described in several studies [24,25], phosphate SI can be determined using the following formula:

$$SI = \frac{\text{Colony diameter} + \text{halo zone diameter}}{\text{Colony diameter}}$$

Evaluation of antagonistic activity of rhizobacteria against pathogenic fungi

The study was conducted to determine the efficacy of *Rhizobium* strains to reduce the incidence of *sclerotinia* stem rot on fenugreek cultivar 'Ryhana'. Seeds were surface disinfected in 2% NaOCl for 3 min, washed three times with sterile distilled water and germinated on autoclaved layers of paper towels in moist chamber at 25°C for 3 days. Germinated seeds, selected for the uniformity (length of radicle), were sown in perlite at a depth of 2 cm. After transplanting of seedling, each pot was drenched with 5 ml of each of the bacterial suspension (10⁸ cells/ml).

Preparation of the fungal inoculants

Two centimeters long surface sterilized fenugreek stem sections were colonized by placing them for two weeks on individual PDA cultures with *Sclerotinia triflorum* isolate. A colonized section was placed against each of three stems per pot and maintained in position with a wooden toothpick. A sterile fenugreek stem section was placed against each of the control stems. Pots were placed in a green house with a temperature regime of approximately 25 °C during the day and 16 °C at night and were irrigated daily. The treatments in the *in vivo* biocontrol experiment were as follows: Plant inoculated with *Sclerotinia triflorum* and rhizobial strains, plants inoculated only with *Sclerotinia triflorum* (control +) and non-inoculated fenugreek (control -). The percentage of dead plants was assessed two weeks after inoculation. The experimental design was a randomized complete block design with 6 replicates (pots) consisting of three plants per pot per isolate. Diseased plants were collected, seedling length shoot dry weight were recorded at the end of the experiment. Samples were kept for 72 h at (60± 2) °C in an oven for measuring the dry weight of seedling.

The virulence of pathogen was evaluated at scale indices from 0 to 4 according Tezcan and Yildiz [26]. The recorder data were based on mass disease index (MDI %) according to the formula :

$$MDI (\%) = \sum_4^1 (n \times i / 4N) \times 100$$

n: number of plant with indices i

N: umber of total plants.

Statistical analysis

All experiments were replicated as completely randomized blocks. Consequently, the data are means ± confidence interval (n = 3, α = 0.05). Statistical analysis (ANOVA) was performed with SPSS 20.0 for windows, followed by Duncan’s multiple range test (P = 0.05).

Results and Discussion

***In vitro* experiments**

Fungal inhibition assays

Among the 32 *Rhizobium* strains tested in dual culture with *Sclerotinia trifoliorum*, 27 inhibited the fungus growth (Figure 1) and reduce

their development more than 50%. Isolates Soli and Tak were the most preferment *in vitro* and caused growth inhibition of *R. solani* more than 90% Control plates without rhizobia were completely covered by the pathogen mycelium showing no growth inhibition of the fungus. The mean mycelium growth inhibition of the most effective bacterial isolates (Figure 1) revealed that the inhibition was highly significant (P = 0.05). No physical contact was observed between any of the antagonistic bacteria tested and *Sclerotinia trifoliorum*; moreover, an inhibitory halo was observed suggesting the presence of fungistatic metabolites secreted by the bacteria. On the other hand, a change in mycelial color was observed closed to the colony end of *Sclerotinia trifoliorum*, being this one of a darker brown than the one observed at the center of colony (Figure 2).




Figure 1: Effect of *Rhizobium* isolates on *Sclerotinia trifoliorum* growth *in vitro*.

(*) = Percent growth inhibition compared to uninoculated control. Values followed by (*) were significant (P = 0.05) by Duncan’s multiple range test.

Figure 2: Illustration of the *in vitro* effect of *Rhizobium* strains on the development of *Sclerotinia trifoliorum*.

Rhizobium volatile effects on *Sclerotinia trifoliorum* growth in vitro and Phosphorus solubilizing efficiency

As to the effect of *Rhizobium* volatiles on pathogen growth, among 19 isolates tested for volatiles activity, 18 were able to reduce the growth of pathogen. The isolates Tak, Soli, Jem, Morg 2 and Ari 2 seemed to be the most effective, giving more than 40% inhibition after 96h of incubation (Figure 3).

Most tested *Rhizobium* isolates were able to solubilize phosphate. Eighteen isolates produced a halo on tricalcium phosphate agar media plates (Figures 4, 5). Four isolates, Tak, Soli, Morg 2 and Ari2 were positive for inorganic phosphate solubilization, volatiles substances production and generate more than 50% of pathogen inhibition. They were classified the most effective isolates *in vitro* trials.




Figure 3: Ability of the most effective bacterial isolates in dual culture to produce volatiles.




Figure 4: Ability of the most effective bacterial isolates in dual culture to solubilize inorganic phosphorus.

Figure 5: Phosphorus solubilization by *Rhizobium* strains.

Greenhouse experiments

To study the rhizobia antagonism activities, the previous trial was replicate in pot under glass house conditions. The results indicate that Soli, INRAT, Mid, Tak, and Morg2 isolates inhibited significantly sclerotinia stem rot of fenugreek caused by *Sclerotinia triflorum* (Figure 5). Mass Disease expression MDI% was calculated according to Tezcan and Yildiz (26) methodology. Inoculation with Sol isolates induced suppression of sclerotinia stem rot more than 80%, while inoculation with Jbn, Mat1/11, Mat3/17, Ment and Morg1 isolates reduced the disease more than 60% (Figure 6). In

these tests; no bacteria protected the plants completely against *Sclerotinia triflorum*, although all isolates increase significantly in fresh weight compared to the infested control (Figure 7, 8). The inhibition of the fungus growth by some of Rhizobium isolates are varied from 60 to 80% (Figure 5). Control plants not treated with bacteria but inoculated with *Sclerotinia triflorum* alone rendered up to 100% sclerotinia stem rot incidence with the majority of plant completely stunted or dead. The fungus resulted in a pronounced decrease in the fresh weight of the shoots compared to the uninoculated control and to some of the treatments with the most effective bacteria isolates (Figure 7).




Figure 6: Antagonism of *Rhizobium* towards *Sclerotinia triflorum* calculated according MDI (%) of sclerotinia stem rot of fenugreek inoculated by the pathogen and different.




Figure 7: Effect of inoculation by *Rhizobium* strains on shoot weight of fenugreek infected with *Sclerotinia triflorum* under glass house conditions 8 weeks after sowing.




Figure 8: Effect of inoculation by *Rhizobium* strains on root weight of fenugreek infected with *Sclerotinia triflorum* under glass house conditions 8 weeks after sowing.

Shoots fresh weight reduction was about 40% in the plant control inoculated with *Sclerotinia triflorum*, alone. Whereas, shoot dry weight reduction, of plants inoculated with both the pathogen and *Rhizobium* strain Morg1 was 10% (Figure 7).

Rhizobium strains *in vivo* under greenhouse conditions. (Control (-) represents un-inoculated fenugreek plants and control (+) plants infested with *Sclerotinia triflorum*)

Discussion

The ability of rhizobia to inhibit certain soil borne plant pathogens and plant parasites [27-29] has increased the importance of rhizobia besides their use in nitrogen fixation. In this study, an experiment was therefore carried out to study the antagonist effect of thirty two *Rhizobium* strains to *Sclerotinia triflorum*. We tested rhizobia for volatiles and solubilizing phosphorus because previous studies suggested that bacteria possessing these traits can increase plant growth [30]. The reduction in fungal growth observed *in vitro*, by the selected *Rhizobium* isolates, was presumably due to metabolites being released from bacteria into the culture medium. These metabolites may include antibiotics and/or cell-wall degrading enzymes. Different studies have implicated antifungal secondary metabolites produced by *Rhizobium* spp. in the control of plant diseases caused by pathogenic fungi [15].

Our experiments showed that four isolates (Tak, Soli, Morg 2 and Ari2) solubilized inorganic phosphorus and significantly inhibited pathogen growth by producing volatiles.

Greenhouse experiments showed that inoculation of fenugreek with *Sclerotinia triflorum* reduced plant growth and caused sclerotinia stem rot of hyhana variety. Several other workers have noticed the beneficial effects of rhizobia on plant growth and reduction of diseases incidence [29,31,32]. Interestingly, application of *Rhizobium* isolates significantly reduced the mass disease index and increased plant growth (Figure 5). These rhizobia also increased fresh weight of roots and shoots. These benefits may be attributed to better disease control in presence of the bacteria and /or to better nutrition, due especially to higher nodulation and phosphorus uptake [33]. In our study, the basic mechanisms behind such protection is not clearly defined, the possibility that competition, antibiosis, direct parasitism and induced resistance by the antagonistic bacteria, may operate synergistically after inoculation with effective *Rhizobium* strain cannot be ruled out. Currently investigations are being conducted to determine the modes of actions of all the promising isolates from this study. To determine whether these promising strains can be developed into commercial inoculants, their biocontrol efficacy must first be confirmed under field's conditions.

Conclusion

Generally, the use of *Rhizobium* have greatly reduced the sclerotinia stem rot incidence as well as enhanced the growth of fenugreek plants. The selected rhizobia could be used effectively as biocontrol agents of fenugreek against *Sclerotinia trifoliorum*. Nevertheless, further studies are needed to characterize the mechanisms involved in *Rhizobium*-inoculated fenugreek resistance to *S. trifoliorum* to improve *Rhizobium* strains uses as biofertilizer and biocontrol agents in fenugreek fields. The use of bioagents is highly beneficial as environmentally friendly application and can be used as an alternative for fungicides to enhance the plant growth and reduce disease incidence, resulting in higher yield.

Conflict of Interest

The authors declare that they have no financial interest and no conflict of interest.

Bibliography

1. Saada N., et al. "The effect of organic composts on the chemical characteristics of fenugreek (*Trigonella foenum graecum*) from Sudan and Tunisia". *Journal of New Sciences* 11 (2014): 1-8.
2. Rekik I and Bergaoui R. "Influence du fenugrec (*Trigonella foenum graecum* L.) sur la production laitière des lapines et la croissance des lapereaux avant sevrage". *Journal of New Sciences* 30 (2016): 1709-1718.
3. Aissa A., et al. "In vitro anthelmintic activity of some Mediterranean plants against *Haemonchus contortus* infective stage". *Options Méditerranéennes* 107 (2013): 183-187.
4. Hfaiedh., et al. "Beneficial effects of fenugreek seeds (*Trigonella foenum graecum* L) added in the diet of diabetic rats". *Journal of Advances in Chemistry* 8 (2014): 1556-1565.
5. Haouala., et al. "Aqueous and organic extracts of *Trigonella foenum-graecum* L. inhibit the mycelia growth of fungi". *Journal of Environmental Sciences* 20 (2008): 1453-1457.
6. Omezzine., et al. "Variation in phytochemical content and antifungal activity of *Trigonella foenum-graecum* L. with plant developmental stage and ploidy level". *South African Journal of Botany* 92 (2014): 120-125.
7. Mohamed IH., et al. "Evaluation of compost and compost extract efficiency as bio-control agents and damping-off disease incidence of fenugreek (*Trigonella foenum-graecum*)". *Zagazig Journal of Agricultural Research* 40.2 (2013): 239-249.
8. Gargouri., et al. "Occurrence of sclerotinia stem rot of fenugreek caused by *Sclerotinia trifoliorum* and *S. sclerotiorum* in Tunisia". *European Journal of Plant Pathology* 149 (2017): 587-597.
9. Peoples MB., et al. "Enhancing legume N₂ fixation through plant and soil management". *Plant Soil* 147 (1995): 83-101.
10. Carrillo GC and Del Rosario VM. "Comparative study of siderophore like activity of *Rhizobium phaseoli* and *Pseudomonas fluorescens*". *Journal of Plant Nutrition* 15 (1992): 579-590.
11. Arora., et al. "Isolation of siderophores-producing strains of *Rhizobium meliloti* and their biocontrol potential against *Macrophomina phaseolina* that causes charcoal rot of groundnut". *Current Science* 81 (2001): 673-677.
12. Essalmani H and Lahlou H. "In vitro antagonistic activity of some microorganisms towards *Fusarium oxysporum* f.sp. *lentis*". *Cryptogamie-Mycologie* 23 (2002): 221-234.
13. Chakraborty U and Purkayastha RP. "Role of Rhizobitoxine in protecting soybean *Glycine-Max* roots from *Macrophomina phaseolina* infection". *Canadian Journal of Microbiology* 30 (1984): 285-289.
14. Ehteshamul-Haque S and Ghaffar A. "Use of rhizobia in the control of root rot diseases of sunflower, okra, soybean and mungbean". *Journal of Phytopathology* 138 (1993): 157-163.
15. Siddiqui IA., et al. "Greenhouse evaluation of rhizobia as bio-control agent of root infecting fungi in Okra". *Acta Agrobotanica* 53.1 (2000): 13-22.

16. Siddiqui ZA and Mahmoud I. "Effects of rhizobacteria and root symbionts on the reproduction of *Meloidog vanica* and growth of chickpea". *Bioresource Technology* 79 (2001): 41-45.
17. Abdelaziz RA., et al. "Biocontrol of *faba bean* root-rot *corrhizae* and its effect on biological nitrogen fixation". *Egyptian Journal of Microbiology* 31 (1996): 273-286.
18. Vincent JM. "A manual for the practical study of the root nodule bacteria", I.B.P. Handbook No.15. Blackwell Scientific Publications, Oxford. (1970): 44.
19. Beck DP, et al. "Practical Rhizobium- Legume Technology Manual". ICARDA, Aleppo, S (1993).
20. Paulitz TC., et al. "Insights into the prevalence and management of soil borne cereal pathogens under direct seedling in the Pacific Northwest, U.S.A". *Canadian Journal of Plant Pathology*. 24 (2002): 416-428.
21. Landa BB., et al. "Antagonistic activity of bacteria from the chickpea rhizosphere against *Fusarium oxysporum* f.sp. *ciceris*". *Phytoparasitica* 25 (1997): 305-318.
22. Fiddman PJB and Rossall S. "The production of antifungal volatiles by *Bacillus subtilis*". *Journal of Applied Bacteriology* 47 (1993): 119-126.
23. Pikovskaya RI. "Mobilisation of phosphorus in soil in connection with the vital activity of some microbial species". *Microbiology* 17 (1948): 362-370.
24. Afzal A and Bano A. "*Rhizobium* and phosphate solubilizing bacteria improve the yield and phosphorus uptake in wheat (*Triticum aestivum* L.)". *International Journal of Agriculture and Biology* 10 (2008): 85-88.
25. Yousefi A., et al. "Phosphate solubilizing bacteria and arbuscular mycorrhizal fungi impacts on inorganic phosphorus fractions and wheat growth". *World Applied Sciences Journal* 15 (2011): 1310-1318.
26. Tezcan H and Yildiz M. "Ege Bolgesinde bazi toprak kaynakli funguslarin neden oldugu kavun kurumalan Uzerinde arastirmalar VI". *Turkiye Fitopatoloji Kongresi* 7-11 Ekim, zmir (1991): 121-124.
27. Arfaoui A., et al. "Biochemical analysis of chickpea protection against *Fusarium* wilt afforded by two *Rhizobium* isolates". *Plant Pathology Journal* 4 (2005): 35-42.
28. Mabrouk Y., et al. "Some compatible *Rhizobium leguminosarum* strains in peas decrease infections when parasitised by *Orobanche crenata*". *Weed Research* 47 (2007): 44-53.
29. Hemissi., et al. "Enhanced Defence Responses of Chickpea Plants Against *Rhizoctonia solani* by Pre-Inoculation with Rhizobia". *Journal of Phytopathology* 161 (2013): 412-418.
30. Glick BR., et al. "A novel procedure for rapid isolation of plant growth promoting *Pseudomonas*". *Canadian Journal of Microbiology* 41 (1995): 533-536.
31. Arfaoui A., et al. "Identification of *Rhizobium* isolates possessing antagonistic activity against *Fusarium oxysporum* F. sp. *ciceris*, the causal agent of *Fusarium* wilt of chickpea". *Journal of Plant Pathology* 88 (2006): 67-75.
32. Hemissi I., et al. "Effects of some *Rhizobium* strains on chickpea growth and biological control of *Rhizoctonia solani*". *African Journal of Microbiology Research* 5 (2011): 4080-4090.
33. Alagawadi AR and Gaur AC. "Associative effect of *Rhizobium* and phosphate-solubilizing bacteria on the yield and nutrient uptake of chickpea". *Plant Soil* 105 (1988): 241-246.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667