



In Vitro Screening of Bioactive Endophytic Microorganisms Isolated from Medicinal Flora of the Himalayan Region

Sharma R, Chamoli S, Negi N, Kumari R, Neha N, Aishwarya A, Chauhan S and Devi S*

Department of Microbiology, Himachal Pradesh University Summerhill, Shimla (171005), India

*Corresponding Author: Devi S, Department of Microbiology, Himachal Pradesh University Summerhill, Shimla (171005), India.

Received: January 08, 2025

Published: January 27, 2025

© All rights are reserved by

Devi S, et al.

Abstract

Plant growth and its defence mechanism have been facilitated by endophytic microorganism, in symbiotic association with bacteria found in various plant sections. This study, focuses on the isolation and characterization of endophytes utilizing different medicinal herbs like *Achyranthes bidentata*, *Asparagus racemosus*, *Centella asiatica*, *Withania somnifera*, and *Viola odorata*, *Urtica dioica*, *Eupatorium adenophorum*, *Carica papaya*, *Thuja occidentalis*, and *Cinnamomum tamala* that are well known for their therapeutic efficacy. Endophytic microorganisms were isolated, out of which some were bacterial and some were fungal species. These endophytic organisms were further screened for their antimicrobial, antioxidant, and plant growth-promoting activities. Among fungi, *Arthrinium* sp. from *Cinnamomum tamala* exhibited maximum free radical scavenging activity of 93.95%, while *Alternaria* sp. showed the minimum of 77.32% free radical scavenging activity. In antifungal assays, *Alternaria* sp. and *Fusarium* sp. inhibited *Aspergillus niger* and *Penicillium notatum*, while *Ulocladium* sp. inhibited *Rhizopus* sp. In antibacterial tests, *Bipolaris* sp. from *Thuja occidentalis* showed the largest zone of inhibition (32 mm), while *Alternaria* sp. showed the smallest (9 mm). Minimum inhibitory concentration (MIC) assays revealed that the fungi ethyl acetate extracts of the effectively inhibited pathogenic bacteria, with MIC values as low as 0.625 mg/100 µl. These insights reveal the potential of endophytic microorganisms in sustainable agriculture and therapeutic development, offering promising leads for combating antibiotic-resistant pathogens.

Keywords: Antibiotic Resistance; Endophytes; Medicinal Plants; Antioxidant

Introduction

Numerous microbial populations, including bacteria and fungus, have been identified as endophytes and linked to the interior tissues of plants. De Bary (1866) first used the word "endophyte" to describe the presence of microorganisms within infected plant tissues that do not negatively impact the host plant [1]. Almost all the plants have been found to be infested with one or more endophytes [2]. By releasing beneficial secondary metabolites, these endophytes shield their hosts against pathogens and unfavorable circumstances [3-4].

Endophytic fungi, in particular, are recognized for their critical physiological [5] and ecological [6] contributions to plant life.

They assist in nutrient acquisition, enhance the plant's stress tolerance, and modulate the host's defence mechanisms. These fungi and bacteria synthesize a diverse range of bioactive compounds, including antimicrobial agents effective against plant pathogens, insect- and pest-repellent molecules, phytohormones that promote plant growth, and metabolites that enable plants to withstand abiotic stresses such as drought, salinity, and temperature extremes [7]. Moreover, endophytes possess the ability to produce unique secondary metabolites with significant potential for application in various fields, including agriculture, pharmaceuticals, and biotechnology [8]. For instance, in agriculture, these metabolites can be utilized as natural biopesticides and biofertilizers to promote

sustainable farming practices. In the pharmaceutical sector, endophytes have emerged as a promising source of novel antibiotics, anticancer compounds, and other therapeutic agents [9]. The ecological importance of endophytes extends to their role in maintaining biodiversity, improving soil health, and facilitating ecosystem stability. Endophytes represent a diverse group of microorganisms with immense potential for contributing to plant health and productivity, as well as offering innovative solutions for challenges in agriculture, medicine, and environmental sustainability [10]. The ongoing exploration of endophytic diversity and their metabolic capabilities is crucial for unlocking their full potential. In the present study, *Urtica dioica*, *Eupatorium adenophorum* (also known as *Ageratina adenophora*), *Carica papaya*, *Thuja occidentalis*, *Cinnamomum tamala*, *Achyranthes bidentata*, *Asparagus racemosus*, *Centella asiatica*, *Withania somnifera* and *Viola odorata*, are some of the valuable curative herbs that were screened. The main objectives are the isolation and identification of endophytes (fungi and bacteria), fermentation and extraction of metabolites (ethyl acetate) from isolated fungal isolates and to investigate *in vitro* antioxidant efficacy of endophytic fungal ethyl acetate extracts by DPPH free radical scavenging method and antimicrobial efficacy of endophytic fungal and bacterial isolates by dual culture, agar well diffusion and minimum inhibitory concentration method.

Material and Methods

Chemical and reagent

All the chemical and reagent utilized in the current analysis were of high purity and analytical grade (AR/GR), and they were collected from Hi-Media Pvt. Ltd. Mumbai, E-Merck, and others companies.

Test organisms

In the present work, test microorganisms were used *i.e.*, Bacterial (*P. aeruginosa*, *S. aureus*, *B. cereus*, *E. coli*, *S. typhi*, *S. flexneri*, *K. pneumoniae*) and fungal strains (*Aspergillus* sp., *Penicillium* sp., *Rhizopus* sp.).

Sample collection

In this study, the roots, stem and leaves of the plants *viz.*, *Urtica dioica*, *Eupatorium adenophorum* (also known as *Ageratina adenophora*), *Carica papaya*, *Thuja occidentalis*, *Cinnamomum tama-*

la, *Achyranthes bidentata*, *Asparagus racemosus*, *Centella asiatica*, *Withania somnifera* and *Viola odorata* were collected from different areas of Himachal Pradesh and Uttarakhand.

Isolation of endophytic microorganisms

For the isolation of endophytes, healthy plant parts were washed in running tap water. A sterile cutter and forceps were used to chop the leaves into little pieces. Following that, each segment was put on nutrient agar (NA) and potato dextrose agar (PDA) media and then the plates were incubated at 37°C. After inoculation period, different fungal and bacterial strains were isolated by subculturing the hyphal tips onto a fresh PDA and nutrient agar media [11].

Identification of isolated endophytic microorganisms

Isolated endophytic bacterial isolates were identified by gram staining and different biochemical tests *viz.*, methyl red, indole test, Voges Proskauer, oxidase test, citrate test catalase test (Figure 1), motility test and fermentation of different sugars for the confirmation of isolate [12,13] and for fungal isolates by Lactophenol picric acid [14].

Preservation of isolated endophytic microorganisms

Pure endophytic fungal and bacterial isolates were then transferred separately to PDA and nutrient agar slants [15]. Growths of endophytic bacterial and fungal strains were observed for 24 hours and 72 hours respectively and were maintained at 4°C till further use [16].

Fermentation and extraction of metabolites from endophytic microorganisms *i.e.*, bacterial extract and fungal ethyl acetate extract

The cultivation of isolated fungal endophytes was conducted in conical flasks containing a sterilized rice medium. The rice was pre-soaked in sterile water within a glass container [17]. After cooling, the solidified rice medium was aseptically inoculated with pure mycelial cultures of the isolated endophytic fungi. The resulting supernatant was evaporated to dryness, yielding crude fungal extracts [18]. Similarly, all endophytic bacterial strains were cultured in nutrient broth, followed by centrifugation of each bacterial culture. The resulting supernatant was collected and stored in collection tubes as bacterial extracts [19].

In Vitro antioxidant activity of endophytic fungal ethyl acetate extract

The free radical scavenging activity of fungal extracts was assessed using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay. Fungal extracts and a positive control (DPPH dissolved in distilled water) were prepared in methanol and then combined with DPPH in methanol [20]. A mixture of methanol and DPPH without the fungal extract was used as the control. The experiment was conducted in triplicate, and the inhibition of DPPH was expressed as a percentage of radical scavenging activity at different extract concentrations [21].

In Vitro antifungal screening of endophytic fungal isolates by dual culture method

Ethyl acetate extracts of endophytic fungal isolates were evaluated for antifungal activity against standard pathogenic fungal strains, including *Aspergillus* sp., *Penicillium* sp., and *Rhizopus* sp., using the dual culture method. Potato dextrose agar (PDA) was utilized as the culture medium [22]. Mycelial plugs of both the pathogens and endophytic fungi were placed at a specific distance from each other on PDA plates. The incubation of each endophytic fungal isolate against pathogenic strains were observed by the growth pattern of fungus by comparing with controls [23].

In Vitro Antibacterial Activity of endophytes

The activity of endophytic ethyl acetate fungal extracts against various pathogenic bacteria was evaluated using the agar well diffusion method. Additionally, the minimum inhibitory concentration (MIC) of the endophytic bacterial extracts was determined using the resazurin dye assay [24].

Antibacterial activity of endophytic fungal ethyl acetate extract by agar well diffusion method

The antimicrobial compounds in the ethyl acetate extract of endophytic fungi were allowed to diffuse into the agar medium, interacting with freshly inoculated test bacteria, including *P. aeruginosa*, *S. aureus*, *B. cereus*, *S. typhi*, *S. flexneri*, and *E. coli*. The inhibition zones formed around the wells were measured using a diameter scale, as described by [25]. Each extract was tested against these pathogenic bacterial strains, along with positive controls, including chloramphenicol, amoxicillin, clindamycin, azithromycin, ciprofloxacin, azlocillin, and imipenem. The diameter of the inhibition zones was recorded following the method of [26,27].

Evaluation of minimum inhibitory concentration (MIC) using the resazurin dye assay

MIC is the concentration of the higher dilution in which the absence of bacterial growth occurs. Sterile microtitre plates (Tarson) were used. Rows were filled with Muller Hinton broth. Fungal extracts showing maximum activity in first column and further 2-fold serial dilution was achieved. Now 10µl of different bacterial suspension were added in each row separately of the microtitre plate [28,29]. Now 1µl of Resazurin dye was added in the entire plate as an indicator dye. Positive and negative controls were also used. Furthermore, a shift in color from purple to pink or a loss of color (colorless) was interpreted as a negative outcome [30].

Result and Discussion

Isolation of endophytes from various plants

Different endophytic microorganisms were isolated from stem, leaves and roots of different plants (Table 1) viz., *Urtica dioica*, *Eupatorium adenophorum* (leaf, stem, root), *Carica papaya*, *Thuja occidentalis*, and *Cinnamomum tamala* which were collected from different district of Himachal Pradesh (Shimla, Solan, Bilaspur), and Uttarakhand (Dehradun, Rudraprayag), *Achyranthes bidentata*, *Asparagus racemosus*, *Centella asiatica*, *Withania somnifera* and *Viola odorata* from H.P.

Identification of Endophytic microorganism isolated from different medicinal plants

The identification of endophytic fungi was done by Lactophenol picric acid and bacteria by gram staining, further isolated bacterium was identified by using different biochemical tests viz., indole test, methyl red, Voges Proskauer, citrate test, oxidase, catalase, motility test and fermentation of different sugars for the identification of isolates. In present study the isolated endophytic fungi were identified on the basis of colony morphology, microscopic examination-like shapes of conidia, mycelium and hyphae and endophytic bacteria were identified by gram staining and by different sugar tests.

Preservation of isolated endophytic microorganisms

The isolated endophytic isolates were preserved by slant preparation method in refrigerator at 4°C on PDA for fungal isolates and Nutrient agar for bacterial isolates (Figure 2).

Table 1: Endophytic microorganisms isolated from different medicinal plants.

Sr. No.	Medicinal plant	Location	Types of isolates	No. of isolates
1.	<i>Urtica dioica</i>	Himachal Pradesh	Bacterial	3
			Fungal	1
		Uttarakhand	Bacterial	1
			Fungal	1
2.	<i>Eupatorium adenophorum</i>	Himachal Pradesh	Bacterial	2
			Fungal	1
		Uttarakhand	Bacterial	2
			Fungal	1
3.	<i>Carica papaya</i>	Himachal Pradesh	Bacterial	-
			Fungal	-
		Uttarakhand	Bacterial	1
			Fungal	-
4.	<i>Thuja occidentalis</i>	Himachal Pradesh	Bacterial	1
			Fungal	1
		Uttarakhand	Bacterial	1
			Fungal	2
5.	<i>Cinnamomum tamala</i>	Himachal Pradesh	Bacterial	2
			Fungal	1
		Uttarakhand	Bacterial	1
			Fungal	-
6.	<i>A. bidentata</i>	Himachal Pradesh	Bacterial	3
			Fungal	2
7.	<i>A. racemosus</i>	Himachal Pradesh	Bacterial	2
			Fungal	-
8.	<i>C. asiatica</i>	Himachal Pradesh	Bacterial	2
			Fungal	-
9.	<i>W. somnifera</i>	Himachal Pradesh	Bacterial	2
			Fungal	2
10.	<i>V. odorata</i>	Himachal Pradesh	Bacterial	1
			Fungal	1
Total			Bacterial	24
			Fungal	13

Fermentation and extraction of metabolites from isolated endophytic fungal ethyl acetate extract

All the isolated endophytic fungal isolates were used for the extraction of fungal metabolites which were extracted by using ethyl acetate. Whole process took 9-10 days. The endophytic ethyl acetate extract was then dissolved in Dimethyl sulfoxide (DMSO) (Figure 3).

In vitro Antioxidant Activity of isolated Endophytic Fungal ethyl acetate extracts

All the isolated endophytic fungal ethyl acetate extract were having antioxidant activity where they showed ability to reduce free radicals. In contrast with the current investigation, Selim and coworkers reported that *Ulocladium* sp. isolated from *Pulicaria undulate* showed 14% antioxidant activity and *Ulocladium* sp. isolated from *Galium sanaicum* don't showed any activity [31].



Figure 1: Results of different biochemical tests of isolated Endophytic bacterial isolates.



Figure 2: Slants of isolated endophytic bacterial and fungal isolates.



Figure 3: (a) Growth of isolated endophytic fungal isolates in solid rice media (b) different isolated endophytic fungal ethyl acetate extracts.




Figure 4: DPPH Radical Scavenging Activity of isolated endophytic fungal ethyl acetate extracts.

Table 2: Antioxidant activity (%) of different isolated endophytic fungal ethyl acetate extracts.

Sr. No.	Sample (Endophytic fungal ethyl acetate extract)	Antioxidant Activity (%)
1.	<i>Arthrinium</i> sp.	93.95
2.	<i>Ulocladium</i> sp.	88.89
3.	<i>Fusarium</i> sp.	91.82
4.	<i>Alternaria</i> sp.	86.75
5.	<i>Verticillium</i> sp.	92.71
6.	<i>Bipolaris</i> sp.	89.38
7.	<i>Chrysosporium</i> sp.	88.84
8.	<i>Geomyces</i> sp.	92.33
9.	<i>Alternaria</i> sp. (ABS)	79.96
10.	<i>Alternaria</i> sp. (ABL)	77.32
11.	<i>Fusarium</i> sp. (VOR)	80.83
12.	<i>Alternaria</i> sp. (WSR)	77.72
13.	<i>Ulocladium</i> sp. (WSS)	83.04

Anti-fungal screening of endophytic fungal isolates by dual culture method

The isolated endophytic fungal isolates were tested against 3 pathogenic fungi viz. *Penicillium notatum*, *Rhizopus stolonifer* and *Aspergillus niger*. In the present study isolated endophytic fungi *Verticillium* sp., *Chrysosporium* sp., *Bipolaris* sp., and *Fusarium* sp. showed antagonism against pathogenic fungus *Aspergillus niger* and *Penicillium notatum* were as *Arthrinium* sp., *Alternaria* sp. showed antagonism against *Aspergillus niger* and *Ulocladium* sp. against *Penicillium notatum*. *Alternaria* sp. from *A. bidentata* and *W. somnifera*, *Fusarium* sp. from *V. odorata* showed antifungal activity against *Aspergillus niger* but *Ulocladium* sp. from *W. somnifera* was unable to inhibit *Aspergillus niger*. In case of *Penicillium nota-*

tum, *Alternaria* sp. from *A. bidentata* and *W. somnifera*, *Fusarium* sp. from *V. odorata* but *Ulocladium* sp. from *W. somnifera* donot showed any inhibitory activity. In case of *Rhizopus stolonifer*, *Alternaria* sp. from *W. somnifera*, *Ulocladium* sp. of *W. somnifera* and *Fusarium* sp. of *V. odorata* showed antagonistic effect but *Alternaria* sp. of *A. bidentata* does not.

Antibacterial activity of endophytic microorganisms: Antibacterial activity of endophytic fungal ethyl acetate extract by agar well diffusion method

In the present study antibacterial activity of isolated fungal ethyl acetate extracts and endophytic bacterial isolate were checked against various pathogenic microorganisms viz., *Bacillus cereus*, *E.*

coli, *Staphylococcus aureus*, *Shigella flexneri*, *Salmonella typhi* and *Pseudomonas aeruginosa* by agar well diffusion method on Muller Hinton Agar. Positive controls were also run (Chloramphenicol, Amoxicillin, Clindamycin, Azithromycin, Ciproflaxin, Azlocillin and Imipenem). *Bacillus cereus*, *E. coli*, *Staphylococcus aureus*, *Shigella flexneri*, *Salmonella typhi* and *Pseudomonas aeruginosa* on Muller Hinton Agar along with positive controls (Chloramphenicol, Amoxicillin, Clindamycin, Azithromycin, Ciproflaxin, Azlocillin and imipenem) and negative control (Sterile nutrient broth). The maximum zone of inhibition was shown by *Bipolaris* sp. isolated from *Thuja*

occidentalis i.e. 32mm against *Shigella flexneri* and minimum zone of inhibition was shown by *Ulocladium* sp. and *Fusarium* sp. isolated from *Urtica dioica* (Uk) and *Eupatorium adenophorum* (UK and HP) i.e. 10 mm against *S. typhi* and *P. aeruginosa*. (Figure 5, 6 and 7 and Table 3 and 4). Similar type of investigation was done where Enany and coworkers reported that *Ulocladium* sp. isolated from *Mentha piperita* showed 8mm zone against *Pseudomonas aeruginosa* isolated from infected freshwater fishes which is lower than the present study [32].

Table 3: Zone of inhibition measured in millimeter against pathogenic bacterial strains.

Zone of inhibition (ZOI) in millimeter						
Bacterial strains	<i>Bacillus cereus</i>	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Shigella flexneri</i>	<i>Salmonella typhi</i>	<i>Pseudomonas aeruginosa</i>
Zone of inhibition of positive controls	Chloram phenicol	Amoxicillin	Clindamycin	Azithromycin	Ciproflaxin	Aziocillin
	30 mm	24 mm	28 mm	33 mm	25 mm	30 mm
Zone of inhibition (ZOI) of isolated endophytic fungal ethyl acetate extract and endophytic bacterium						
<i>Verticillium</i> sp.	16 mm	11 mm	20 mm	26 mm	13 mm	19 mm
<i>Chrysosporium</i> sp.	-	-	14 mm	15 mm	-	-
<i>Bipolaris</i> sp.	23 mm	17 mm	20 mm	32 mm	20 mm	23 mm
<i>Arthrinium</i> sp.	17 mm	12 mm	19 mm	23 mm	-	12 mm
<i>Fusarium</i> sp.	13 mm	-	22 mm	24 mm	-	10 mm
<i>Alternaria</i> sp.	19 mm	11mm	22 mm	25 mm	13 mm	21 mm
<i>Ulocladium</i> sp.	23 mm	-	20 mm	28 mm	10 mm	16 mm
<i>Geomyces</i> sp.	28 mm	17 mm	24 mm	30 mm	20 mm	28 mm
<i>Bacillus acidicer</i>	-	24 mm	23 mm	-	23 mm	22 mm
<i>Paenibacillus apiaries</i>	-	20 mm	23 mm	-	21 mm	21 mm




Figure 5: Activity of Endophytic fungal ethyl acetate extract against different pathogenic bacteria on the basis of ZOI formation.




Figure 6: Activity of Endophytic bacterium against different pathogenic bacteria on the basis of ZOI formation.

Table 4: Zone of inhibition of isolated Endophytic fungal ethyl acetate extract and Bacterial extracts measured in millimeter against pathogenic bacterial strains.

Zone of inhibition (ZOI)						
Bacterial strains	<i>B. cereus</i>	<i>E. coli</i>	<i>S. aureus</i>	<i>S. lexineri</i>	<i>S. typhi</i>	<i>P. aeruginosa</i>
Zone of inhibition (ZOI) of positive controls	Chloramphenicol	Amoxicillin	Clindamycin	Azithromycin	Ciproflaxin	Aziocillin
	27 mm	24 mm	19 mm	16 mm	25 mm	22 mm
Zone of inhibition (ZOI) of endophytic fungal ethyl acetate extract and bacterial extract						
<i>Alternaria</i> sp.	15 mm	14 mm	14 mm	12 mm	12 mm	15 mm
<i>Alternaria</i> sp.	14 mm	12 mm	10 mm	14 mm	9 mm	10 mm
<i>Ulocladium</i> sp.	20 mm	11 mm	13 mm	29 mm	16 mm	17 mm
<i>Bacillus</i> sp.	11 mm	-	-	11 mm	10 mm	10 mm
<i>Micrococcus</i> sp.	-	10 mm	14 mm	12 mm	10 mm	12 mm
<i>Serratia</i> sp.	-	-	15 mm	10 mm	10 mm	10 mm




Figure 7: Activity of endophytic fungal ethyl acetate extract and endophytic bacterial extracts against different pathogenic bacteria on the basis of ZOI formation.

Determination of minimum inhibitory concentration (MIC) by Resazurin dye

All the endophytic fungal ethyl acetate extract and bacterial extract were checked for their MIC by Resazurin dye method. Enany and coworkers reported that *Ulocladium* extract was not effective

against pathogenic bacteria which was isolated from *Mentha piperita* [32], but in this study *Ulocladium* sp. was found more effective against all pathogenic bacterial strains (Table 5, 6 Figure 8).

Conc. of extract Mg/100µl	<i>S. aureus</i>		<i>E. coli</i>		<i>B. cereus</i>		<i>S. typhi</i>		<i>S. flexneri</i>		<i>P. aeruginosa</i>	
	Paenibacillus apiarius	Bacillus acidicereler	Paenibacillus apiarius	Bacillus acidicereler	Paenibacillus apiarius	Bacillus acidicereler	Paenibacillus apiarius	Bacillus acidicereler	Paenibacillus apiarius	Bacillus acidicereler	Paenibacillus apiarius	Bacillus acidicereler
10	-	-	-	-	-	-	-	-	-	-	-	-
5	-	-	-	-	-	-	-	-	-	-	-	-
2.5	-	-	-	-	-	-	-	-	-	-	+	-
1.25	+	+	+	-	-	-	+	-	+	-	+	-
0.625	+	+	+	+	+	-	+	+	+	-	+	+
0.312	+	+	+	+	+	+	+	+	+	+	+	+
0.156	+	+	+	+	+	+	+	+	+	+	+	+
0.078	+	+	+	+	+	+	+	+	+	+	+	+
0.039	+	+	+	+	+	+	+	+	+	+	+	+
0.0195	+	+	+	+	+	+	+	+	+	+	+	+
MIC (mg/100µL)	2.5	2.5	2.5	1.25	1.25	0.625	2.5	1.25	1.25	0.625	5	1.25

- = no bacterial growth, + = bacterial growth
 Paenibacillus apiarius Bacillus acidicereler

Table 5: MIC of ethyl acetate extract of Endophytic bacterial Isolates against Bacterial pathogens.

Conc. of extract mg/100µl	<i>B. cereus</i>					<i>E. coli</i>					<i>P. aeruginosa</i>					<i>S. aureus</i>					<i>S. flexneri</i>					<i>S. typhi</i>																
	Ulocladium sp.					Serratia sp.					Micrococcus sp.					Bacillus sp.					Alternaria sp. (A. bidentata)					Alternaria sp. (W. somnifera)																
10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
2.5	+	-	-	-	-	+	+	+	+	+	+	+	+	+	+	+	+	-	-	+	+	+	+	+	+	+	+	+	-	-	+	+	+	+	+	+	+	+	+	+	+	+
1.25	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
0.625	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
0.312	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
0.156	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
0.078	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
0.039	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
0.0195	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
MIC(mg/100µl)	5	2.5	2.5	2.5	2.5	5	2.5	2.5	5	5	5	5	2.5	2.5	5	5	5	5	2.5	2.5	5	5	5	2.5	2.5	1.2	5	5	5	5	5	2.5	2.5	2.5	2.5	5	5	5				

- = no bacterial growth, + = bacterial growth
 Alternaria sp. (A. bidentata) Alternaria sp. (W. somnifera) Ulocladium sp. Bacillus sp. Micrococcus sp. Serratia sp.

Table 6: MIC of ethyl acetate extract of endophytic Fungal isolate against Bacterial pathogens.




Figure 8: Determination of MIC of different endophytic fungal ethyl acetate extract against various pathogenic bacterial strains (a) *Alternaria* sp. (b) *Ulocladium* sp. (c) *Bacillus* sp. (d) *Serratia* sp.

Conclusion

The current research findings on endophytic isolates indicate that they can be used to produce bioactive compounds for pharmacological purposes without affecting human beings. Seasonal variations play a key role in plant's antimicrobial and antioxidant properties. Phytoconstituents are the constituent molecules which helps plants in exhibiting bioactive potential. Phenolic compounds and saponin levels appear to be higher in winter and autumn season. The availability and concentration of phytochemicals in plants are affected by seasonal fluctuations. These changes in phytochemicals have been widely attributed to variations in environmental variables or abiotic factors such as composition of soil, responses against a variety of season specific pathogens and changes in temperature and rainfall. Biotic factors such as development of the plant and plant genotype also change rapidly and these too may affect the composition of the plant parts. These variations seem to influence the harvest time which ensure optimum quality and yields of medicinal plants. From the Current research work, it can be concluded that endophytic microbes can be a good source for herbal formulation. These organisms can be harmless and can be used to produce bioactive compounds for pharmaceutical use. The obtained results indicates that Endophytic organisms can be a good source for obtaining biofertilizers, biocontrol agents and antioxidants agents. Therefore, the plants from Uttarakhand exhibited good antimicrobial potential in comparison to the plants from Himachal Pradesh due to the environmental variations such as composition of soil, temperature of that particular area, rainfall, microflora of plant's soil, plant's genotype etc.

Bibliography

1. Gouda Sushanto., *et al.* "Endophytes: a treasure house of bioactive compounds of medicinal importance". *Frontiers in Microbiology* 7 (2016): 1538.
2. Grabka Rachel., *et al.* "Fungal endophytes and their role in agricultural plant protection against pests and pathogens". *Plants* 11.3 (2022): 384.
3. Elhamouly Neveen Atta., *et al.* "The hidden power of secondary metabolites in plant-fungi interactions and sustainable phytoremediation". *Frontiers in Plant Science* 13 (2022): 1044896.
4. Pandey Praveen., *et al.* "Deciphering the mechanisms, hormonal signalling, and potential applications of endophytic microbes to mediate stress tolerance in medicinal plants". *Frontiers in Plant Science* 14 (2023): 1250020.
5. Chaubey Anand Kumar., *et al.* "Induction of Stress Tolerance in Plants by Metabolic Secretions of Endophytes for Sustainable Development". *Microbial Bioactive Compounds: Industrial and Agricultural Applications*. Cham: Springer Nature Switzerland, (2023): 225-248.
6. Santoyo Gustavo., *et al.* "Trichoderma and Bacillus multifunctional allies for plant growth and health in saline soils: recent advances and future challenges". *Frontiers in Microbiology* 15 (2024): 1423980.
7. Selvasekaran Pavidharshini and Ramalingam Chidambaram. "Agriculturally important fungi for crop protection". *Agriculturally Important Fungi for Sustainable Agriculture: Volume 2: Functional Annotation for Crop Protection* (2020): 1-53.

8. Chaudhary Parul, *et al.* "Endophytes and their potential in biotic stress management and crop production". *Frontiers in Microbiology* 13 (2022): 933017.
9. Anand Kumar, *et al.* "Bioactive molecules from fungal endophytes and their applications in pharmaceutical industries: challenges and future scope". *Journal of Basic Microbiology* 63.7 (2023): 690-708.
10. Muhammad Murad, *et al.* "Harnessing bacterial endophytes for environmental resilience and agricultural sustainability". *Journal of Environmental Management* 368 (2024): 122201.
11. Huang Rong, *et al.* "Antagonistic effects of endophytic fungi from *Camellia reticulata* pedicels on yeasts: implications for antimicrobial mechanism of nectar". *Frontiers in Plant Science* 15 (2024): 1494855.
12. Abedinzadeh Motahhareh, *et al.* "Characterization of rhizosphere and endophytic bacteria from roots of maize (*Zea mays* L.) plant irrigated with wastewater with biotechnological potential in agriculture". *Biotechnology Reports* 21 (2019): e00305.
13. Ek-Ramos María J, *et al.* "Bioactive products from plant-endophytic gram-positive bacteria". *Frontiers in Microbiology* 10 (2019): 463.
14. Dhale D A, *et al.* "Production of IAA by Endophytic Fungi as *Phyllanthus emblica* Linn". *Current Agriculture Research Journal* 11.3 (2023).
15. Sallam Nashwa, *et al.* "Endophytic fungi associated with soybean plants and their antagonistic activity against *Rhizoctonia solani*". *Egyptian Journal of Biological Pest Control* 31 (2021): 1-9.
16. Mahfooz Mohd, *et al.* "Evaluation of antifungal and enzymatic potential of endophytic Fungi isolated from *Cupressus torulosa* D. Don". *International Journal of Current Microbiology and Applied Sciences* 67 (2017): 4084-4100.
17. Malhadas Cynthia, *et al.* "Antimicrobial activity of endophytic fungi from olive tree leaves". *World Journal of Microbiology and Biotechnology* 33 (2017): 1-12.
18. El-Nahas Gehad A, *et al.* "Scrutinizing harsh habitats endophytic fungi and their prospective effect on water-stressed maize seedlings". *International Microbiology* (2024): 1-17.
19. Singh Ritu, *et al.* "Isolation and characterization of endophytes bacterial strains of *Momordica charantia* L. and their possible approach in stress management". *Microorganisms* 10.2 (2022): 290.
20. Kusmiati Kusmiati, *et al.* "Profile and in silico analysis of metabolite compounds of the endophytic fungus *Alternaria alternata* K-10 from *Drymoglossum piloselloides* as antioxidants and antibacterials". *Heliyon* 10.6 (2024).
21. Khalil Doaa, *et al.* "Phytochemical screening and antioxidant potential of endophytic fungi isolated from *Hibiscus sabdariffa*". *Journal of Applied Biotechnology Reports* 7.2 (2020): 116-124.
22. Shehata Hanan R, *et al.* "Relevance of in vitro agar based screens to characterize the anti-fungal activities of bacterial endophyte communities". *BMC Microbiology* 16 (2016): 1-7.
23. Colombo Elena Maria, *et al.* "Selection of an endophytic *Streptomyces* sp. strain DEF09 from wheat roots as a biocontrol agent against *Fusarium graminearum*". *Frontiers in Microbiology* 10 (2019): 2356.
24. Zihad SM, *et al.* "Isolation and characterization of antibacterial compounds from *Aspergillus fumigatus*: an endophytic fungus from a mangrove plant of the Sundarbans". *Evidence-Based Complementary and Alternative Medicine* 2022.1 (2022): 9600079.
25. Snoussi Mejdi, *et al.* "Chemical composition and antibiofilm activity of *Petroselinum crispum* and *Ocimum basilicum* essential oils against *Vibrio* spp. strains". *Microbial Pathogenesis* 90 (2016): 13-21.
26. Gonelimali Faraja D, *et al.* "Antimicrobial properties and mechanism of action of some plant extracts against food pathogens and spoilage microorganisms". *Frontiers in Microbiology* 9 (2018): 1639.
27. F Ahmad Abeer and Heba A Shehta. "Assessment of the effects of different extraction methods on the phytochemicals, antimicrobial and anticancer activities of *Eruca sativa* extracts". *Novel Research in Microbiology Journal* 4.3 (2020): 825-844.
28. Santra Hiran K and Debdulal Banerjee. "Broad-spectrum antimicrobial action of cell-free culture extracts and volatile organic compounds produced by endophytic fungi *Curvularia eragrostidis*". *Frontiers in Microbiology* 13 (2022): 920561.

29. Caruso Daniel. "Endophytic Fungi of Native Australian Eremophila Plants as a Source of Antibacterial Compounds". Diss. Swinburne, (2024).
30. Elshikh Mohamed S., *et al.* "Diclofenac removal from the wastewater using activated sludge and analysis of multi-drug resistant bacteria from the sludge". *Environmental Research* 208 (2022): 112723.
31. Selim Khaled A., *et al.* "Antiviral and antioxidant potential of fungal endophytes of Egyptian medicinal plants". *Fermentation* 4.3 (2018): 49.
32. Enany Mohamed E., *et al.* "Antibacterial Activity of Bioactive Compounds from Endophytic Fungi against *P. aeruginosa* isolated from Freshwater Fishes". *Egyptian Journal of Aquatic Biology and Fisheries* 26 (2022): 469-490.