



The Effect of Liming on Carbon Mineralization in Acid Soils

S Suvana^{1,3*}, P Jha², Muralidhar M¹, Indoria AK³ and Anusha E³

¹ICAR- Central Institute of Brackish water Aquaculture, India

²ICAR- Indian Institute of Soil Science, Bhopal, India

³ICAR- Central Research Institute for Dryland Agriculture, Hyderabad, India

*Corresponding Author: S Suvana, ICAR -Central Research Institute for Dryland Agriculture, Hyderabad, India.

DOI: 10.31080/ASAG.2025.09.1468

Received: January 23, 2025

Published: January 31, 2025

© All rights are reserved by

S Suvana, et al.

Abstract

Application of lime to acid soils is a common practice to ameliorate soil acidity, the short term and long-term effect of which has not been looked upon much. To study the short-term effect of liming on acid soils an incubation study was conducted with different doses of lime for 48 days in the soils of Raipur. Long-term effects were observed by studying the total organic carbon and computing carbon input in the long term fertilizer experiment (LTFE) samples of Ranchi. In both cases, it was observed that liming decreased carbon reserves in the soil by promoting carbon mineralization and microbial respiration. In the study incubation, cumulative carbon that was mineralized was maximum in higher doses of lime and in LTFE soils, 100% NPK and lime treated plots had lower total organic carbon compared to other treatments even though carbon input was high. This shows that liming promotes carbon mineralization and additional management practices are required to balance the losses.

Keywords: Lime; Acid Soils; Carbon Mineralization; Total Organic Carbon; Carbon Reserves

Introduction

Low pH significantly impairs soil capability, with acidic soils hindering plant growth in nearly 50% of the world's cultivable lands [1,2]. While soil acidification occurs naturally, human activities have substantially accelerated this process in recent decades. Current food production systems contribute to ongoing acidification as plant biomass removal depletes soil cations. Acidic soils are known to disrupt soil carbon and nutrient cycles, negatively impact plant and soil organism growth, and jeopardize ecosystem functions like net primary production and species diversity [3-5].

Liming, which has shown considerable potential for improving soil pH and crop yields, is widely employed as a common remedy for soil acidification [6,7]. The application of lime boosts plant growth by raising soil pH, which enhances nutrient availability while reducing aluminum toxicity [1,8]. Additionally, liming facilitates the immobilization of harmful heavy metals and modi-

fies plant nutrient transformation and uptake, thereby influencing ecosystem productivity [9,10]. Nevertheless, conflicting outcomes regarding liming's effects on plant growth and yields have been documented. While some research has demonstrated significant improvements in crop yields [11,12], other studies have observed reductions in crop production, particularly when soils were limed or treated with excessive amounts [7,13]. Beyond its impact on plant productivity, pH changes induced by liming also affect soil microbial biomass, composition, and activity, as well as soil C and N availability. Furthermore, liming may influence soil organic C (SOC) concentrations and stocks.

Liming affects the preservation and breakdown of SOC. It can accelerate carbon depletion soil by increasing carbon solubility, microbial activity, and carbon decomposition rates [14,15]. However, lime-induced enhancements in plant growth, both above and below

ground, and the resulting increase in organic matter input to the soil, may lead to observed SOC accumulation [16,17]. The overall change in total SOC content after liming depends on the balance between SOC gains and losses. Typically, liming is anticipated to boost soil microbial populations and their activities, thereby increasing organic matter mineralization. Consequently, researchers have observed increases in soil respiration and decreases in SOC stocks [18,19]. In addition to pH directly impacting soil microbial processes like denitrification [20], alterations in soil microbial parameters and substrate availability also influence the generation, consumption, and emission of greenhouse gases, specifically N_2O , methane, and carbon dioxide [7,21-23]. The worldwide rise in soil acidification and liming practices is expected to have a cumulative effect on the global carbon cycle [24]. On the other hand, the liming-induced enhancement of plant growth increases C inputs through litter and root exudation, which may surpass increased C losses due to SOC mineralization, ultimately resulting in SOC accumulation [25,26]. Assessing how liming impacts the mineralization of soil organic carbon (SOC) is crucial for anticipating shifts in soil carbon balance, emissions of greenhouse gases, and the long-term viability and economic benefits of agricultural crop systems [1].

The application of lime is recognized for its immediate impact on enhancing soil biological activity, while also potentially contributing to long-term increases in soil organic matter content. The short-term effects of liming on microbial activity in both field studies and laboratory experiments [27] has been reported. The temporary boost in microbial activity following lime addition to acidic soils is well-established [28]. Studies have shown that liming can enhance various soil properties, including microbial biomass content, soil respiration rate, the microbial metabolic quotient, soil enzyme activity (specifically dehydrogenase, sulphatase, and protease), and the net mineralization of organic N and S in soil [27].

The aim of this study was to investigate whether addition of lime in acid soil affects carbon mineralization. It was hypothesized that liming promotes carbon mineralization. The study revealed that addition of lime in acid soil has an impact on carbon mineralization in long term fertilizer experiments as well as in short lab incubation studies.

Materials and Method

Soil samples were collected from different sites i.e. from Raipur and the farm of the Birsa Agricultural University, Ranchi, India ($23^{\circ}17' N$, $85^{\circ}19' E$ and 625 m above mean sea level) under the All India Coordinated Research Project on Long-term Fertilizer Experiments (AICRP-LTFE) of the Indian Council of Agricultural Research. The experiment was initiated in 1972. The major cropping system followed here is soybean- wheat. The treatment details are as follows: control, 50% NPK, 100% NPK, 100% NP, 100% N, 100% NPK+FYM, 100% NPK + Lime. The soil of the study site is Typic Haplustalf with pH of 5.3 (soil: water, 1:2). Total carbon was determined using a TOC analyser.

A short term laboratory incubation study was conducted with the samples collected from Raipur for a period of 48 days. The pH of the sample was 4.5 (soil: water, 1:2). An experiment was conducted using 100g of soil kept at 70% field capacity. Various lime rates (0, 0.25, 0.50, and 1.0 g per 100 g of soil) were applied, and the samples were incubated at room temperature. The study was replicated three times. Additionally, a separate sample with only 0.5g of lime was incubated to examine CO_2 efflux solely from lime. Carbon mineralization, measured as CO_2 , was assessed using the NaOH trap method at specific intervals (1, 4, 7, 14, 28, 34, 41, and 48 days). Excess NaOH was back-titrated with 0.5 N HCl, using saturated $BaCl_2$ and phenolphthalein as indicators. After each titration, the vials were replenished with fresh NaOH and sealed with parafilm to prevent CO_2 loss. To determine the exact strength of the acid, the HCl was standardized using 0.5 N sodium carbonate with methyl red as an indicator [29].

The amount of C evolved was calculated using the formula
 $C \text{ evolved (mg/100 g soil)} = (B-S) * N * 6 (\text{equivalent wt of C})$

Where B is the amount of HCl consumed in blank sample, S is the amount of HCl consumed in each sample and N is the normality of HCl.

Statistical analysis

One way ANOVA analysis was carried out on the data to study the effect of liming.

Results and Discussion

The results from the incubation of Raipur soil indicate that during the incubation period, carbon mineralization was higher in lime amended soils compared to non-amended soils. The cumulative carbon dioxide evolved from 1.0 g lime treated soil was found

to be significantly higher compared to other treatments (Figure 1). The carbon dioxide emanating from the control and only lime added samples were low throughout the incubation. The cumulative carbon dioxide released was in the order 1.0 g lime > 0.5g lime> 0.25g lime > blank> lime blank.




Figure 1: Cumulative carbon mineralized from soils of Raipur with different doses of lime.

The cumulative CO₂ efflux observed at the end of incubation was 0.84 mg CO₂-C/g soil for 1.0 g lime amended soils, 0.65 mg CO₂-C/g soil for 0.5g lime, 0.59 mg CO₂-C/g soil for 0.25g lime, 0.31 mg CO₂-C/g soil for control and 0.21 mg CO₂-C/g soil for control samples with 0.5g lime. The cumulative CO₂ efflux from 1.0g lime treated soils was significantly higher than 0.5g and 0.25g lime amended soils. There was no significant difference between 0.5g and 0.25g lime amended soils. The CO₂ efflux from only lime added samples show that lime does contribute to CO₂ evolution from the samples studied, though it was marginal compared to control samples.

The Total Organic Carbon (TOC) (%) and C input (kg C/ha/yr) of the soils of Ranchi under different treatments are given in table 1. The TOC under different treatment in 0-15 cm depth varied from 0.64 to 0.94 and in 15-30 cm depth varied from 0.57 to 0.73. The carbon input ranged from 602 (kg C/ha/yr) in control plots to 6094 (kg C/ha/yr) in 100% NPK + FYM treated plots. The TOC was found to be highest in 100 % NPK + FYM treated plots corresponding to the highest C input. The C input in lime treated plots was found to be higher than in other treatments excluding FYM treated plots, but no corresponding increase in TOC was observed

in this treatment. Liming typically yields beneficial outcomes by mitigating aluminum and occasionally manganese toxicity, and/or addressing calcium deficiency. The primary symptom of aluminum toxicity is stunted growth, particularly in roots and shoots. The application of lime can significantly enhance root and shoot development, leading to increased carbon return to the soil via decaying roots and plant residues. Over time, these effects may contribute to enhanced soil structure [28] due to the resulting increase in soil organic matter [30]. The total organic carbon (TOC) in lime-treated plots was measured at 0.74%, which was lower compared to other treatments. Comparable findings were observed in 34-year-old low-input and cultivated trial plots, where liming reduced soil organic carbon (SOC) in both bulk soil and aggregates [31]. It was noted that total SOC in the 0-10 cm layer either decreased or remained constant following long-term lime application [32]. The study also found that changes in each carbon type were influenced by alterations in TOC, with lower content of each carbon type in limed soils compared to unlimed soils.

This indicates that even though addition of lime promotes increase in carbon input to the soil, the carbon sequestered in the soil is less as it drives carbon mineralization.

Treatments	TOC (%)		C input (kg C/ha/yr)*
	0-15 cm	15-30 cm	
50% NPK	0.76	0.71	1530
100% NPK	0.78	0.65	1944
100% NP	0.78	0.68	1377
100% N	0.74	0.62	294
100% NPK+ FYM	0.94	0.73	6094
100% NPK+LIME	0.74	0.64	2231
Control	0.64	0.57	602

Table 1: Effect of different treatments on total organic carbon concentration (%) and carbon input under LTFE of Ranchi soil.

The decrease in soil organic carbon (SOC) caused by liming is primarily attributed to increased carbon mineralization, which results from higher carbon solubility, enhanced microbial activity, or a combination of both factors [14,15,18,33]. The growth in microbial biomass and activity in limed soils may have contributed to greater SOC mineralization. The reduction in SOC content following liming could be linked to increased microbial breakdown of SOC at optimal soil pH levels [34], which in turn speeds up SOC turnover rates. However, if the rise in biomass inputs due to liming is insufficient to counterbalance the resulting accelerated SOC turnover rate, a net loss of carbon would occur. The positive effects of liming acidic soil on above- and below-ground plant biomass production have been well-established in scientific literature [35,36].

Conclusion

This study demonstrated that short term (48 days incubation) as well as long term liming decreased SOC and promoted carbon mineralization in acid soils. Carbon mineralization increased with increase in dosage of lime (maximum at 1.0g lime/100g soil). In long term fertilizer experiments, supplementation of lime increased plant biomass and carbon input to the soil, but there was no corresponding increase in TOC in these soils. The results indicate that impact of liming on carbon reserves depends on the balance between the carbon input and the rate of carbon mineralization and microbial respiration. Thus, amelioration of soil acidity with liming should be carried out with additional management practices to maintain the carbon reserves in the soil.

Conflict of Interest

The authors have no conflict of interest to declare.

Bibliography

- Grover SP, *et al.* "Liming and priming: the long-term impact of pH amelioration on mineralisation may negate carbon sequestration gains". *Soil Security* 3 (2021): 100007.
- Kochian LV, *et al.* "Plant adaptation to acid soils: the molecular basis for aluminium resistance". *Annual Review of Plant Biology* 66 (2015): 571-598.
- Meng C., *et al.* "Global soil acidification impacts on below-ground processes". *Environmental Research Letters* 14.7 (2019).
- Stevens CJ., *et al.* "Contribution of acidification and eutrophication to declines in species richness of calcifuge grasslands along a gradient of atmospheric nitrogen deposition". *Functional Ecology* 24.2 (2010): 478-484.
- Yang Y., *et al.* "Significant soil acidification across northern China's grasslands during 1980s-2000s". *Global Change Biology* 18 (2012): 2292-2300.
- Eze S., *et al.* "Soil organic carbon stock in grasslands: Effects of inorganic fertilizers, liming and grazing in different climate settings". *Journal of Environmental Management* 223 (2018): 74-84.
- Wang Y., *et al.* "Potential benefits of liming to acid soils on climate change mitigation and food security". *Global Change Biology* 27.12 (2021): 2807-2821.
- Weil RR and Brady NC. "The nature and properties of soils. Pearson, Boston".
- Whitmore AP, *et al.* "Technologies for increasing carbon storage in soil to mitigate climate change". *Soil Use and Management* 31.S1 (2015): 62-71.
- Fornara DA., *et al.* "Increases in soil organic carbon sequestration can reduce the global warming potential of long-term liming to permanent grassland". *Global Change Biology* 17 (2011): 1925-1934.
- Li Y., *et al.* "Liming effects on soil pH and crop yield depend on lime material type, application method and rate, and crop species: A global meta-analysis". *Journal of Soils and Sediments* 19.3 (2019): 1393-1406.
- Crusciol CAC., *et al.* "Lime and gypsum combination improves crop and forage yields and estimated meat production and revenue in a variable charge tropical soil". *Nutrient Cycling in Agroecosystems* 115 (2019): 347-372.
- Holland JE., *et al.* "Yield responses of arable crops to liming -An evaluation of relationships between yields and soil pH from a long-term liming experiment". *European Journal of Agronomy* 105 (2019): 176-188.
- Hénault C., *et al.* "Management of soil pH promotes nitrous oxide reduction and thus mitigates soil emissions of this greenhouse gas". *Scientific Reports* 9 (2019): 1-11.
- Bezdicsek DF, *et al.* "Subsoil ridge tillage and lime effects on soil microbial activity, soil pH, erosion, and wheat and pea yield in the Pacific Northwest, USA". *Soil and Tillage Research* 74.1 (2003): 55-63.
- Fuentes JP, *et al.* "Microbial activity affected by lime in a long-term no-till soil". *Soil and Tillage Research* 88.1 (2006): 123-131.

17. Bronick CJ and Lal R. "Soil structure and management: A Review". *Geoderma* 124.1 (2005): 3-22.
18. Briedis C., *et al.* "Soil organic matter pools and carbon-protection mechanisms in aggregate classes influenced by surface liming in a no-till system". *Geoderma* 170 (2012): 80-88.
19. Ahmad W., *et al.* "Inorganic and organic carbon dynamics in a limed acid soil are mediated by plants". *Soil Biology and Biochemistry* 57 (2013): 549-555.
20. Biasi C., *et al.* "Direct experimental evidence for the contribution of lime to CO₂ release from managed peat soil". *Soil Biology and Biochemistry* 40 (2008): 2660-2669.
21. Bakken LR and Frostegård Å. "Emerging options for mitigating N₂O emissions from food production by manipulating the soil microbiota". *Current Opinion in Environmental Sustainability* 47 (2020): 89-94.
22. Khaliq MA., *et al.* "Soil liming effects on CH₄, N₂O emission and Cd, Pb accumulation in upland and paddy rice". *Environmental Pollution* 248 (2019): 408-420.
23. Royer-Tardif S., *et al.* "Can alkaline residuals from the pulp and paper industry neutralize acidity in forest soils without increasing greenhouse gas emissions?" *Science of the Total Environment* 663 (2019): 537-547.
24. Shaaban M., *et al.* "Effects of dicyandiamide and dolomite application on N₂O emission from an acidic soil". *Environmental Science and Pollution Research* 23 (2016): 6334-6342.
25. Fisher J., *et al.* "Quantifying the acid balance for broad-acre agricultural systems". *Handbook of Soil Acidity* (2003): 117-133.
26. Abalos D., *et al.* "Trade-offs in greenhouse gas emissions across a liming-induced gradient of soil pH: Role of microbial structure and functioning". *Soil Biology and Biochemistry* 150 (2020): 108006.
27. Liang C., *et al.* "The importance of anabolism in microbial control over soil carbon storage". *Nature Microbiology* (2017).
28. Haynes RJ and Swift RS. "Effects of lime and phosphate additions on changes in enzyme activities, microbial biomass and levels of extractable nitrogen, sulphur and phosphorus in an acid soil". *Biology and Fertility of Soils* 6.2 (1988): 153-158.
29. Haynes RJ. "Lime and phosphate in the soil-plant system". *Advances in Agronomy* 37 (1984): 249-315.
30. Mishra PK., *et al.* "Sustainable Land Management: Issues". *Problems and Prospects* (2019).
31. Haynes RJ and Naidu R. "Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: A Review". *Nutrient Cycling in Agroecosystems* 51.2 (1998): 123-137.
32. Aye NS., *et al.* "The impact of long-term liming on soil organic carbon and aggregate stability in low-input acid soils". *Biology and Fertility of Soils* 52.5 (2016): 697-709.
33. Wang X., *et al.* "Long-term effect of lime application on the chemical composition of soil organic carbon in acid soils varying in texture and liming history". *Biology and Fertility of Soils* 52.3 (2016): 295-306.
34. Kemmitt SJ., *et al.* "pH regulation of carbon and nitrogen dynamics in two agricultural soils". *Soil Biology and Biochemistry* 38.5 (2006): 898-911.
35. Ahmad W., *et al.* "Temperature sensitivity and carbon release in an acidic soil amended with lime and mulch". *Geoderma* 214 (2014): 168-176.
36. Hati KM., *et al.* "Impact of long-term application of fertilizer, manure and lime under intensive cropping on physical properties and organic carbon content of an Alfisol". *Geoderma* 148.2 (2008): 173-179.
37. Kostic L., *et al.* "Liming of anthropogenically acidified soil promotes phosphorus acquisition in the rhizosphere of wheat". *Biology and Fertility of Soils* 51.3 (2015): 289-298.