



Demonstration of Integrated Applications of site specific NPS Fertilizer Recommendation with Vermicompost for Bread Wheat Production in the Burka Jiren Watershed Under Lime Condition

Bati Dube*, Amanuel Kejela, Dechasa Mengistu, Dagne Chimdessa, Gedefa Sori, Mosisa Mezgebu and Belay Beyene

Oromia Agricultural Research Institute (IQQO), Bedele Agricultural Research Center (BeARC), Bedele, Ethiopia

***Corresponding Author:** Bati Dube; Oromia Agricultural Research Institute (IQQO), Bedele Agricultural Research Center (BeARC), Bedele, Ethiopia.

Received: July 21, 2025

Published: October 28, 2025

© All rights are reserved by

Bati Dube., et al.

Abstract

Degradation of physical properties caused as a result of the loss of soil organic matter and reduced presence and activity of soil organisms also occur in heavily polluted soils. The addition of vermicompost is used to increase organic matters in the soil. Therefore, this study was conducted to demonstrate integrated applications of site specific NPS fertilizer recommendation with vermicompost for Bread wheat production in the Burka Jiren watershed under lime condition. The experiment was implemented during 2024 crop season in Gechi district at Burka Jiren community watershed. The treatments consisted of two levels; namely (T1) Blanket Recommendation (1:1 ratio of urea and NP respectively) and (T2) 1/2 Recommended vermicompost + 1/2 recommended N + recommended NPS. The collected data was analyzed using T-test and interpreted as mean and standard deviation by SAS software version 9.0. Results indicated that the grain yield of bread wheat was significantly ($P < 0.05$) influenced between two treatments. The highest grain yield (47.224qt/ha) was recorded from treatment two which was treated by 1/2 Recommended vermicompost + 1/2 recommended N + recommended NPS. However, the lowest grain yield (23.625qt/ha) was obtained from blanket recommendation of NPS fertilizer alone. The integrated use of Vermicompost with NPS fertilizer significantly increased the mean of soil pH from 5.0 to 5.26. Both treatments presented similar levels of organic carbon and total nitrogen. Farmers in the Burka Jiren community watershed should consider integrating agricultural lime with site specific NPS fertilizer to mitigate soil acidity and enhance crop productivity.

Keywords: Vermicompost; Bread Wheat; NPS Fertilizer; Urea

Introduction

Degradation of physical properties caused as a result of the loss of soil organic matter and reduced presence and activity of soil organisms also occur in heavily polluted soils [1]. It has been estimated that soil pH has reduced globally an average of 0.26 due to acid deposition and N additions. Soil acidification also enhances the mobilization and bioavailability of certain trace elements, increas-

ing the risk to human health and the environment and decreasing crop growth [2].

Smallholder farmers in Africa typically only have access to blanket fertilizer recommendations which are defined over very broad areas and may not be optimal for local production conditions. Blanket fertilizers recommendations are used to encourage farm-

ers adopt particular blends of fertilizer in Ethiopia. These recommendations, however, may not be optimal (or even suitable) for particular soil types or agro-climatic conditions within broad areas receiving such recommendations [3].

Soil degradation and nutrient depletion have been serious threats to agricultural productivity and food security in Ethiopia. Over the years, soil fertility has also declined due to the increase in population size and decline in plot size. Nitrogen (N) and phosphorus (P) are the nutrients that are most lacking [4,3]. In 2007, the Ministry of Agriculture and Natural Resources (MoANR) and Agricultural Research Centers together developed regional or blanket fertilizer recommendations (fertilizer types and application rates for different crops). These recommendations inform farmers how much fertilizer in kilograms they should apply per hectare without performing any soil test and regardless of agro-ecological zones. Since nutrient management requirements of farms vary across crop type, soil type and other agro-ecological characteristics, these blanket fertilizer recommendations may not be suitable for all farmers. The adoption of fertilizer remains low and the average application rates by those who do use fertilizer are generally lower than recommended rates [3]. Bread wheat (*Triticum aestivum* L.) is one of the most important cereal crops playing a vital role in the country's economy, food security, and consumption. In Ethiopia, wheat covers an estimated area of 1.7 million ha with total annual grain production of 4.6 million tons by 4 million smallholder farmers. In terms of area of production, wheat ranks fourth after tef (*Eragrostis tef* [Zucc.] Trotter), maize (*Zea mays* L.) and Sorghum (*Sorghum bicolor* L.) and third in total grain production after tef and maize in Ethiopia [5]. However, the productivity of wheat is still low with a national average yield of 3 t ha⁻¹ stations, i.e., 5 t ha⁻¹ compared to yields obtained from the research [6].

Integrated soil fertility management is a set of soil fertility management practices that necessarily include the use of fertilizer, organic inputs, and improved germplasm combined with the knowledge on how to adapt these practices to local conditions, aiming at maximizing agronomic use efficiency of the applied nutrients and improving crop productivity [7].

Lime is a soil conditioner and controls the soil acidity by neutralizing the effects of acids from nitrogen (N) fertilizer, slurry and high rainfall. Other benefits include an increase in earthworm activity, improvement in soil structure and grass is more palatable to livestock [3].

The primary reason to apply agricultural lime is to correct the high levels of acidity in the soil as acid soils reduces plant growth by inhibiting the intake of major plant nutrients -nitrogen, phosphorus and potassium. Some plants, for example legumes, will not grow in highly acidic soils.

Soils become acidic in a number of ways: Leaching of land caused by high rainfall levels, Minerals loss over time caused by crop removal Application of modern chemical fertilizers, which are the major contributors of acidified soil. Soil with pH below 5.5 and below 70% of saturation requires liming. Agricultural lime has good effects on soil as it increases the pH of acidic soil, provides a source of calcium and magnesium for plants, Permits improved water penetration for acidic soils and Improves the uptake of major plant nutrients (nitrogen, phosphorus, and potassium) of plants growing on acid soils.

Application of the mineral P fertilizer with other amendments can be used to improve P deficiency in acid soils. Even though the chemical fertilizers including mineral P are used to increase productivity for a certain time, their negative impacts coupled with their high cost have prompted the interest in the use of organic fertilizers as source of nutrients. Organic fertilizer application has been reported to improve crop growth by supplying plant nutrients as well as improving soil physical, chemical, and biological properties [8,9].

Vermicompost is one of the stabilized, finely divided organic fertilizers with a low C: N ratio, high porosity, and high water-holding capacity, in which most nutrients are present in forms that are

readily available for plants [10,11]. Application of Vermicompost improves overall physical and biochemical properties, and at the same time, decreases exchangeable acidity which can support a release of plant nutrients in the acidic soils, promotes growth of plants and positive effect on growth and productivity of cereals and legumes. When it is compared with conventional compost, different research on Vermicompost indicated that Vermicompost very important in promoting plant growth from 50 to 100% over conventional compost and from 30 to 40% over chemical fertilizers. Therefore this demonstration initiated with these objectives: To demonstrate integrated use of site specific NPS fertilizer recommendation with vermicomposting and under lime condition as acid soil management practices and to improve farmers' awareness about integrated use of site specific NPS fertilizer recommendation with vermicomposting for Bread wheat production and under lime condition as acid soil management

Materials and Methods

Descriptions of the Study Area

The watershed is situated in the Gechi district of Buno Bedele zone, southwestern Ethiopia (Figure 1). The specific place of the watershed is Bido Jiren Kebeles, and so the local communities named the watershed Burka Jiren Community Watershed. The watershed lies between 8°12'50.4" or 8°14'52.8 N latitudes and between 36°25'27.48" or 36°26'52.8" E longitudes. Burka Jiren Community Watershed is about 15 km from Gechi town in Gechi district and about 35 km from Bedele town in Buno Bedele zone. The watershed covers about 535.13 hectares, and 148 smallholder households benefit from the watershed. According to the Google Earth Pro software, the length of the Burka Jiren Community Watershed is about 6 kilometers from the watershed in-late to the outlet. Based on slope gradient, Burka Jiren Community Watershed is characterized mostly by slop, gently slop, and strongly slop. The dominant soil type of the watershed is nitisils, and the common crops produced by the communities are tef and wheat.




Figure 1: Map of Burka Jiren Community Watershed.

Sites, farmer's selection and Technology promotion events

The activity was conducted in purposively on four selected farmers from the watershed based on wheat production potential, for two years initial available phosphorus level of soil and their accessibility for determine the amount of P_2O_5 from NPS was determined. One FREG having 15 members (male and female) was established in each upper, middle and lower parts of the watershed. After, established of FREG four host farmers were selected among the FREG members and the activity were implemented on their farm land. The farmers Selection was done with collaboration of

district experts and development agents. After the establishment of FREGs Mini-field day and training was provided for development agents, subject matter specialists (SMS) and FREG members on the use of site specific NPS Fertilizer with vermicompost on wheat production, management and its dissemination methods.

Experimental design

The experiment consisted of two treatments namely (T1) Blanket Recommendation (1:1 ratio of urea and NP respectively) and (T2) 1/2 Recommended vermicompost + 1/2 recommended N +

recommended NPS). The plot size was 10 m x 10 m (100 m²) for each plot and the liben wheat variety was planted in rows. Recommended lime was broadcast manually and mixed thoroughly in upper soils at 20cm depth (plow depth) to each plots. Amount of recommended N fertilizer kg/ha in vermicompost equivalent ratio of N in urea was calculated and was applied the recommended amount at sowing time. The experiment was conducted on four sites in sub-watershed.

Data to be collected

Soil data (pH, organic carbon (OC), total N, available P, CEC and exchangeable acidity) before and after applications. Agronomic data of grain yield and farmer's perceptions on integrated use of site specific NPS fertilizers and vermicompost was collected.

Data Analysis

The collected data was analyzed using T-test and interpreted as mean and standard deviation. Partial budget analysis was done to identify economically feasible recommended site specific NPS fertilizer with vermicompost and NPS fertilizer alone.

Results and Discussions

Soil Chemical Properties before Planting of Experimental Field

The soil PH (H₂O) of the study sites were strongly to slightly acidic the value ranged from 4.7 to 5.52 with mean 5.0 (Table 1) according to the rating indicated by FAO, 2006. The available phosphorus contents of the soil were very low which ranged from 0.74 to 2 ppm with mean 1.3ppm (Table 1) this results agreed with FAO (2006) indicated that the available Phosphorus which put under less than 5ppm was very low. The results of organic carbon indicated medium with value ranged from 3.25 to 4.75% with mean 4.1% of organic carbon (Table 1) this obtained results linked with FAO (2006) [12]. Looking this result the study area of soil needed additional organic material to full fill the educated level of the organic carbon. The gain results of Total Nitrogen of the study sites before planting was medium with value ranged 0.28 to 0.41% with mean 0.4% of TN (Table 1) it is also, agreed with FAO (2006) and Cation Exchange capacity of the soil ranged from 14.5 to 23 cmol(+)/kg soil, with the mean 16.6 cmol(+)/kg soil (Table 1) results shows medium level.

Sites	Site 1	Site 2	site 4	site 4	Mean
pH(H ₂ O)	4.9	4.7	4.8	5.52	5.0
Exchangeable Acidity(cmol+)/kg soil)	2.96	2.69	2.32	0.12	2.0
Available Phosphorus(ppm)	1.64	2	0.74	0.98	1.3
Organic Carbon (%)	4.03	4.75	4.23	3.25	4.1
Total Nitrogen (%)	0.35	0.41	0.36	0.28	0.4
Cation Exchange Capacity(cmol+)/kg soil)	14	23	14.7	14.5	16.6

Table 1: Soil Chemical Properties before Planting of Experimental Field.

The soil chemical properties of experimental sites after harvesting wheat, comparing two treatments:

The application of vermicompost significantly increased the mean of soil pH from 5.0 to 5.26, (Table 1,2) demonstrating its effectiveness in reducing soil acidity. Organic Carbon and Nitrogen: Both treatments presented similar levels of organic carbon and

total nitrogen, telling that Vermicompost application did not negatively influence these soil health indicators. The combined use of site specific NPS fertilizer and vermicompost improved soil pH (reducing acidity) without compromising organic carbon or nitrogen content, supporting better wheat production in acidic soils.

Treatments	Mean of soil PH	Mean of organic carbon	Mean of total nitrogen
Blanket recommended of NPS and urea	4.98	3.36	0.26
Site specific NPS + ½RN from urea + ½ RN from vermicompost	5.26	3.88	0.33

Table 2: Soil Chemical properties of experimental sites after harvesting.

Participatory mini-Field Visit Based Training for Participants

Participatory mini-field visit based training was given to participants on the concept and principles of FREG, the role and accountability of the FREG members in management of the trial, application of Site specific NPS Fertilizer with vermicompost and alone NPS fertilizer on wheat production and influence of vermicompost on selected soil chemical properties. Participants visit the trial field

as all FREG members practice on their own field and whereas others share the experience. A total of 37 participants among which 33 farmers (29 male and 4 female), 2 DAs (1 male and 1 female) and 2 other concerned stakeholders (2 male and 0 female) were participated on field visit based training.

Watershed	Participant	Male	Female	Total
Burka Jiren Community	Farmers	29	4	33
	DAs	1	1	2
	Other stakeholders	2	0	2
Total		32	5	37

Table 3: Stakeholders participated on field visit based training events.

Effects of site specific NPS-fertilizer with vermicompost and NPS-fertilizer alone on Wheat grain yield

The results of these study indicated that significance ($P < 0.05$) difference between mean grain yield of bread wheat. The lowest mean grain yield (23.625 qt/ha) in table 3 was observed from treatment one (T1) which was treated by farmer practice (blanket recommended of NPS and urea). Whereas, the highest mean grain yield (47.224 qt/ha) in table 3 was obtained from treatment T2 (that treated by NPS + half of recommended N from urea + half of N from vermicompost recommended). The integrated approach (T2) nearly doubled the grain yield compared to T1, demonstrating its effectiveness for bread wheat production. The higher standard deviation in T2 suggests greater variability in yield outcomes, possibly due to factors like soil heterogeneity or management practices. Combining site-specific NPS fertilizer with vermicompost under lime conditions can substantially enhance wheat productivity. Similarly, the pre-sowing application of the vermicompost extract had the evidential impact on the grain yield [13]. This results also,

agreed with [14], who stated that combined application of vermicompost and NPS fertilizers significantly increases grain yield. This suggests that the application of NPS fertilizer along with vermicompost (T2) had a real and measurable positive effect on wheat grain yield compared to the blanket recommendation (T1).

Farmer's preference of wheat production

Farmers selected T2 which treated by Site specific NPS + ½RN from urea + ½ RN from vermicompost rather than blanket recommendation depending on the result obtained.

Economic partial budget analysis

The partial budget analysis of the treatments was done considering all variable costs and all benefits of grain yield. The analysis of marginal rate of return (MRR), revealed that the rate of return per unit cost of production was highest for application of site specific NPS + half of recommended N from urea + half of N from vermicompost recommended (% MRR =626). This showed that it would yield

Treatments	Rank	Reason of farmers preference
Blanket recommended of NPS and urea	2 nd	Medium performance of head, medium length, and medium headed yield
Site specific NPS + ½RN from urea + ½ RN from vermicompost	1 st	Very good performance than T1, full head, attractive green color, longer than T1 and long headed yield, changed soil color from red to black and believed to improve soil fertility and structure

Table 5: Reason for preference of result obtained on wheat grain yield.

gave 6.26 Ethiopian Birr for every Birr invested. Thus, applications of site specific NPS + half of recommended N from urea + half of N from vermicompost recommended ha⁻¹ of is economically benefi-

cial as compared to the blanket recommendation of NPS with Urea treatments, because the highest net benefit (NB) and marginal rate of return (MRR) was above the minimum level (100%).

Variables	Treatments	
	Bread Wheat with Vermicompost application	Bread Wheat without Vermicompost application
Yield obtained (qt ha ⁻¹)	47.224	23.625
Sale price (ETB/qt)	8,000	8,000
Total Revenue (Price * Qt)	377,792	189,000
Variable Costs/ha in ETB.		
Land preparation	4,000	4,000
Seed purchase	6,500	6,500
Fertilizers purchase	7,838	7,838
Cost of lime	500	0
Cost of labor for weeding	3,200	3,200
Cost of labor for Vermicompost application	1,000	0
Cost of Labor for harvesting	3,300	3,300
Cost of Labor for threshing	4,000	4,000
Total Variable Costs	30,338	28,838
Cost of Land ETB/ha	30,000	30,000
Total cost (TVC + TF C)	60,338	58,838
Gross Margin (GM) = TR-TVC	347,454	160,162
Net Profit = TR-TC or GM-TFC	317454	130,162
Benefit-cost ratio = Total Revenue/Total Cost	6.26	3.21

Table 6: Partial budget analysis for the site specific NPS and Agricultural Lime on bread wheat grain yield.

Conclusions and Recommendations

The integrated use of site-specific NPS and vermicompost (T2) significantly increased wheat grain yield compared to the alone Site-specific NPS fertilizer (T1).

Farmers could potentially double their wheat yield by adopting this treatment, though further field trials may be needed to confirm consistency across different conditions.

Conflict of Interest

The authors declare that they have no conflict of interest.

Acknowledgements

The authors would like to thanks Climate Action through Land scape Management (CALM P4R) for their funding of this research with collaboration Oromia Agricultural Research Institute and Be-

dele Agricultural Research Center for providing all the necessary facilities required respectively. Additional laboratory analysis team of Bedele Agricultural Research Center for their support and unreserved effort to provide reliable samples analysis data on time.

Bibliography

1. Korkina IN and Vorobeichik EL. "Humus Index as an indicator of the topsoil response to the impacts of industrial pollution". *Applied Soil Ecology* 123 (2018): 5-463.
2. Tian D and Niu S. "A global analysis of soil acidification caused by nitrogen addition". *Environmental Research Letters* 10.2 (2015): 024019.
3. Ayalew Hailemariam., *et al.* "Site-specific agronomic information and technology adoption: A field experiment from Ethiopia". *Journal of Development Economics* (2022).
4. Fassil Kebede and Charles Yamoah. "Soil Fertility Status and Numass Fertilizer Recommendation of Typic Hapluusterts in the Northern Highlands of Ethiopia". *World Applied Sciences Journal* 6.11 (2009): 1473-1480.
5. CSA. "Agricultural Sample Survey Report on Area of Production of Major Crops Volume I, in: The Federal Democratic Republic of Ethiopia Central Statistical Agency. CSA Bulletin No 584, Addis Ababa.
6. H Molla., *et al.* "Response of bread wheat to integrated application of vermicompost and NPK fertilizers". *African Journal of Agricultural Research* 13.1 (2018): 14-20.
7. Mugwe JN., *et al.* "Integrated soil fertility management in sub-Saharan Africa: Evolving paradigms toward integration. Springer Nature Switzerland, AG (2019): 1-13.
8. Mengistu DK and Mekonnen LS. "Integrated agronomic crop managements to improve tef productivity under terminal drought". *Water Stress. Vienna Tech* 1 (2011): 235-254.
9. Melese A., *et al.* "Effects of lime, wood ash, manure and mineral P fertilizer rate on acidity related chemical properties and growth and P uptake of wheat (*Triticum aestivum* L.) on acid soil of Farta district, northwestern highlands of Ethiopia". *International Journal of Agriculture and Crop Sciences* 8.2 (2015): 256-269.
10. NQ Arancon., *et al.* "Effects of vermicompost produced from food waste on the growth and yields of greenhouse peppers". *Bio-Resources Technology* 93.2 (2004): 139-144.
11. J Dominguez. "State of the art and new perspectives on vermicomposting research". in *Earthworm Ecology*, C. A. Edwardspp. 401-424, CRC Press, Boca Raton, FL, USA, 2nd edition (2004).
12. Food and Agricultural Organization (2006).
13. Kováčik Peter., *et al.* "Impact of vermicompost extract application into soil and on plant leaves on maize phytomass formation". *Journal of Ecological Engineering* 16.4 (2015): 143-153.
14. Kinfe Tekulu., *et al.* "Effect of vermicompost and blended fertilizers rates on yield and yield components of Tef (*Eragrostis tef* (Zucc.) Trotter)". *Journal of Soil Science and Environmental Management* 10.6 (2019): 130-141.