



A Review of Nuclear Emergency System Adapted for Food Chain Model

Priscilla Oforiwa^{1*}, Nanaadom Abayie Nyarko², Manchun Liang¹,
Guofeng SU¹ and Li KE¹

¹Department of Engineering Physics, Tsinghua University, PR China

²Department of Industrial Engineering and Engineering Management, Western
New England University, Springfield MA, USA

*Corresponding Author: Priscilla Oforiwa, Department of Engineering Physics,
Tsinghua University, PR China.

Received: July 15, 2021

Published: May 31, 2022

© All rights are reserved by Priscilla
Oforiwa, et al.

Abstract

Food chain modeling development is a vital part of emergency planning and response in the field of nuclear industries, chemical industries, petroleum and fire. The adverse effects of some elements and their properties are of great importance in developing a decisive policy approach. Terrestrial food chain modelling gives the predictive approach and forecasted for effects of food substances to the general population. Nuclear Emergency Response and planning is a major block in the safety development of Nuclear Power plant. To react rapidly and adequately to a radiological crisis, significant level coordination is required between various stake-holders in higher organizational level.

In this paper, Nuclear Emergency systems are reviewed and analyzed; from their application in atmospheric dispersion to its application in terrestrial models (food chain approach) and countermeasures. This study is a review of the existing Nuclear Emergency Planning and Response platform such RODOS, ARAC and WSPEDI; the dynamics and the similarities of modelling approaches and its effectiveness in Food chain modelling and policies.

The conclusion from the study illustrate the need to develop a Robust Emergency system that for food chain predictions in radiological situation.

Keywords: Emerging Prediction; Radiological; Food Chain; Modelling

Introduction

Nuclear emergency response and Preparedness plays an important role in the safe operations of nuclear powerplant. However, previous incidents such as the Chernobyl accident, Fukushima accident and weapon fall out sparked a new interest in the accidental release of radionuclides to the atmosphere [1]. These atmospheric releases are further dispersed on soil or vegetative cover either by wet deposition or dry deposition [2].

Various studies have been performed over the years from the Chernobyl accident deposition, however, due to the complexities of the environmental factors such as soil properties, lithological properties, organic matter content and the physiochemical properties of the

soil and plant, there is the need for adverse research pertaining to this area in different geographical location to circumspect the disadvantage in food in order to strengthen emergency planning should the unexpected happens [3,4].

In view of the current growth of this industry in the world, there is an absolute need for nuclear emergency decision system to ensure maximum safety and to give adverse security [5]. The various emergency systems are analyzed and compared with their fundamental use on emergency situation; How these systems can be adapted for Food chain predictions in a radiological event. The system analyzed are Rodos or JRodos system, Speedi/Wspeedi system and the Arac/Narac system

Rodos system is an online emergency decision supporting system that incorporates atmospheric dispersion models and the food dose model for terrestrial environment [6]. However, in recent years, there has been the need for model development in the system as the larger part of the systems values were built on with the Chernobyl influences and the geographic conditions in Europe. Various challenges were faced in trying to adapt the model with the various soil and food dynamics of China. In essence, a comparative study was made with the results and conclusions made [5,7].

Nuclear accident in history

In reference to the increasing status of nuclear energy in the world, the threat of radioactive material leakage has become a concern. Although nuclear power technology is constantly improving, the safety level of nuclear power is constantly improving, and the probability of nuclear accidents is also reduced. But Murphy's law tells us that if anything is possible to go wrong, it will go wrong. Once a nuclear accident occurs, it often has a significant impact on the environment, population, economy and society, resulting in incalculable consequences [5].

There are three far-reaching nuclear accidents in the account of nuclear power development. The first one was the core meltdown accident at the Three Mile Island nuclear power plant in the United States on March 28, 1979, which caused the release of some radioactive materials and the release of about 1017 BQ of rare gases into the atmosphere. During the accident, about 140000 people were evacuated within 20 miles. Although no death was caused, it caused panic and protest among the public, causing long-term social impact. It is the most serious nuclear accident in the United States.

The second was the nuclear accident at the Chernobyl nuclear power plant on April 26, 1986, which resulted in 31 deaths in three months, 60000-80000 deaths in 15 years, 134000 people affected by radiation diseases and 115000 people forced to evacuate within 30 kilometers. The aggregate sum of radioactive materials delivered during the accident is about 12×10^{18} BQ, making it the most serious nuclear accident ever. The third is 2011 On March 11, 2011, the Fukushima nuclear power plant leakage accident caused by the magnitude 9 earthquake and huge tsunami in Japan. During the accident, the evacuation ranges gradually increased, and nearly 80000 people were evacuated within a radius of 20 kilometers. It is estimated that Iridium 131 released to the atmosphere reached 1.6×10^{17} Bq, $^{137}\text{Cesium}$ reached 1.5×10^{16} Bq.

In addition, there are also a large number of radioactive effluents leaking into the ocean, which is the largest marine nuclear pollution event at present. The leaked radioactive materials have caused global impact on both aquatic and terrestrial Foodchain [8].

Real-time online decision support system (RODOS)

Following the Chernobyl mishap, expanding assets were apportioned in numerous nations to the improvement of plan of off-site crisis reaction in case of a Nuclear accident. The Real-time on-line decision support system, Rodos was initiated in the 1990 [9]. Rodos is a non-commercial comprehensive decision support system for nuclear and radiological emergencies. The principle goals of Rodos is to give the methodological premise, foster models and information bases and introduce the equipment and programming structure of a framework which offers an extensive choice help from the beginning phases of a mishap up to numerous years after the delivery and from the assortment of the site to far off regions unperturbed by public limits. The system is capable of providing consistent and comprehensive information at local, regional and national levels [10].

The system is also a useful tool in the no accident phase thus it can be used for preparing a possible future event, planning and training and scenario development.

Briefly, the system contains models for the atmospheric and aquatic pathways.

The post-Chernobyl radioecological studies have confirmed the long residence time of certain radionuclides, these are Cesium-134 and cesium-137 in a semi-natural ecosystem [7]. The atmospheric dispersion and deposition uses meteorological information, close-in distances and long-range. The source term serves as an input. In the RODOS system, fixed data nuclides which has been currently implemented includes all radionuclides in light water reactor (LWR) accidents, extra radionuclides for supplement of some conceivably significant radioactive rot chains. Some actuation items or other radionuclides that are generally significant for LWR reactors in typical activity mode, Fast breeder reactors or fusion reactors. However, some radionuclide compounds such as tritiated hydrogen (HT) and tritiated water(HTO) for fusion reactors [11]. The incident in England in 2006 added Po -210 as a potentially relevant nuclide for calculations involving radiological emergencies.

In the RODOS system, simulation for an early dose is done with the Emersim system and the terrestrial food chain uses FDMT [12].




Figure 1: System Design of RODOS System [12].




Figure 2: General system overview of RODOS [7].




Figure 3: The process flow chart for RODOS.

The Rodos system gives a wide variety of models, however the input of the system is from the atmospheric dispersion which is estimation of individual and collective doses. The deposition input data are calculated using the DEPOM model.




Figure 4: Food chain model of the RODOS system.

In the RODOS system, terrestrial food chain and dose model assess doses to population through all relevant pathways such as inhalation and ingestion, external exposure from plume and ground. The FDMT is generally created on the previous dynamic model ECOSYS-87(improving models) [6,13] that was initially carried out inside the Microsoft Dominate. In any case, a significant part of the formative work including the mathematical particular of a significant number of the boundaries utilized in ECOSYS-87 was finished in the 1980's and thus much thought isn't given enormous quantities of radioecology intends brought about by the Chernobyl accident in 1986 [6]. Again, the first boundary condition is mostly explicit to southern Germany horticultural conditions albeit the

model was taken into account adaptability and variation to different conditions [14].

Comparing the deliverables of the RODOS system to the radioecological parameters of the FDMT (Food chain Dose Modelling Transfer) to the international recommendation according to the IAEA, an implementation of the I131, Cs134,137 and Sr90 the influence of each region must be investigated [15]. The starting point of the FDMT computations are the yields from the atmospheric dispersion models. In Root take-up is determined by means of the utilization of soil-to-plant focus proportion, Fv, (as characterized in IAEA (2010)) however referred to as TFi inside FDMT (Bq/kg new mass (FM) plant to Bq/kg dry mass (DM) soil); note in IAEA (2010) Fv is characterized on a dry mass plant premise) [16]. By definition, Fv, suggests balance or semi harmony conditions in the dirt plant framework numerous months post-mishap (IAEA 2009) [17]. In the event that deposition happens

during the growth season, a decreased root take-up is accepted by means of the use of a decrease factor (characterized by the proportion of the time frame from statement to collect the entire developing period, or 50d if the developing period is longer than this) [11].

The exchange of radionuclides from grub into animal product is depicted by the feed transfer coefficient, Fm (for milk), Ff (for meat), alluded to as TFm. in FDMT [18]. The conditions used to compute radionuclide activity concentration in the animal (or product) with time represent the unique idea of the framework by considering the admission of action from feedstuffs by animals throughout different time stretches and taking into account the misfortune/depuration of the radionuclide by means of natural discharge [7] been asserted recently with reference placed on the theme several national and international organization such as the environmental protection agency, world health organization.




Figure 5: System overview of Rodos software in atmospheric dispersion.

Scientific limitations of the JRODOs system

The JRodos is the nuclear decision support system with a wide range of uses for nuclear emergency. It uses ranges from atmospheric dispersion models to food chain modelling, however in the event proceeding from the Fukushima nuclear accident proved the existence of limitations for the JRODOS system [11].

In case of Radiological accident and the result of long haul recovery measure managing uncertain data is a characteristic issue for dynamic model. Uncertain limitation related to for instance, incomplete source term and the prevailing whether results in dose assessment that differ dramatically from reality [19].

The food chain process in the JRODOS system proves complications with its large amount of parameters which may

contain significant error. These parameters and dose might be difficult to verify. In JRODOS only deterministic outcomes disregarding uncertainty groups are accessible for clients. In the event that the client needs to research vulnerabilities of the source term, numerous models run with various info which should be performed. Vulnerability forecasts are not an indispensable yield related with the expectations endpoint of JRODOS [20].

The post Fukushima and Chernobyl accidents proved a high limitation with radio-ecological models which failed to predict areas of the Europe region where high radio cesium transfer through the food chain has persisted for decades for following Chernobyl [2].

Application models limitation based on Fukushima Case

In JRODOS, radionuclide transfer parameters used in model predictions were frequently adaptable (IAEA 2010). The parameterization of food chain [21] of JRodod system is centered on ECOSYS-87 model [6]. This model originates before the worldwide assemblage of radionuclide move boundaries upsides of the IAEA; which consolidates the enormous of information acquired in post-Chernobyl studies [22].

The JRodod uses natural half-life values for animal inferred food products where there is the requirement for move boundaries for explicit food items which are additionally proper from nearby conditions [23].

Radioactive iodine has proven to be a major health risk after a nuclear accident, however its environmental behavior has been poorly studied for the modelling of the food chain in JRodod. The comparative short-lived of this radionuclides $t_{1/2}$ 8yrs. However, in the data used for JRodod (IAEA 2010) contains no data for transfer of I to crops [19].

Underlying limitations are

- Lack of transfer boundaries for explicit foodstuffs.
- Lack of transfer parameters appropriate for local conditions
- Variability in transfer parameters.

Software modelling limitations (Process based models)

JRodod predicts the transmission of radionuclides to human food-stuffs which uses stability absorption to define the transfer

from soil to plants [9]. These are parametric generic data with soil categorization.

Recent use of the JRodod model under natural conditions found in Fukushima influenced regions in Japan featured a few issues. Contrasting grass focus proportions with model with values estimated in a few Fukushima influenced regions noticed that the determined fixation proportion esteems digressed from noticed information. The model extensively thought little of the noticed radiocesium take-up into grass when utilizing upsides of soil PIP and soil arrangement Potassium (K) focus assessment from qualities [24,25].

The model significantly disparaged the noticed radiocesium take-up into grass when utilizing upsides of soil Tear and soil arrangement potassium (K) focus.

- The mathematical mean (GM) of the model focus proportion esteems was in excess of a significant degree lower than the GM of the noticed qualities. The GM of the model focus proportion esteems was just about multiple times more noteworthy than that of the noticed qualities when utilizing the deliberate Tear and the K fixation [25].
- The calculated RIP and K factors of 10 and 3 separately overestimation of radio cesium sorption could be credited to various elements, the condition in the model used to ascertain the Tear had been gotten from European soil whose dirt qualities are extraordinarily unique in relation to those of the soils in the Fukushima regions [26].
- Agricultural soil from Fukushima had multiple times lower RIP per unit clay than ordinary European soils
- Secondly, the overestimation of K soil arrangement focuses was likewise credited to the overestimation of k soil arrangement fixations was additionally ascribed to overestimation of other model boundaries, specifically the centralization of calcium and magnesium in the dirt which was multiple times higher than the action esteems. The reliance of Ca and Mg fixation on soil ph. as expected by these models couldn't be shown for the Japanese soils [23].

Software modelling limitations (Data)

Another impediment lies in the way that it's anything but practicable to create a powerful, affectability examination utilizing

the current adaptation model. (further developing models). Muller and Prohl [6] give an primary concern of improbability of the default ecosystem-87 values. This parameter is a site specific [27].

FDMT isn't presently set up to permit the client to settle complex powerful frameworks as basically scientific arrangements are accommodated essential differential conditions and improving on suppositions are made concerning, contributions to and misfortunes from, different parts of the displayed framework. The FDMT has many default parameters which are not founded on data, more transparency is needed on how these qualities have been determined and they ought to be considered with the most recent IAEA information given. (further developing models). On account of move to creatures there is proof that the tissue-diet fixation proportion perhaps a more powerful boundary than the exchange coefficient. (Further developing models and gaining from Post-Fukushima contemplates) [3].

To illustrate some parametric replications using the Rodos model, a system model was developed using Matlab software tools for data testing in the JRodos software. Five radionuclides were considered for this study; Cs 137, Cs 134, Sr 90, Sr89 and Ir131; Food compartments Tubers, Roots, Rice, Fruits, vegetables, Non-leafy vegetables and herbs. Figure 5B indicates the availability of data for Herbs (7,3) and Rice (3,3) in Cs137 concentration.




Figure 6: Simulation of 24hr food chain model on Rodos system (China-HTR10) (A,B).

The JRodos system uses the geographic population by census which do not take into consideration the population of the region by grid. Moreover, the data for population is duely inherent since it does not take into account the site specific population or region data.

The summary for data limitation of the JRodos system

- Not availability of the post Fukushima data
- Use of geographic population by census instead, these Geographic populations do not give an account of people migrated from the place overtime since update of I is not done concurrently.
- Site specific data for analysis and evaluation
- The empirical approach model is adopted in the RODOS system, which gives a complex and not systematic evaluation of data

System design for WSPEEDI

SPEEDI and WSPEEDI are two sorts of PC based choice emotionally supportive networks that have been produced for continuous portion appraisal in radiological crises. SPEEDI is created for homegrown neighborhood range crisis engaged with public scale crisis reaction program connected to Japanese emergency response [22]. The WSPEEDI is a worldwide version which has been created to foresee the radiological effects on the Japanese individuals of a nuclear accident in an a neighbouring country. SPEEDI is generally known as a framework for forecast of ecological crisis portion data [28].

SPEEDI codes are systematized in combination with databases conversational software and graphic software. Data is divided into two parts which are time dependent data like meteorological information and invariant data. Meteorological information and release information are expected to be the input in real-time from a data communication network and keyboard [29]. The worldwide version of SPEEDI was developed as remediation for expansion of area to assess long reach transport of airborne radioactivity because of an extreme atomic mishap in an unfamiliar country. The purpose of SPEEDI is to estimate the long range transport and deposition of airborne radioactivity over synoptic and hemispheric areas [30].

WSPEEDI is a recreation framework that figures air fixation and surface testimony of radionuclides and radiological dosages by progressive utilization of a meteorological expectation model and a Lagrangian molecule scattering model (Refreshed examination of Fukushima unit 3-Fukushima). The WSPEEDI has a limitation for estimating average release over long period of time, that is between 0.5-3.5hrs [22].




Figure 7: Computational flow of SPEEDI and workflow chart of Wspeedi.

Terrestrial model for WSPEEDI

WSPEEDI was created based on SPEEDI. The distinction among SPEEDI and WSPEEDI is that WSPEEDI is intended to reproduce the long-range transport of radionuclides up to the side of the equator scale (Speedi expectation devices) in an upward measurement, the computational area been stretched out to the highest point of the lower atmosphere. In the WSPEEDI a molecule scattering model is utilized to reproduce atmospheric model component. The modelling system of the software was modified with Chernobyl monitoring data [22]. The dispersion and dose model of the WSPEEDI technically follows the same model such as the GEAR (Speedi prediction tools). The radioactive model is expressed with a mass practical with each position been computed in the system. For easy operation of computational codes in emergency situations, SPEEDI codes are systematized in combination with

databases, where the codes are grouped into time dependent such as the meteorological and release information and invariant data [4].




Figure 8: Terrestrial Model of WSPEEDI.

System design of atmospheric release advisory capability (ARAC)

Atmospheric Release Advisory Capability (ARAC) is a system established by Lawrence more national laboratory. The ARAC system uses three dimensional atmospheric transport models to simulate the discharge of contaminants contained in a regional scale flow systems and to prepare calculations for dissemination to local accident emergency response officials [30].

The fundamental goal of the ARAC framework that prompted its origin in 1973 was to give constant expectations of portion levels and degree of surface tainting from mishap arrival of radionuclides from atomic offices. The ARAC framework is utilized by authorities specialists to show portion and testimony computation access, wellbeing perils, define departure plans and concentrate estimation and tidy up endeavors [31].

The various tools and capabilities involve in the ARAC system is the atmospheric transport and diffusion models; the Langrangian operational dispersion integrator which is a three dimensional scattering model created by NARAC to mimic the cycle of shift in weather conditions, tempestuous dispersion, radioactive decay, first-request substance responses, wet statement, gravitational settling, dry testimony and light energy plum rise (NARAC: An emergency reaction source for predicting the atmospheric dispersion) [30].

The Langrangian approach is used more than the Eulerian model because the Langrangian approach avoids numerical dispersion and can resolve point sources [32].




Figure 9: System modelling of the ARAC system.

Software system of ARAC

NARAC's latest version of the software became operational in 2000. It has the client-server system framework with web situated advancements that can deal with various simultaneous clients and occasions. The product utilizes a multi-layered disseminated programming engineering that gives ongoing admittance to the worldwide meteorological and graphical information base atmosphere modelling. It has two working component [4]. The NARAC focal system (NCS) and the NARAC undertaking framework (reference; the public air discharge warning focus demonstrating and choice emotionally supportive network for radiological and atomic crisis readiness and reaction) [28]. In the NARAC central system (NCS), there is a combination of three major subsystems demonstrating and choice framework for radiological and atomic crisis readiness and reaction. In the NARAC Enterprise system, there is a combination of three major subsystems (geospatial, metadata and the modelling execution subsystem) that has the surrounding for advanced scientific analysis and visualization. The metadata subsystem allows temporary and what's more, topographically significant information to be separated for use in the model execution subsystem. NARAC uses maps to show areas where dose limits are exceeded, areas with protective actions (sheltering, evacuation and relocation) thresholds are attained, estimated areas affected population and geographic reference data. (NARAC; An emergency response resource for predicting the atmospheric dispersion and assessing the consequence Michael M. Bradley) [11].

Limitation of the NARAC system

The ARAC now NARAC system specializes in the atmospheric transport and diffusion. (Atmospheric dispersion models). NARAC framework tends to these requirements by giving apparatuses and administrations that foresee and guide the likely spread of unsafe material inadvertently or globally into the climate (The public barometrical delivery warning focus demonstrating and choice emotionally supportive network for radiological and atomic crisis readiness John S. Nasstrom). The main usage of NARAC system is to provide real-time predictions of atmospheric dispersion globally [22].

Atmospheric dispersion models need a source term that portrays attributes, for example, the mass or movement delivered to the environment, the discharge rate, height, spatial appropriation

and molecule size circulation. The NARAC software tools integrate interface with numerous Gaussian plume and puff models [33].




Figure 10: Diagram of the ARAC emergency response.

Analysis of system classification on food chain

Real time consequence assessment model is a valuation tool for emergency response because it can predict the impact on the environment and public. Nuclear accident consequence and decision support system is an important part of nuclear emergency preparedness and is necessary technology in the process of decision [20].




Figure 11: Model chain and data stream of an Emergency Plan and Policy system.

The results illustrate the higher application of the systems in atmospheric modelling and dispersion with Rodos system being the widely used model. In the use in the terrestrial and ecological models its application is sublimely with a lot of influences from the atmospheric dispersion model.

Conclusion

In a nuclear emergency, Protective actions such as evacuation, sheltering and food bans can be taken to prevent the radioactivity release Following the events of the Fukushima accident, it is required to ensure that all maximum system is in placed to in the events of an emergency. New technologies and systems are always in placed to ensure that the public is safe from any unforeseen circumstances. From our study, it can be concluded that Rodos system gives a minimal approach to food safety in the event of emergency, Further studies is needed in the development of food chain models with relevant policy approach and countermeasures for a robust emergency response in the Nuclear Industry.

Bibliography

1. Radiation Monitoring and dose Estimation of the Fukushima Nuclear Accident, Tokyo: Springer (2014).
2. S Y N C D Tarsitano., et al. "Evaluating and reducing a model of radiocaesium soil-plant uptake". *Journal of Environmental Radioactivity* 102.3 (2011): 262-269.
3. SFA KNA Beresford. "Thirty years after the chernobyl accident; what lessons have we learnt". *Journal of Environmental Radioactivity* 157 (2016): 77-89.
4. M Chino., et al. "Speedi and wspeedi: japanese emergency response system to predict radiological impacts in local and worldwide areas due to a nuclear accident". *Radiation Protection Dosimetry* 50 (1993): 145-152.
5. R Mu., et al. "China Approach to Nuclear Safety-from the perspective of policy and institutional system". *Journal of Energy Policy* 76 (2015): 161-172.
6. GH Muller., et al. "ECOSYS-87; a dynamic model for assessing radiological consequences of nuclear accident". *Journal of Health Physics* 64.3 (1993): 232-252.
7. D T M W R I levdin., et al. "RODOS re-engineering; aims and implementation details". *Radioprotection* 45.5 (2010).
8. PA Speed. "The Governance of Nuclear Power in China". *Journal of World Energy Law and Business*, Oxford 13.1 (2020): 23-46.

9. N B A N G C B J Howard., *et al.* "The strategy project; decision tools to aid sustainable restoration and long-term management of contaminated agricultural ecosystems". *Journal of Environmental Radioactive* 83.3 (2005): 275-295.
10. Tecnatom. "www.tecnatom.es". Tecnatom, 18 November (2019).
11. W R W T C Landman., *et al.* "A proposed countermeasure simulation model for the new ICRP recommendation". *EDP Sciences* (2013).
12. A W J Ehrhardt., *et al.* "RODOS; Decision support system for off-site Nuclear Emergency management in Europe, Final Report of the RODOS project". *European Commission* (2000).
13. V K O C L R G T Brit Salbu., *et al.* "Challenges associated with the behaviour of radioactive particles in the environment". *Journal of Environmental Radioactivity* 186 (2018): 101-115.
14. J B H N B H O C T J A F nisbet., *et al.* "Decision aiding handbooks for managing contaminated food production systems, drinking after and inhabited areas in europe". (2010).
15. C W M K K J R E Tipping. "Solid-solution distribution of radionuclides in acid soils; application of the WHAM chemical speciation model". *Environmental Science Technology* 29.5 (1995): 1365-1372.
16. IAEA. "Handbook of Parameters Values for the Prediction of Radionuclide transfer in temperate region". IAEA, Vienna (1994).
17. IAEA. "Handbook of Parameters Values for the prediction of Radionuclide Transfer in terrestrial and freshwater environment". technical reports Series No 472, Vienna (2010).
18. N C J A S W A G Gillett., *et al.* "Temporal and spatial prediction of radiocaesium transfer to food products". *Radiation and Environmental Biophysics* 40 (2001): 227-235.
19. B N R Center. "Biosphere impact studies unit". Boeretang 200, Belgium (2000).
20. S F K N Papamichail., *et al.* "Design and evaluation of an intelligent decision support system for nuclear emergency". *Decision Support Systems* 41.1 (2005): 84-111.
21. F G H Muller. "Documetation of the terrestrial foodchain and dose module FDMT in ROPDOS PV6.0". Rodos Report (2003).
22. "Developing of a worldwide version of system for prediction of environmental emergency dose information; WSPEEDI 111". *Nuclear Science and Technology* (1994): 969-978.
23. L Fernandez-Moguel. "Updated analysis of Fukushima Unit 3 with Melcor 2.1 Part 2; Fission Product Release and Transport Analysis". *Annals of Nuclear Energy* 130 (2019): 93-106.
24. S Y n C J P Absalom. "Predicting soil to plant transfer radiocaesium using soil characteristics". *Environmental Science* 33.8 (1999): 1218-1223.
25. F J A N V W Raskob. "Overview and main achievements of the EURANOS project" (2010).
26. E L J W M V h S Uematsu. "Predicting Radiocaesium sorption characteristics with soil Chemical properties for the japanese soils". *Journal of Science and Environment* 524 (2015): 148-156.
27. J B P B A G M Van Der Perk. "A GIS based environmental Decision support system to assess the transfer of long-lived radiocaesium through food chains in areas contaminated by the chernobyl accident". *International Journal of Geographical Information Science* 15.1 (2001).
28. H ISHIKAWA. "Development of Worldwide Version of System for Prediction of Environmental emergency Dose Information; WSPEEDI (III)". *Nuclear Science and Technology* 31 (1993): 969.
29. S Gerdan. "GIS-based Decision Support system Applications in Disaster Management". 25.3 (2018).
30. M M Bradley. "NARAC: Emergency response resource for predicting the atmospheric dispersion and assessing the consequences of airborne radionuclides". *Journal of Environmental Radioactivity* 96 (2007): 116-121.
31. N A H L J S S H S French. "Presenting Uncertain Information in Radiological Emergencies". UK Atmospheric Dispersion Modelling Laison Committee (2016).
32. G Sugiyama. "Atmospheric Dispersion Modelling: Challenges of the Fukushima Daiichi response".
33. R A M G M Norden. "Bioavailability in the BORIS assessment model". *Radioprotection* 40 (2005): 107-111.
34. F Gering. "Data Assimilation Methods for Improving the Prognoses of Radionuclide Deposition from Radioecological Models with Measurements". *Dissertation* (2005).

35. V B M T J Geldermann. "Multi- Criteria decision support and Evaluation of Strategies for nuclear Remediation management". *Omega* 37.1 (2009): 238-251.
36. P Speed. "The Governance of Nuclear Power In China". *Journal of world Energy Law and Business*, Oxford (2020).
37. J Mu. "China's Approach to nuclear safety- From the perspective of policy and Institutional system". *Journal of Energy Policy*, China (2014).
38. N B Howard., *et al.* "The Strategy Project; Decision tools to aid sustainable restoration and long term management of contaminated agricultural ecosystems". *Journal of Environmental Radioactivity* 83.3 (2005): 275-295.
39. W C Landman., *et al.* "A Proposed Countermeasure Simulation Model for the new ICRP Recommendations". *EDP Sciences* (2013).
40. A Ehrhardt. "RODOS; Decision Support System for off-site Nuclear Emergency management in Europe, Final Report of the RODOS Project". *European Commission* (2000).
41. F V B W Raskob. "Approaches o Visualisation of uncertainties to decision makers in an operational decision support system". Gothenburg, Proceedings of the 6th International Conference on Information systems for Crisis Response and management (2009).
42. GH Muller., *et al.* "ECOSYS-87; a dynamic model for assessing radiological consequences of nuclear accident". *Journal of Health Physics* 64.3 (1993): 232-252.