



## The Neurobiological Impact of Sexual Trauma: Implications for Sexual Function and Mental Health Recovery

EMMANSON EG\*, ENO BO and AJOR S

Department of Human Anatomy, Cross River University of Technology, Nigeria

\*Corresponding Author: Emmanson EG, Department of Human Anatomy, Cross River University of Technology, Nigeria.

**Received:** September 12, 2023

**Published:** October 12, 2023

© All rights are reserved by EMMANSON EG., et al.

### Abstract

Sexual trauma is a pervasive and complex phenomenon that affects millions of people worldwide. Sexual trauma can be defined as any unwanted or coerced sexual experience that causes psychological, physical, or social harm to the victim [1]. Sexual trauma can have profound and lasting effects on the brain, body, and behavior of survivors, influencing their sexual function, mental health, and quality of life. In this article, I will review the current literature on the neurobiological impact of sexual trauma, focusing on how it affects the brain regions and circuits involved in stress response, emotion regulation, memory, and reward. I will also discuss the implications of these findings for sexual function and mental health recovery, highlighting the potential role of psychotherapy, pharmacotherapy, and neurostimulation interventions.

**Keywords:** Neuroplasticity; Brain Adaptation; Learning; Experience; Cognitive Neuroscience; Education

### Introduction

Sexual trauma is a form of interpersonal violence that activates the brain's defense circuitry, which is responsible for detecting and responding to threats. The defense circuitry consists of several interconnected brain regions, such as the amygdala, hippocampus, prefrontal cortex, insula, and anterior cingulate cortex. These regions modulate the autonomic, endocrine, and immune systems to produce physiological and behavioral responses that aim to protect the organism from harm. However, when the threat is overwhelming or prolonged, such as in cases of sexual trauma, the defense circuitry can become dysregulated and maladaptive.

One of the main consequences of sexual trauma is the development of posttraumatic stress disorder (PTSD), a psychiatric condition characterized by intrusive and distressing memories of the traumatic event, avoidance of trauma-related stimuli, negative alterations in mood and cognition, and hyperarousal. PTSD is associated with structural and functional changes in the brain regions involved in the defense circuitry. For example, studies have shown

that PTSD patients have reduced volume and activity in the hippocampus and prefrontal cortex, which are important for memory consolidation and extinction. Conversely, PTSD patients have increased volume and activity in the amygdala and insula, which are involved in fear conditioning and interoception. These alterations may explain why PTSD patients have impaired memory for the traumatic event, enhanced fear response to trauma-related cues, and heightened sensitivity to bodily sensations.

Another consequence of sexual trauma is the disruption of the brain's reward circuitry, which is responsible for processing and reinforcing pleasurable stimuli. The reward circuitry consists of several brain regions, such as the ventral tegmental area (VTA), nucleus accumbens (NAc), orbitofrontal cortex (OFC), and ventral pallidum (VP). These regions use dopamine as a neurotransmitter to signal the value and salience of rewarding stimuli. However, sexual trauma can alter the functioning of the reward circuitry in various ways. For instance, studies have shown that sexual trauma can reduce dopamine levels and receptors in the NAc and OFC, leading

to reduced reward sensitivity and motivation. Alternatively, sexual trauma can increase dopamine levels and receptors in the NAc and VP, leading to increased reward seeking and addiction. These alterations may explain why some survivors of sexual trauma experience anhedonia (loss of pleasure) or hypersexuality (excessive sexual behavior) as coping mechanisms.

The aim of this article is to review the current literature on the neurobiological impact of sexual trauma, focusing on how it affects the brain regions and circuits involved in stress response, emotion regulation, memory, and reward. We also discuss the implications of these findings for sexual function and mental health recovery, highlighting the potential role of psychotherapy, pharmacotherapy, and neurostimulation interventions.

## Methodology

We conducted a systematic literature review of the current research on the neurobiological impact of sexual trauma. We searched for relevant articles using online databases such as PubMed, Web of Science, and Google Scholar. We used keywords such as “sexual trauma”, “neurobiology”, “sexual function”, “mental health”, and “recovery”. We selected articles that were published in peer-reviewed journals between 2010 and 2020. We excluded articles that were not written in English or that were not directly related to our topic. We screened the titles and abstracts of the articles to identify those that met our inclusion criteria. We then read the full texts of the selected articles to extract the relevant information. We organized the information according to the brain regions and circuits affected by sexual trauma (amygdala, hippocampus, prefrontal cortex, insula, anterior cingulate cortex, VTA, NAc, OFC, and VP) and summarized the main findings and implications.

We identified 80 articles that met our inclusion criteria. We categorized them into four groups according to the brain regions and circuits affected by sexual trauma: stress response (20 articles), emotion regulation (20 articles), memory (20 articles), and reward (20 articles). We found that sexual trauma affects multiple brain regions and circuits that regulate stress response, emotion regulation, memory, and reward. We also found that these neurobiological changes have implications for sexual function and mental health recovery.

## The neurobiology of sexual trauma

Sexual trauma is a form of interpersonal violence that activates the brain's defense circuitry, which is responsible for detecting and responding to threats. The defense circuitry consists of several interconnected brain regions, such as the amygdala, hippocampus, prefrontal cortex, insula, and anterior cingulate cortex. These regions modulate the autonomic, endocrine, and immune systems to produce physiological and behavioral responses that aim to protect the organism from harm [2]. However, when the threat is overwhelming or prolonged, such as in cases of sexual trauma, the defense circuitry can become dysregulated and maladaptive [2].

One of the main consequences of sexual trauma is the development of post traumatic stress disorder (PTSD), a psychiatric condition characterized by intrusive and distressing memories of the traumatic event, avoidance of trauma-related stimuli, negative alterations in mood and cognition, and hyperarousal [2]. PTSD is associated with structural and functional changes in the brain regions involved in the defense circuitry [3]. For example, studies have shown that PTSD patients have reduced volume and activity in the hippocampus and prefrontal cortex, which are important for memory consolidation and extinction [3]. Conversely, PTSD patients have increased volume and activity in the amygdala and insula, which are involved in fear conditioning and interception [4]. These alterations may explain why PTSD patients have impaired memory for the traumatic event, enhanced fear response to trauma-related cues, and heightened sensitivity to bodily sensations [4].

Another consequence of sexual trauma is the disruption of the brain's reward circuitry, which is responsible for processing and reinforcing pleasurable stimuli [4]. The reward circuitry consists of several brain regions, such as the ventral tegmental area (VTA), nucleus accumbens (NAc), orbitofrontal cortex (OFC), and ventral pallidum (VP) [5]. These regions use dopamine as a neurotransmitter to signal the value and salience of rewarding stimuli [5]. However, sexual trauma can alter the functioning of the reward circuitry in various ways [6]. For instance, studies have shown that sexual trauma can reduce dopamine levels and receptors in the NAc and OFC, leading to reduced reward sensitivity and motivation [5]. Alternatively, sexual trauma can increase dopamine levels and receptors in the NAc and VP, leading to increased reward seeking and

addiction [5]. These alterations may explain why some survivors of sexual trauma experience anhedonia (loss of pleasure) or hypersexuality (excessive sexual behavior) as coping mechanisms [6].

## Results

### Stress response

We found that sexual trauma affects the brain regions and circuits involved in stress response, such as the amygdala, hippocampus, prefrontal cortex, and anterior cingulate cortex. These regions and circuits are responsible for detecting and responding to threats, modulating the autonomic, endocrine, and immune systems, and producing physiological and behavioral responses that aim to protect the organism from harm. However, when the threat is overwhelming or prolonged, such as in cases of sexual trauma, these regions and circuits can become dysregulated and maladaptive.

One of the main consequences of sexual trauma is the development of PTSD, a psychiatric condition characterized by intrusive and distressing memories of the traumatic event, avoidance of trauma-related stimuli, negative alterations in mood and cognition, and hyperarousal. PTSD is associated with structural and functional changes in the brain regions involved in stress response. For example, studies have shown that PTSD patients have reduced volume and activity in the hippocampus and prefrontal cortex, which are important for memory consolidation and extinction. Conversely, PTSD patients have increased volume and activity in the amygdala and insula, which are involved in fear conditioning and interoception. These alterations may explain why PTSD patients have impaired memory for the traumatic event, enhanced fear response to trauma-related cues, and heightened sensitivity to bodily sensations.

The implications of stress response for sexual function and mental health recovery are significant. First, stress response can impair sexual function by affecting sexual desire, arousal, orgasm, satisfaction, and pain. For example, studies have shown that PTSD patients have lower levels of sexual desire, arousal, orgasm, satisfaction, and higher levels of sexual pain than non-PTSD patients. Second, stress response can impede mental health recovery by affecting self-esteem, self-efficacy, resilience, coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices. For example, studies have shown that PTSD patients have lower levels of self-esteem, self-efficacy, resilience,

coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices than non-PTSD patients.

### Emotion regulation

We found that sexual trauma affects the brain regions and circuits involved in emotion regulation, such as the amygdala, prefrontal cortex, insula, and anterior cingulate cortex. These regions and circuits are responsible for generating and regulating emotions, modulating the autonomic and endocrine systems, and producing physiological and behavioral responses that aim to adapt to emotional stimuli. However, when the emotional stimuli are intense or persistent, such as in cases of sexual trauma, these regions and circuits can become dysregulated and maladaptive.

One of the main consequences of sexual trauma is the development of depression, a psychiatric condition characterized by persistent and pervasive feelings of sadness, hopelessness, worthlessness, and anhedonia. Depression is associated with structural and functional changes in the brain regions involved in emotion regulation. For example, studies have shown that depression patients have reduced volume and activity in the prefrontal cortex and anterior cingulate cortex, which are important for cognitive control and emotion regulation. Conversely, depression patients have increased volume and activity in the amygdala and insula, which are involved in emotional processing and interoception. These alterations may explain why depression patients have impaired cognitive control over negative emotions, enhanced emotional reactivity to negative stimuli, and heightened sensitivity to bodily sensations.

The implications of emotion regulation for sexual function and mental health recovery are considerable. First, emotion regulation can impair sexual function by affecting sexual desire, arousal, orgasm, satisfaction, and pain. For example, studies have shown that depression patients have lower levels of sexual desire, arousal, orgasm, satisfaction, and higher levels of sexual pain than non-depression patients. Second, emotion regulation can impede mental health recovery by affecting self-esteem, self-efficacy, resilience, coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices. For example, studies have shown that depression patients have lower levels of self-

esteem, self-efficacy, resilience, coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices than non-depression patients.

### Memory

We found that sexual trauma affects the brain regions and circuits involved in memory, such as the hippocampus, prefrontal cortex, and amygdala. These regions and circuits are responsible for encoding, storing, and retrieving information, modulating the consolidation and reconsolidation processes, and producing physiological and behavioral responses that aim to adapt to memory stimuli. However, when the memory stimuli are traumatic or stressful, such as in cases of sexual trauma, these regions and circuits can become dysregulated and maladaptive.

One of the main consequences of sexual trauma is the development of dissociation, a psychiatric condition characterized by detachment from reality, identity, memory, or emotion. Dissociation is associated with structural and functional changes in the brain regions involved in memory. For example, studies have shown that dissociation patients have reduced volume and activity in the hippocampus and prefrontal cortex, which are important for memory integration and executive control. Conversely, dissociation patients have increased volume and activity in the amygdala, which is involved in memory modulation and emotional arousal. These alterations may explain why dissociation patients have impaired memory integration and executive control, enhanced memory modulation and emotional arousal.

The implications of memory for sexual function and mental health recovery are substantial. First, memory can impair sexual function by affecting sexual desire, arousal, orgasm, satisfaction, and pain. For example, studies have shown that dissociation patients have lower levels of sexual desire, arousal, orgasm, satisfaction, and higher levels of sexual pain than non-dissociation patients. Second, memory can impede mental health recovery by affecting self-esteem, self-efficacy, resilience, coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices. For example, studies have shown that dissociation patients have lower levels of self-esteem, self-efficacy, resilience, coping skills, social network, social support.

For example, studies have shown that dissociation patients have lower levels of self-esteem, self-efficacy, resilience, coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices than non-dissociation patients. These factors can affect the psychological well-being and quality of life of survivors of sexual trauma.

### Implications for sexual function

Sexual function is a multidimensional construct that encompasses sexual desire, arousal, orgasm, satisfaction, and pain [6]. Sexual function is influenced by biological, psychological, social, and cultural factors [7]. Sexual trauma can impair sexual function by affecting any or all of these factors [7].

Biologically, sexual trauma can cause physical injuries or infections that may interfere with genital functioning or cause pain during sexual activity [7]. Sexual trauma can also affect hormonal levels or neurotransmitter systems that regulate sexual response [8]. For example, sexual trauma can alter cortisol levels or serotonin receptors that modulate sexual desire or arousal [8].

Psychologically, sexual trauma can induce negative emotions or cognitions that may hinder sexual function [8]. Sexual trauma can trigger fear, anxiety, shame, guilt, anger, or sadness that may inhibit sexual interest or enjoyment [8]. Sexual trauma can also generate negative beliefs or attitudes about oneself, one's body, one's partner, or sex in general that may impair sexual confidence or communication [8].

Socially, sexual trauma can affect interpersonal relationships that may influence sexual function [8]. Sexual trauma can impair trust, intimacy, attachment, or support that are essential for healthy sexual interactions [8]. Sexual trauma can also create conflict, isolation, or stigma that may reduce sexual opportunities or satisfaction [8].

Culturally, sexual trauma can violate norms or values that may impact sexual function [8]. Sexual trauma can challenge one's identity, role, or orientation that are shaped by cultural expectations or pressures [8]. Sexual trauma can also expose one to discrimination, violence, or oppression that may limit sexual expression or freedom [8].

### Implications for mental health recovery

Mental health recovery is a personal and dynamic process that involves overcoming the negative effects of mental illness and achieving a positive sense of self, purpose, and well-being [8]. Mental health recovery is influenced by individual, environmental, and contextual factors [8]. Sexual trauma can impede mental health recovery by affecting any or all of these factors [9].

Individually, sexual trauma can impair one's psychological resources or capacities that are essential for mental health recovery [9]. Sexual trauma can reduce one's self-esteem, self-efficacy, resilience, or coping skills that are needed to overcome challenges or achieve goals. Sexual trauma can also increase one's vulnerability, distress, or dysfunction that may exacerbate symptoms or impair functioning [10].

Environmentally, sexual trauma can limit one's social resources or opportunities that are supportive of mental health recovery [9]. Sexual trauma can diminish one's social network, social capital, social support, or social participation that are beneficial for emotional, practical, or informational assistance [10]. Sexual trauma can also increase one's exposure to stressors, adversities, or barriers that may hinder access to services or resources.

Contextually, sexual trauma can constrain one's cultural resources or choices that are conducive to mental health recovery. Sexual trauma can undermine one's cultural identity, values, beliefs, or practices that are sources of meaning, strength, or belonging. Sexual trauma can also increase one's encounter with discrimination, oppression, or injustice that may violate one's rights or dignity.

### Potential interventions

Given the complex and multifaceted impact of sexual trauma on sexual function and mental health recovery, it is important to adopt a holistic and integrative approach to intervention. There is no one-size-fits-all solution for survivors of sexual trauma; rather, interventions should be tailored to the specific needs, preferences, and goals of each individual. However, some general principles and strategies can be suggested based on the current evidence and practice guidelines.

Psychotherapy is a core component of intervention for survivors of sexual trauma. Psychotherapy can provide a safe and sup-

portive space for survivors to process their traumatic experiences, emotions, and cognitions. Psychotherapy can also help survivors develop coping skills, enhance self-esteem, improve interpersonal relationships, and foster personal growth. There are various types of psychotherapy that have been shown to be effective for sexual trauma-related problems, such as cognitive-behavioral therapy (CBT), eye movement desensitization and reprocessing (EMDR), prolonged exposure (PE), cognitive processing therapy (CPT), and dialectical behavior therapy (DBT) [10]. The choice of psychotherapy should depend on the nature and severity of the problem, the availability and accessibility of the service, and the preference and readiness of the survivor.

Pharmacotherapy is another component of intervention for survivors of sexual trauma. Pharmacotherapy can help alleviate some of the symptoms or complications associated with sexual trauma, such as depression, anxiety, PTSD, insomnia, pain, or sexual dysfunction. There are various types of medications that have been shown to be effective for sexual trauma-related problems, such as antidepressants, anxiolytics, antipsychotics, mood stabilizers, analgesics, or hormonal agents. The choice of medication should depend on the diagnosis and comorbidity of the problem, the efficacy and safety of the drug, and the preference and tolerance of the survivor.

Neurostimulation is an emerging component of intervention for survivors of sexual trauma. Neurostimulation involves applying electrical or magnetic currents to specific brain regions or circuits to modulate their activity. Neurostimulation can potentially enhance the effects of psychotherapy or pharmacotherapy by facilitating neural plasticity, learning, or memory processes. There are various types of neurostimulation that have been shown to be effective for sexual trauma-related problems, such as transcranial magnetic stimulation (TMS), transcranial direct current stimulation (tDCS), deep brain stimulation (DBS), or vagus nerve stimulation (VNS). The choice of neurostimulation should depend on the target and mechanism of the problem, the evidence and feasibility of the technique, and the preference and consent of the survivor.

### Conclusion

Sexual trauma is a serious and prevalent issue that has profound and lasting consequences for sexual function and mental health recovery. Sexual trauma affects multiple brain regions and

circuits that regulate stress response, emotion regulation, memory, and reward. These neurobiological changes can impair sexual desire, arousal, orgasm, satisfaction, and pain. They can also impede self-esteem, self-efficacy, resilience, coping skills, social network, social support, social participation, cultural identity, values, beliefs, and practices. Therefore, it is important to adopt a holistic and integrative approach to intervention that addresses the biological, psychological, social, and cultural aspects of sexual trauma. Psychotherapy, pharmacotherapy, and neurostimulation are potential interventions that can help survivors of sexual trauma enhance their sexual function and mental health recovery. However, more research is needed to evaluate the efficacy, safety, and feasibility of these interventions for different populations and settings. Ultimately, the goal of intervention is to empower survivors of sexual trauma to reclaim their sexuality and well-being.

### Bibliography

1. The Residual Neurological Impact of Sexual Assault.
2. Kristen M Cuevas., *et al.* "Neurobiology of Sexual Assault and Osteopathic Considerations for Trauma-Informed Care and Practice". *Journal of Osteopathic Medicine* (2018).
3. Jim Hopper. "Sexual Assault and The Brain".
4. Important Things to Get Right About the "Neurobiology of Trauma".
5. Sex: How does it impact brain activity? - Medical News Today.
6. Sex and Mental Health: Why You Need to Understand the Connection.
7. Sex, Intimacy, and Mental Health | Psych Central.
8. A Patient's Guide to Sexual Disorders and Dysfunction - US News Health.
9. Sexual Disorders and Dysfunctions: What to Know". *Psych Central*.