

Assessment of Knowledge of Cervical Cancer Transmission and Prevention among the Mothers of Daughters Aged Below 10 Years

Sharmin Zaman Khan*

Green Life Medical College, Bangladesh

*Corresponding Author: Sharmin Zaman Khan, Green Life Medical College, Bangladesh.

Received: August 10, 2018; Published: August 14, 2018

Abstract

Introduction: Cervical Cancer is the second most common cancer among women worldwide. It is one of the few preventable human cancers. Maternal knowledge is required for vaccination of their adolescent girls because they are key decision makers and potentially a major source of information for their daughters. Bangladesh Government will introduce HPV vaccine under EPI program for adolescent girls aged 10 years from 2018. For this reason, it is important to assess the knowledge level of mothers of daughters aged below 10 years.

Objective: The objective of this study was to assess the knowledge of Cervical Cancer transmission, prevention and HPV vaccine among mothers of daughters aged below 10 years.

Method: A cross sectional descriptive study has been conducted among the mothers of daughters aged below 10 years, live in the Mohakhali 'Sat Tola' slum area of Dhaka city. The pre-tested questionnaire was used to collect information from the respondents by face to face interview at house-hold level. Questions were on socio-demographic characteristics, knowledge about Cervical Cancer transmission, prevention and HPV vaccination. Convenient sampling technique was used to select the sample. The sample size was 100 in number. The thesis protocol was approved by Institutional Review Board of BSMMU. The total questions on knowledge was 22. Knowledge score was divided into 3 category: poor, average and good knowledge. Poor knowledge score was 0 to 7, average knowledge score was 8 to 14 and good knowledge score was 15 to 22. Frequency, percentage, mean and chi-square test was done by using SPSS version 21.

Result: Majority respondents' age was 21 to 29 years, most of them were primary school educated, most of them were house-wife and monthly family income was 11,000 to 15,000 taka. Respondent's mean age at marriage was 15 years and mean age was 17 years when their first child born, majority had two children. The respondents' mean knowledge score was 7.19 out of 22. In total, 55% respondents had poor knowledge, 45% respondents had average knowledge and 2% respondents had good knowledge. Among them 98% respondents heard about cancer, 77% respondents heard about Cervical Cancer, 8% respondents knew about sign and symptom and only 2% respondents knew the causes of Cervical Cancer. On the other hand, 3% respondents told prevention of Cervical Cancer is possible but 12% respondents heard about VIA test. Around 18% respondents heard about HPV vaccine but nobody knew in which age this vaccine should be administered. All respondents wanted information about Cervical Cancer and HPV vaccine.

Conclusion: The knowledge of Cervical Cancer transmission, prevention and HPV vaccination among mothers who live in slum of Dhaka city is poor. Providing knowledge to these mothers is essential.

Keywords: Cervical Cancer Transmission; Mothers; Daughters; HPV Vaccination

Introduction

Cervical Cancer is one of the few preventable human cancers [1]. It is the second most common cancer among women worldwide and about 86% of Cervical Cancer occur in developing countries [2]. More than half a million women are newly diagnosed with Cervical Cancer each year, with approximately 90% of Cervical Cancer-related death occurring in developing countries [3]. However,

in developing countries, it remains the most common cause of cancer related deaths among women. Cervical Cancer which is one of the preventable cancers and the important public health problem worldwide and especially in developing countries [4].

Bangladesh, at 142 million people, is the 9th most populous country in the world. There are 13 to 15 lakh cancer patients in Bangladesh, with about 2 lakh patients newly diagnosed with can-

cer each year. As an overview, in women, Cervical Cancer and breast cancer are most prevalent [5]. Bangladesh has annual incidence of Cervical Cancer is about 11,956. According to World Health Organization (WHO) statistics, incidence of Cervical Cancer cases in Bangladesh has been estimated at 167 per 100,000 populations and 6,582 women die every year in the country for this cancer. The problem of Cervical Cancer in Bangladesh is particularly acute because of poverty, early age marriage, multiple marriages, high parity and illiteracy. In developing countries, women's knowledge about risk factors of Cervical Cancer is very limited. However, it is considered one of the most preventable cancers [6].

Despite the high prevalence of Cervical Cancer, many studies have shown that women's knowledge about HPV and Cervical Cancer is very low [1]. The incidence of Human Papilloma Virus (HPV) infection and Cervical Cancer can be reduced by increasing vaccination for HPV [7]. In Bangladesh, there is little awareness among the general population, health care professionals and policy makers about Cervical Cancer prevention. Immunization against the most prevalent HPV types affecting the country may be an efficient means to long term prevention [8]. Maternal knowledge is required for vaccination of adolescent girls, they are key decision makers and potentially a major source of information for their daughters [9]. Therefore, improving the knowledge of the mothers who have daughters aged below 10 years regarding Cervical Cancer transmission, prevention and organism is one of the most important steps in enhancing the vaccination coverage among Bangladeshi females.

Objective of the Study

The objective of this study was to determine the knowledge of Cervical Cancer transmission, prevention and organism among the mothers of any slum area of Dhaka city to identify the main barrier to get information about Cervical Cancer, so that possible interventions could be taken to increase the level of knowledge of mothers of Cervical Cancer.

Methods

This was a cross-sectional descriptive study design. This study was done in 'Sat Tola' slum in Mohakhali, Dhaka, Bangladesh. The study was conducted from January, 2016 to December, 2016. Data collection was done for four weeks. Population was the mothers of daughters aged below 10 years, were defined as study population. Mothers who have daughters aged below 10 years were taken because, Cervical Cancer vaccine is given between 12 years to 18 years, so before 10 years if a mother know about Cervical Cancer

and its vaccine then she can took her daughters for vaccine for Cervical Cancer prevention. The mothers were selected by using convenient sampling technique. The mothers who gave voluntary consent were taken as sample. Inclusion criteria is mothers who have daughters aged below 10 years and mothers who are willing to participate in the study. sample size was 100. Pre-tested questionnaire was used to collect information from the respondents by face to face interview at house hold level. The study was performed after getting clearance from Institutional Review Board of BSMMU. Written permission was taken from ward counselor of ward no-20 for data collection. Informed written consent was taken from the mothers before taking interview. After data collection data entry was done. Then data editing, cleaning and data analysis was performed. Statistical data analysis was done using SPSS software version 21.0. After completion of data collection the researcher visited the study area and distributed a leaflet, which contains information about Cervical Cancer. This component is not included in a part of research.

Results

In table 1 shown, 41% respondents' age was between 21 - 29 years, 63% were primary school educated, 80% were house-wife, 43% respondents' monthly family income was 11,000 - 15,000 taka, mean age at marriage was 14.62 ± 2.673 years, mean age during first child born was 17.13 ± 2.866 years, 39% respondent had two children, mean age of last child was 63.4 ± 39.92 months and mean knowledge score was 7.19 out of 22.

In table 2 shown, 98% respondents heard about Cancer but only 77% mothers heard about Cervical Cancer where 54.9% respondents heard from neighbor. Only 8% respondents had knowledge about sign symptom and only 2% respondents had knowledge about causes of Cervical Cancer.

In table 3 shown, 12% respondents heard about VIA test, only 3% check-up Cervix by VIA test, 18% respondents heard about Cervical Cancer vaccine. Among them 55.6% heard from health worker. Only 2% respondents took vaccine and only 6% heard about Human Papilloma Virus.

In figure 1 shown, 60% respondents did not know whether Cervical Cancer prevention is possible or not, 36% told prevention is possible and 4% told prevention is not possible.

In figure 2 shown, 87 respondents wanted to know about Cervical Cancer from health worker, 12 respondents from hospital and 7 respondents from television.

Socio-demographic characteristics	Frequency (n)	Percentage (%)
Age of respondents in year		
< 20	17	17%
21 - 29	41	41%
30 - 39	30	30%
40 and above	11	11%
Don't know	1	1%
Mean age of respondents in year ± SD 29.08 ± 9.621		
Educational status		
No formal education	33	33%
Primary	63	63%
Secondary and above	4	4%
Employment status		
House wife	80	80%
Service holder	10	10%
Business	2	2%
Maid servant	8	8%
Monthly house-hold income in taka		
< 5000	15	15%
6,000 - 10,000	38	38%
11,000 - 15,000	43	43%
> 15,000	4	4%
Age at marriage of respondents in year (mean ± SD) 14.62 ± 2.673		
Age during first child born in year (mean ± SD) 17.13 ± 2.866		
Number of children of respondents		
One	28	28%
Two	39	39%
Three	17	17%
Four	8	8%
Five or more than five	8	8%
Age of last child in month (mean ± SD) 63.4 ± 39.92		
Knowledge score (mean) 7.19 out of 22		

Table 1: Distribution of respondents by socio-demographic characteristics (n = 100).

This study used a questionnaire, consists of 22 questions about knowledge of Cervical Cancer, Cervical Cancer transmission, prevention and HPV. Among the questions 7 questions about knowledge of Cervical Cancer and Cervical Cancer transmission, 11 questions about knowledge of Cervical Cancer prevention and 4 questions about knowledge of HPV. Each question was scored ‘1’ point for the correct response and ‘0’ for the wrong answer. An overall total

Variables	Frequency (n)	Percentage (%)
Heard about Cancer		
Yes	98	98%
No	2	2%
Heard about Cervical Cancer		
Yes	77	77%
No	23	23%
Sources of knowledge about Cervical Cancer (n = 77)		
Health worker	14	17.1%
Television	10	12.2%
Neighbor	45	54.9%
Relatives	13	15.9%
Knowledge about sign symptom of Cervical Cancer (N = 100)		
Yes	8	8%
No	92	92%
Knowledge about causes of Cervical Cancer (N = 100)		
Yes	2	2%
No	92	92%

Table 2: Knowledge about Cervical Cancer transmission (N = 100).

Variables	Frequency (n)	Percentage (%)
Heard about VIA test		
Yes	12	12%
No	88	88%
Check-up Cervix by VIA test		
Yes	3	3%
No	97	97%
Heard about Cervical Cancer vaccine		
Yes	18	18%
No	82	82%
Source of knowledge about Cervical Cancer vaccine (n = 18)		
Health worker	10	55.6%
Television	1	5.6%
Neighbor	5	27.8%
Relative	2	11.1%
Take Cervical Cancer vaccine		
Yes	2	2%
No	98	98%
Heard about Human Papilloma Virus		
Yes	6	6%
No	94	94%

Table 3: Knowledge about Cervical Cancer prevention and organism (N = 100).




Figure 1: Opinion about prevention of Cervical Cancer.




Figure 2: Desire source of knowledge..

knowledge score was calculated by adding up the correct score of each respondent across all 22 questions. Knowledge score was divided into 3 categories which is described below.

In table 4 shown, knowledge score is divided in three categories.

Category	Range of knowledge score
Poor knowledge	0 to 7
Average knowledge	8 to 14
Good knowledge	15 to 22

Table 4: Category of knowledge score of respondents.

In figure 3 shown, 55% respondents had poor knowledge, 43% had average knowledge and only 2% had good knowledge.

Figure 3: Knowledge score of respondents (n = 100).

Discussion

Knowledge on Cervical Cancer among different groups, professionals, populations and students were evaluated by survey worldwide in several studies. In Bangladesh, socio-epidemiological data on Cervical Cancer status are scarce, particularly Cervical Cancer knowledge of mothers are not available at national level. The present study intended to provide such information among the mothers of daughters aged below 10 years in a slum of Dhaka city in Bangladesh. The mean age of the study subjects was 29.08 ± 9.621 years. In a study conducted by Fernandez., *et al.* [7], it was seen that mothers’ mean age was 47.9 ± 5.5 years. In our study, 63% mothers had primary education, 3% mothers had secondary and 1% mothers had higher secondary education. In a study conducted by Kose., *et al.* [4], 63% of the mothers were primary school, 25.7% were high school and 11.4% were university graduates. In this study, it was found out that 77% mothers heard of Cervical Cancer. In a study conducted by Ahmed., *et al.* [10], it was seen that 66.9% of the mothers had heard of Cervical Cancer. In our study, only 2% mothers knew about causes of Cervical Cancer where in a study, conducted at Nigeria by Ahmed., *et al.* [10], it was seen that 62.5% mothers knew about causes of Cervical Cancer. In our study, 12% mothers heard about VIA test and 18% mothers heard about HPV vaccine. In other studies conducted it is seen that 83.5% of the mothers [4] and 53.7% of the mothers [11] had no knowledge about HPV vaccination. Among 18 mothers who expressed that they knew about vaccination, 55.6% (n = 10) stated that they learned about it from a health worker, 27.8% (n = 5) of them learned it from neighbor, 11.1% (n = 2) learned it from relative and 5.6% (n = 1) learned it from television. In a study conducted by Kose., *et al.* [4], 27.1% (n = 35) stated that

they learned about it from a health professional, 61.2% (n = 79) of them learned it from communication tools like television, radio, newspapers or internet, 11.6% (n = 15) learned it from friends or relatives. In our study, only 2% (n = 2) mothers took HPV vaccine, 100% (n = 100) among 100 mothers did not know the age of HPV vaccine administration. In a study conducted by Kose., *et al.* [4], found out that a total of 0.7% mothers who had daughters were found out to be vaccinated against HPV, 83.2% of the mothers did not know age of vaccination was applied. In our study, it was seen that 94% of the mothers who have daughters aged below 10 years had never heard of HPV before. In a study conducted by Marlow., *et al.* [12], it was seen that HPV knowledge 72.8% of the mothers who have daughters between the ages of 8 - 14 had never heard of HPV before. In our study the knowledge score of Cervical Cancer showed that there was good knowledge (2%) about Cervical Cancer among the mothers, living in slum area. Study conducted by Ahmed., *et al.* [10], where the knowledge score of Cervical Cancer showed that there was a fair knowledge (43.5%) of Cervical Cancer among respondents living in slum area. In this study there was no significant association between mothers' education level and their knowledge score. In a similar study conducted by Ozan., *et al.* [13], there was no significant difference. However, in a similar study conducted by Kose., *et al.* [4], there was a significant association between the mothers' education level and their knowledge. According to mothers' economic conditions, there is no significant association between their knowledge about HPV and economic condition. However, there was a significant relationship between high income and the mothers' awareness of HPV was found by Sanders Thompson., *et al* [14].

Conclusion and Recommendation

This study showed that most of the respondents heard about Cancer and Cervical Cancer, among mothers who heard about Cervical Cancer most of them heard from neighbor. But most of the respondents did not know about sign symptoms and the causes of Cervical Cancer. Majority of the respondents did not know whether Cervical Cancer prevention is possible or not. In this study, study was done on mothers who have daughters aged below 10 years at slum area of Dhaka city where under-privileged people lived, most of them uneducated, from lower socio-economic condition, they live there in unhygienic environment and they are deprived from health services such as home visit from female health worker from government sector that's why the mothers cannot get proper information about health so if we want to increase the knowledge level of mothers living in a slum area of Dhaka city at first we have to provide

information to increase awareness among mothers about Cervical Cancer, Cervical Cancer transmission, prevention and HPV. The mothers who had some information about Cervical Cancer stated that they learned about the subject from neighbor, health worker, relatives and television. Based on this result, it can be suggested that doctors and nurses should be informed about this situation and mothers be trained by them on this subject. Despite low knowledge of Cervical Cancer, all mothers were willing to get information about this disease, where majority of them wanted to get information from health worker. We recommend improving mothers' knowledge through health education.

Bibliography

1. Assoumou SZ., *et al.* "Awareness and knowledge regarding of Cervical Cancer, pap smear screening and Human Papilloma Virus infection in Gabonese women". *Journal of Negative Results in Biomedicine* 15 (2015): 37.
2. Johnson DC., *et al.* "Knowledge and awareness of Human Papilloma Virus (HPV), Cervical Cancer and HPV vaccine among women in two distinct Nepali communities". *Asian Pacific Journal of Cancer Prevention* 15.19 (2014): 8287- 8293.
3. Kim K., *et al.* "Knowledge perceptions and decision making about Human Papilloma Virus vaccination among Korean American women: A focus group study". *Women's Health Issues* 25.2 (2015): 112-119.
4. Kose D., *et al.* "Mothers' knowledge and attitudes about HPV vaccination to prevent Cervical Cancer". *Asian Pacific Journal of Cancer Prevention* 15.17 (2014): 7263-7266.
5. Hussain SMA. "Comprehensive update on cancer scenario of Bangladesh". *South Asian Journal of Cancer* 2.4 (2013): 279-284.
6. Papri FS., *et al.* "Knowledge and awareness about risk factors of Cervical Cancer, its screening and vaccination among the women attending Chittagong Medical College Hospital". *Chattagram Maa-O-Shishu Hospital Medical College Journal* 14.2 (2015): 57-60.
7. Fernandez ME., *et al.* "Knowledge, attitudes and beliefs about Human Papilloma Virus (HPV) vaccination among Puerto Rican mothers and daughters: a qualitative study". *Preventing Chronic Disease* 11 (2014): E212.
8. Banik U., *et al.* "High risk Human Papilloma Virus type 16 and 18 infection in the cervical lesions of women with epithelial cell abnormality in pap smear: A cytohistomorphologic association in Bangladeshi women". *Cyto Journal* 10 (2013): 14.

9. Haesebaert J., *et al.* "French women's knowledge and attitudes towards Cervical Cancer prevention and the acceptability of HPV vaccination among those with 14-18year old daughters: a quantitative-qualitative study". *BioMed Central Public Health* 12 (2012): 1034.
10. Ahmed SA., *et al.* "Knowledge, attitude and practice of Cervical Cancer screening among market women in Zaria, Nigeria". *Nigeria Medical Journal* 54.5 (2013): 316-319.
11. Bulbul S., *et al.* "Thoughts of mothers with children 0-15 years of age about Cervical Cancer and HPV vaccination in Anatolia". *Turkish Journal of Pediatrics* 2 (2013): 73-78.
12. Marlow LAV., *et al.* "Parental attitudes to pre-pubertal HPV vaccination". *Vaccine* 25.11 (2007): 1945-1952.
13. Ozan H., *et al.* "Detecting the knowledge level of patients who admitted to obstetrics and gynecology departments about Human Papilloma Virus and HPV vaccination". *Urology University Journal Faculty of Medicine* 37 (2011): 145-148.
14. Thompson SVLS., *et al.* "African American parents' attitudes towards HPV vaccination". *Ethnicity and Disease* 21.3 (2011): 335-341.

Volume 2 Issue 7 September 2018

© All rights are reserved by Sharmin Zaman Khan.