



## Prevalence and Determinants of Dyslipidaemia among Children and Adolescents in India: Findings from a Large Representative Survey

Abu Bashar\*

Department of Community Medicine, Institute of Medical Sciences, BHU, India

\*Corresponding Author: Abu Bashar, Department of Community Medicine, Institute of Medical Sciences, BHU, India.

Received: October 14, 2020

Published: November 30, 2020

© All rights are reserved by Abu Bashar.

### Introduction

It is estimated that 31% of all deaths globally i.e. 17.5 million deaths occurred due to cardiovascular disease in 2014 [1]. In India, mortality due to cardiovascular disease was expected to rise by 103% in men and by 90% in women from 1985 to 2015 [2].

Dyslipidaemia is defined as abnormal plasma lipid levels [3]. Dyslipidaemia is the disorder of lipoprotein metabolism that result in elevated levels of total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), triglycerides, or decreased high-density lipoprotein cholesterol (HDL-C) [4]. Dyslipidaemia is a primary, widely established as an independent major risk factor for coronary artery disease (CAD) and may even be a prerequisite for CAD, occurring before other major risk factors come into play [5]. It is, therefore, considered as an important modifiable risk factor for cardiovascular disease (CVD) and leading cause of morbidity and mortality worldwide [6].

There is a strong pathophysiological association of raised LDL cholesterol with initiation and progression of coronary atherosclerosis. Robust data are available that shows that lowering its levels can regress and stabilize atherosclerotic vascular disease. Similar, though not as robust, data are also available for raised triglycerides and low HDL cholesterol. Studies have reported that atherosclerosis begins silently during childhood [7]. Fatty streaks - the precursors of mature atherosclerotic lesions - are the first visible arterial lesions as reported in autopsy studies [8]. Pathogenesis of atherosclerosis is related to total cholesterol levels. Reduced serum high-density lipoprotein cholesterol level and increased serum low-density lipoprotein cholesterol level are independent risk factors for cardiovascular disease [9].

The link between childhood dyslipidaemia and occurrence of

atherosclerosis and its sequels in adulthood are well-documented [10-13]. Considering that the number of children at risk of developing cardiovascular disease is increasing, there has been no population-based study from India to determine the prevalence of dyslipidaemia. Few hospitals based and school-based studies with limited samples from India report the prevalence of dyslipidaemia among children and adolescents ranging from 10% - 70% [14-22].

With this background, this study aims to determine the prevalence of dyslipidaemia and its association with socio-demographic characteristics and clinical and laboratory findings in children and adolescent from India.

### Bibliography

1. The World Health Organization. Global status report on non-communicable diseases (2014).
2. Bulatao RA and Stephens PW. "Global estimates and projections of mortality by cause, 1970-2015". Washington: Population Health and Nutrition Department, World Bank (1992).
3. Maahs DM., *et al.* "Dyslipidemia in youth with diabetes: to treat or not to treat?" *The Journal of Pediatrics* 153 (2008): 458-465.
4. Haney EM., *et al.* "Screening for Lipid Disorders in Children and Adolescent". Rockville, MD: Agency for Healthcare Research and Quality (US) (2007).
5. Genest JG Jr. "Dyslipidemia and coronary artery disease". *Canadian Journal of Cardiology* 16 (2000): 3A-4A.
6. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents; National Heart, Lung, and Blood Institute. Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and adolescents: Summary report". *Pediatrics* 128.5 (2011): S213-S256.

7. Holman RL, *et al.* "The natural history of atherosclerosis. The early aortic lesions as seen in the middle of 20th century". *The American Journal of Pathology* 34 (1958): 209-235.
8. Sary HC. "Evolution and progression of atherosclerotic lesions in coronary arteries of children and young adults". *Arteriosclerosis* 9.1 (1980): 119-132.
9. Gotto AM Jr. "Low high-density lipoprotein cholesterol as a risk factor in coronary heart disease: a working group report". *Circulation* 103 (2001): 2213-2218.
10. Daniels SR. "Screening and treatment of dyslipidemias in children and adolescents". *Hormone Research in Paediatrics* 76.1 (2011): 47-51.
11. Psaty BM and Rivara FP. "Universal screening and drug treatment of dyslipidemia in children and adolescents". *The Journal of the American Medical Association* 307 (2012): 257-258.
12. Hatami M, *et al.* "Adolescent lipoprotein classifications according to National Health and Nutrition Examination Survey (NHANES) vs. National Cholesterol Education Program (NCEP) for predicting abnormal lipid levels in adulthood in a Middle East population". *Lipids in Health and Disease* (2012): 11-107.
13. Zachariah JP and De Ferranti SD. "NHLBI integrated pediatric guidelines: Battle for a future free of cardiovascular disease". *Future Cardiology* 9 (2013): 13-22.
14. Rajesh Dholarpuria, *et al.* "Atherosclerosis risk factors in adolescents". *Indian Journal of Pediatrics* (2007): 823-826.
15. Anita K, *et al.* "Lipid profile norms in Indian children". *Indian Pediatrics* 4 (1995): 1177-1180.
16. Ambady R, *et al.* "Insulin resistance and clustering of cardio-metabolic risk factors in urban teenagers in Southern India". *Diabetes Care* (2007): 1828-1833.
17. Parmar IB, *et al.* "Lipid profile in the progeny of parents with ischchemic heart disease". *Indian Journal of Pediatrics* 68 (2001): 617-621.
18. Anand NK, *et al.* "Lipid profile in children of 5-15 years age with parental history of IHD, Hypertension, Diabetes Mellitus". *Indian Pediatrics* 39 (2002): 168-172.
19. Khalil A, *et al.* "Insulin resistance and lipid profile in the children of young ischemic parents". *Indian Pediatr* 40 (2003): 556-560.
20. Ghulati S and Saxena A. "Study of lipid profile in children of parents with premature coronary artery disease". *Indian Pediatrics* 40 (2003): 556-560.
21. Agarwal A, *et al.* "Prevalence of dyslipidemia in students aged 6-16 years in a private school of rural Jaipur". *Indian Journal of Basic and Applied Medical Research* 6.1 (2016): 658-662.

#### Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

**Website:** [www.actascientific.com/](http://www.actascientific.com/)

**Submit Article:** [www.actascientific.com/submission.php](http://www.actascientific.com/submission.php)

**Email us:** [editor@actascientific.com](mailto:editor@actascientific.com)

**Contact us:** +91 9182824667