



The Relationship Between Neuromuscular Dysfunction of the Bladder and Pathological Narcissism Category: Mini Review

Melehin Aleksey Igorevich*

PhD, Associate Professor, Clinical Psychologist of the Highest Qualification Category, Consultant Somnologist, Cognitive Behavioral Therapist, Psychoanalyst of Psychosomatic Orientation (Paris School, IPSO), Psychoncologist, Body-Oriented Therapist, Titular Member of the Academy of Cognitive Therapy (USA), Full Member of the International Psycho-Oncology Society, USA

***Corresponding Author:** Melehin Aleksey Igorevich, PhD, Associate Professor, Clinical Psychologist of the Highest Qualification Category, Consultant Somnologist, Cognitive Behavioral Therapist, Psychoanalyst of Psychosomatic Orientation (Paris School, IPSO), Psychoncologist, Body-Oriented Therapist, Titular Member of the Academy of Cognitive Therapy (USA), Full Member of the International Psycho-Oncology Society, USA.

Received: January 04, 2024

Published: February 09, 2024

© All rights are reserved by **Melehin Aleksey Igorevich**.

Abstract

The results of our study indicate how important it is to pay attention to the characteristics of personality organization (pathological narcissism) in the clinical picture of CPPS. Pathological narcissism contributes to the development of somatic symptoms due to the patient's increased sensitivity to physical sensations, which, in turn, affects the perception of somatic distress. However, the results of our study force us to state that we are still far from controlling all the pathological aspects of CPPS, low tolerance for coping with stress, deterioration of mood, catastrophic beliefs, narcissistic vulnerability, which makes the experience of pain truly a source of suffering and requires, in addition to medical, psychotherapeutic tactics aimed at strengthening psychological stability in this group of patients.

Keywords: Urge; Overactive Bladder; Anxiety; Chronic Pelvic Pain Syndrome; Pain; Depression

Introduction

In neurology, neuromuscular dysfunction of the bladder, as well as hyperactive (stress-induced) bladder syndrome, refers to chronic pelvic pain syndrome (CPPS). This disorder in moderate to severe form is observed in 6.5-10% of men in the general population and should be understood as the result of complex interactions of psychological factors with somatic dysfunctions. It can be perceived as a somatoform disorder [1]. To this day, there is uncertainty about the etiology of this disorder. Research in the field of neurology has so far focused mainly on somatic factors. However, according to the phenotyped multimodal diagnostic system UP-OINT, the interaction of infectious, genitourinary, neurological, and psychological factors that are represented by the patient's mental state (depression, hypochondriacal disorder, anxiety spectrum disorders, somatoform disorder, PTSD) should be considered [2].

To this day, the psychological factors involved in CPPS in men have not been sufficiently studied in Russian practice, unlike our foreign colleagues [4]. To assess psychological factors in the UP-

OINT system, attention is often paid to symptoms of mental ill-being, but little is paid to the patient's personal organization. It has been shown that men with chronic prostatitis often (up to 63%) have borderline personality functioning, personality disorders in the narcissistic sphere predominate, narcissistic personality disorder (narcissism disorder) is present with difficulties in understanding both their own mental state and other people (the ability to mentalize [3-6].

Changes in the narcissistic configuration, pathological narcissism in a patient in neurology is an important and underestimated clinical problem associated with significant functional outcomes during treatment, and the risks of developing a refractory course.

According to the hierarchical model of A. Pincus and M. Lukovitsky, pathological narcissism refers to a deficit in the regulation of self-esteem, including maladaptive mechanisms for restoring and maintaining one's own "I" or self-image [6].

Characterized by a combination of three phenomena [7]

- Dysfunctional self-regulation (eg, tendency to focus on, intensify, and misinterpret bodily sensations);
- Failure in emotional regulation (formation of pain behavior, muscle-tonic manifestations);
- Difficulties in interpersonal relationships, which are often observed in men with CPPS

Also, this form of narcissism is represented by two types of dysfunctional phenotypes: grandiosity and vulnerability, which were adapted for patients with CPPS.

Patients with poorly defended narcissism have trouble regulating their behavior and emotions, especially in the context of negative emotions and social functioning. People with narcissistic vulnerability have been shown to react more intensely to pain (physical and psychological) and are more open in their communications about negative experiences than grandiose narcissists. The above-mentioned aspects can be summarized in terms that a person has a threatening part in the self. In addition, patients with poorly defended narcissism are concerned about their appearance and demonstrate increased somatic, illness-oriented preoccupation, which can be viewed from the point of view that they have a large part of the hypochondriacal view of the self. Poorly defended narcissistic patients show a more common profile of poor adaptation to rapidly changing situations in everyday life, reduced autonomy, self-efficacy, and a tendency to depression than those with well-defended narcissism [7]. In urological practice, if a patient has narcissistic vulnerability, there is an increased catastrophizing of pain, kinesophobia, and a spectrum of reinsurance and avoidance behavior. They have psychological (social) pain, which most often occurs when the patient is ostracized - excluded and ignored. Social pain activates the same brain structures as physical pain (such as the anterior cingulate cortex), meaning that experiencing social pain can be just as painful as experiencing physical pain.

Well-protected narcissism is more closely related to psychological well-being, whereas poorly protected narcissism correlates with a lack of self-confidence and with negative emotions such as depression and anxiety, problems with appearance, sensitivity to cause pain, lower self-esteem (inflict psychological wounds on oneself), reduce life satisfaction), as well as with an increase in somatic problems. The potential relationship between narcissistic grandiosity and vulnerability and severity of CPPS has not yet been studied. Although narcissistic grandiosity in men can serve as a defense against perceived weakness or infirmity, narcissistic vulnerability can exacerbate the physical experience of exhaustion and fragmentation [6]. Understanding such phenomena is important

for clinicians who often encounter patients with refractory CPPS. The present study was conducted as a preliminary study of the relationship between pathological narcissism and CPPS in a clinical sample of men.

The participants of the study

The main group consisted of 55 men, with an average age of 47.3 ± 9.5 years, who turned to a clinical psychologist on the recommendation of urologists, neurologists with a diagnosis of heading N41.1 according to the International Classification of Diseases of the 10th revision for psychological examination and psychotherapy. Non-inflammatory chronic pelvic pain syndrome (prostatodynia) and chronic abacterial prostatitis prevailed in patients. Patients complained of discomfort ("pulling", "burning"), pain in the perineum, lower abdomen or during ejaculation, changes in urination, "burning near the navel", lower back discomfort. The control group consisted of 55 men who had no urological manifestations, according to the results of an examination by a urologist, the average age was 41.0 ± 7.2 years.

Exclusion criteria

The presence of concomitant organic disease from the urinary system, gastrointestinal tract; severe concomitant pathology (for example, oncology, coronary heart disease, metabolic, autoimmune diseases); had autonomic dysfunction (for example, peripheral neuropathy, vagotomy); took medications that could change the functioning of the central nervous system (for example, anticholinergic, antiarrhythmic drugs, beta-blockers); underwent abdominal surgery, with the exception of appendectomy and/or cholecystectomy; have had or are currently experiencing serious mental disorders, substance abuse or alcohol abuse.

Research methods

The scale of pathological Narcissism (Pathological Narcissism Inventory, PNI), which includes 52 questions. The assessment of grandiosity includes exploitative interpersonal behavior, immersion in grandiose fantasies and presumptuous altruistic behavior. Vulnerability consists of admiring self-esteem, hiding perceived flaws and needs, devaluing others, and having anger at limitations.

Somatization was assessed using a health assessment scale (Patient Health questionnaire, PHQ-15), which allows you to assess which of the 15 symptoms, including gastrointestinal ones, the patient has been concerned about in the last 3 months.

The presence and degree of symptoms of anxiety and depression were assessed using the Patient Health Questionnaire-9 (PHQ-9) and Generalized Anxiety Disorder-7 (GAD-7) scales.

Pain Catastrophizing Scale.

The results of the study

Men with CPPS have pronounced symptoms of generalized anxiety disorder and depression (feeling of emptiness), which indicates the presence of a state of threat to the “I” (the threatening part in the “I”). The tendency to somatization of affect as a protective mechanism and a way of avoiding internal distorted standards.

On the part of pathological narcissism, men with CPPS experience fluctuations between assertive grandiosity and vulnerability, and there may be bouts of insecurity that destroy a sense of grandiosity or singularity (uniqueness).

More pronounced changes are observed on the part of narcissistic fragility (vulnerability) than grandiosity, which reflects the vulnerability of the “I” (hypersensitivity), a tendency to anxiety arousal, disorders in emotional self-regulation (fluctuations in self-esteem, feelings of shame). Following H. Kogut, it can be said that these patients have diffuse narcissistic vulnerability. The difficulty for these patients lies in the pain and suffering that accompany the presence of such disparate “split off” or non-integrated parts of the Self, which leads to the defensive use of maladaptive intrapersonal and interpersonal tactics to maintain a “stable” sense of self.

The presence of narcissistic fragility in patients with CPPS suggests that they carry traumatic narcissism (narcissistic trauma), which poses a risk of developing PTSD, since this injury is perceived as a threat to the “I”.

There are high rates of self-sacrifice and self-aggrandizement, as the deliberate use of altruistic acts to maintain an “inflated” image of “Me” (for example, “I work”, “I eat”, “I try”), as well as the presence of grandiose fantasies (to fight, “win”, “achieve”, “to prove yourself”, which perform the function of artificial mobilization. In order to cope with unstable self-esteem, patients with CPPS can create an exaggerated sense of superiority, uniqueness, plunging into grandiose fantasies.

On the part of narcissistic fragility as a form of maladaptive or poorly protected narcissism, there is a pronounced demanding rage, which reflects a tendency to put pressure on oneself (self-sabotage), to be dissatisfied with oneself, to react with anger (anger, resentment, irritation) in cases where expectations from a significant other are not satisfied or do not meet internal strict standards (“norms”). Patients can drown out this demanding rage with alcohol.

Unstable self-esteem prevails in the form of fluctuations in the sphere of self-esteem, recognition of its failures in the absence of

external sources of recognition, “reassurance”. support, admiration (feedback from another, for example, close people, doctors). Devaluation manifests itself in the absence of spontaneous, flexible interest in other people who do not demonstrate the necessary admiration, desired behavior. There is a sense of shame for the excessive need for recognition from those who once disappointed, often a father figure. The presence of narcissistic fragility can be understood as a pathological defense against negative emotions and reflects their reduced resilience.

Patients with CPPS seem to lack appropriate strategies to deal with strong emotions, and they rely only on protective mechanisms (e.g. somatization, denial) to regulate their emotions and self-esteem. They feel ashamed of their needs and ambitions.

Significant positive associations have been found between somatic urological symptoms and symptoms of depression, anxiety, catastrophization of pain, as well as between somatic urological symptoms and narcissistic grandiosity and vulnerability in patients with CPPS.

Narcissistic fragility in patients with CPPS is associated with the presence and severity of symptoms of depression, which is consistent with evidence that this form of pathological narcissism is associated with a wide range of symptoms of disorders of both the affective and psychotic spectrum. Anxiety in poorly protected narcissistic individuals (with fragility) to be separated from others and their desire to be recognized can lead to an excessive search for conformity, which can lead to somatoform symptoms (abdominal pain, gastrointestinal specific anxiety, nausea, itching), and they can be associated with a sense of meaninglessness, which refers to psychological insecurity or negative emotionality.

Note that the presence of self-sacrifice in patients with CPPS/self-elevation, unstable self-esteem and demanding rage affect all indicators of psychological well-being, as well as the catastrophization of pain.

The relationship between grandiosity and somatic symptoms in men is less consistent with theoretical assumptions. According to early research, men with grandiose features were more likely to approve of physical integrity and even superiority. However, the men in our study voluntarily agreed to psychological examination and psychotherapy. Perhaps for men with grandiose facial features, the expression of physical complaints is preferable to the experience of crumpled self-perception. Thus, physical pain can perform a protective function, saving men from further erosion of self-representation. It may also conform to socialized gender norms that

emphasize men's physical attractiveness and discourage men from expressing emotional distress.

It has been shown that the tendency to somatize affect in the form of painful manifestations is influenced by pathological narcissism, the presence and severity of anxiety disorder, depression and pain catastrophizing.

The results of our study indicate how important it is to pay attention to the characteristics of personality organization (pathological narcissism) in the clinical picture of CPPS. Pathological narcissism contributes to the development of somatic symptoms due to the patient's increased sensitivity to physical sensations, which, in turn, affects the perception of somatic distress. However, the results of our study force us to state that we are still far from controlling all the pathological aspects of CPPS, low tolerance for coping with stress, deterioration of mood, catastrophic beliefs, narcissistic vulnerability, which makes the experience of pain truly a source of suffering and requires, in addition to medical, psychotherapeutic tactics aimed at strengthening psychological stability in this group of patients.

Bibliography

1. Ehlert U., *et al.* "Chronic pelvic pain as a somatoform disorder". *Psychotherapy and Psychosomatics* 68 (1999): 87-94.
2. Fischer-Kern M., *et al.* "Psychische Struktur bei chronischen Schmerzpatienten". *Zeitschrift für Psychosomatische Medizin und Psychotherapie* 56 (2010): 34-46.
3. Giannantoni A., *et al.* "Resilience in the face of pelvic pain: A pilot study in males and females affected by urologic chronic pelvic pain". *Neurourology and Urodynamics* 40.4 (2020): 1011-1020.
4. Kealy D., *et al.* "Investigating the Link Between Pathological Narcissism and Somatization". *The Journal of Nervous and Mental Disease* 206.12 (2018): 964-967.
5. Kealy D., *et al.* "Pathological narcissism and somatic symptoms among men and women attending an outpatient mental health clinic". *International Journal of Psychiatry in Clinical Practice* 20.3 (2016): 175-178.
6. Keltikangas-Järvinen L., *et al.* "Personality Pathology Underlying Chronic Prostatitis". *Psychotherapy and Psychosomatics* 37 (1982): 87-95.
7. Schalkwijk F., *et al.* "Narcissistic Personality Disorder: Are Psychodynamic Theories and the Alternative DSM-5 Model for Personality Disorders Finally Going to Meet?". *Frontiers in Psychology* 12 (2021): 676733.