



Rethinking Postoperative Pain After Nonsurgical Root Canal Treatment: Understanding, Prevention, and Management-Review

Mothanna Al Rahabi*

Associate Professor, College of Dentistry, Taibah University, Madinah Al Munawwarah, Saudi Arabia

***Corresponding Author:** Mothanna Al Rahabi, Associate Professor, College of Dentistry, Taibah University, Madinah Al Munawwarah, Saudi Arabia.

Received: July 12, 2018; **Published:** August 10, 2018

Abstract

Background: Postoperative pain after root canal treatment is frustrating to practitioners and patients. The incidence of postoperative pain could be changed with recent advances in endodontic techniques and equipment. The aim of this review was to highlight key aspects of postoperative pain after nonsurgical root canal treatment, including causes, related factors, effects of recent advances, and management of postoperative pain.

Methods: Electronic databases were searched for publications on postoperative pain after nonsurgical root canal treatment. The search parameters were postoperative pain, nonsurgical treatment, single visit, recent advances in endodontics, and management of postoperative pain of endodontic treatment.

Results: The results of this review revealed that the incidence of postoperative pain after nonsurgical root canal treatment ranged from 3% to 58%. The main reason for postoperative pain was microorganisms, and there was no significant difference in postoperative pain between single-visit root canal treatment and multiple-visit treatment. The use of recent endodontic techniques and devices will reduce postoperative pain. To control and manage postoperative pain after root canal treatment, a so-called flexible plan for drug administration can be used based on the severity of pain.

Conclusions: Postoperative pain after root canal treatment ranged from mild to moderate, and it can happen even after high-standard root canal treatment. The management of postoperative pain with a correct approach is often considered a sign of clinical excellence.

Keywords: Root Canal Treatment; Postoperative Pain; Review; Acetaminophen; Analgesics; Ibuprofen; Narcotic Medications

Introduction

Pain can be considered one of the most important elements in dentistry because fear of pain is a major reason for dental apprehension [1]. Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage or described in terms of such damage [2]. There is a common belief that root canal treatment is the most painful dental treatment [3-5]. Although root canal treatment alleviates pain, postoperative pain after root canal treatment unfortunately is still a common problem [6]. The incidence of this problem ranges from 3% to 58% [7-11]. The variance among these results can be explained by the use of different criteria in postoperative pain assessment, the use of different materials and techniques, and the failure to consider preoperative pain as a variable [12]. The severity of postoperative pain can range from mild to severe [13]. In one study, 12% of patients reported severe pain after finishing treatment using a visual analogue scale (VAS) [14]. The period of postoperative pain can last from one day to several weeks and can be a major cause of patient dissatisfaction [13]. The purpose of this review is to examine postoperative pain in nonsurgical root canal treatment, including related factors and management of postoperative pain.

Causative and related factors of pain after nonsurgical root canal treatment

The main reason for postoperative pain after nonsurgical root canal treatment is microorganisms; pain generally results from acute inflammation of peri-radicular tissues. Mechanical or chemical damage to peri-radicular tissues can be considered among other causes [15]. Mechanical factors, such as over instrumentation or extrusion of root canal obturation materials, have been related to postoperative pain after endodontic treatment [6,16]. Postoperative pain following nonsurgical root canal treatment is multifactorial and is related to several factors such as presence of periapical pathosis, missed canals, inadequate cleaning and shaping, apical extrusion of debris, apical patency during instrumentation, extrusion of irrigant and intracanal medications, overbite restorations [17], the type of tooth [18,19], and gender [5]. There is strong evidence of a correlation between preoperative and postoperative pain, and patients who have acute preoperative pain are likely to have more severe postoperative pain [19-22]. There is controversy regarding the correlation between pulpal status and postoperative pain, with some reports mentioning that pulpal status has an impact on postoperative pain [12,20,23], and others failing to dem-

onstrate an influence of pulpal status [18,21,24]. Several reports identified no relationship between age and postoperative pain after nonsurgical root canal treatment [23,25,26]. Various studies have concluded that women had more postoperative pain after nonsurgical root canal treatment than did men [9,12,14,21,22,27,28]. The biological differences between men and women that result from changes in serotonin and non-adrenalin hormones may be the cause of these differences [29,30]. Cortisol hormone organises the feeling of pain. This hormone is secreted more in males than in females [31,32]. Generally, men have been used as subjects in most clinical research, and women have been excluded [33]. Oestrogen and women’s menstrual cycle effects, as well as societal expectations, may contribute to differences between the genders in their physiological reaction to pain [34,35]. The incidence of postoperative pain after nonsurgical root canal treatment was higher in the mandibular arch and in molar teeth [19,28]. This difference may result from the dense trabecular pattern structure of the mandible, which decreases blood circulation and concentrates infection, delaying healing [21]. This difference results from the complex anatomy of the lower molars [5,36].

Number of visits and postoperative pain

Studies have shown that most clinicians complete root canal treatment in multiple visits [20,26,37]. However, in the last few decades, over 70% of schools in all geographic areas and various studies have advocated single-visit root canal treatment [9,38-40]. Single-visit root canal treatment is an efficient procedure, but the incidence of postoperative pain is still a source of concern [41]. Several studies mentioned found no significant difference in postoperative pain between single-visit and multiple-visit root canal treatment [9,13,20,27,38,42-46], whereas some researchers reported that postoperative pain was less in single-visit root canal treatment [47]. Table 1 summarises the relationship between number of visits and postoperative pain.

Recent advances in endodontics and postoperative pain after root canal treatment

In recent years, the practice of endodontics has benefitted from several new technologies that can improve its efficiency, safety, and quality [48]. The use of recent endodontic techniques and devices, such as operative microscopes, electronic apex locators, and rotary nickel titanium systems, will shorten the treatment time and increase the success rate of endodontic treatment [49]. Studies have shown that crown-down instrumentation that allows the practitioner to reach the apical terminus gradually, starting from the coronal aspect of the root canal, significantly reduces apically extruded debris, which is considered one of the important etiological factors of postoperative pain and flare-ups [50]. In addition, nickel–titanium engine-driven instrumentation extrudes less debris than do stain-

less steel K-files manipulated by hand, and use of this instrumentation will decrease postoperative pain [51-53]. Irrigant activation could be an effective method for minimising postoperative pain [54]. The use of a negative apical pressure irrigation device can reduce postoperative pain significantly [6].

Prevention and management of postendodontic treatment pain

Prevention and management of postoperative pain after nonsurgical root canal treatment is considered a complementary part of root canal treatment. Providing patients with information regarding expected postoperative pain, and managing pain by prescribing medications will improve patient confidence in their dentists, increase patients’ pain threshold, and improve their attitude regarding dental treatment in the future [55,56]. The synthesis of prostaglandin is prevented by nonsteroidal anti-inflammatory drugs (NSAIDs) through reducing the activity of the enzymes cyclooxygenase (Cox) 1 and 2. Several NSAIDs such as ibuprofen, aspirin [57], flurbiprofen [58], ketorolac [59,60], and etodolac [61] have been used to manage postoperative pain after root canal treatment. The best way to control moderate to severe pain is a combination of two or more drugs, so that a lower dose of each drug is used with fewer side effects [60]. The combination of an NSAID and acetaminophen can provide additional analgesia for management of dental pain. If this combination does not control pain, a narcotic analgesic will give additional relief [62,63]. To control and manage postoperative pain after root canal treatment, a so-called flexible plan for drug administration can be used based on the severity of pain [64]. Drugs are administered four times per day until pain subsides (Figure 1).




Figure 1: A flexible analgesic plan

Study and Reference No.	Type of Tooth	Pulp Status	No. of Cases	Percentage of Cases		Incidence of Postoperative Pain
				Single	Multiple	
Wong Amy Wai-Yee., et al. 2015 [13]	All types	Vital and nonvital	538 teeth	51%	49%	No significant difference was found in postoperative pain between single and multiple visits for root canal treatment.
Onay Emel O., et al. 2015 [47]	All types	Vital and nonvital	1819 teeth	31%	69%	The incidence of flare-up was minimal when teeth were treated in single visit. A peri-apical lesion in necrotic teeth is a significant risk factor for flare-ups.
Singh, Smita, and Aniket Garg. 2012 [45]	Single-rooted teeth	Vital and nonvital	200 teeth	50%	50%	No significant difference was found in postoperative pain between single and multiple visits for root canal treatment.
El Mubarak AH., et al. 2010 [20]	All types	Vital and nonvital	230	50%	50%	No significant difference was shown in postoperative pain between single and multiple visits for endodontic treatment.
Wang C., et al. 2010 [46]	Permanent anterior teeth	Vital	100 teeth	50%	50%	In teeth with vital pulp and a single canal, there was no significant difference in incidence and intensity of postoperative pain after one- or two-visit endodontic treatment.
Al-Negrish., et al. (2006) [27]	Central incisor teeth	Nonvital	120 teeth	50%	50%	There were no significant differences in the incidence and severity of postoperative pain between one- and two-visit root canal therapy.
Eleazer Paul D and Kristen R Eleazer (1998) [38]	Molar teeth	Nonvital	402 teeth	50%	50%	This study showed an advantage for one-visit treatment at a 95% confidence level.
Albashaireh ZSM and AS Alnegrish 1998 [9]	All types	Vital and nonvital	300 teeth	50%	50%	Pain was significantly higher in the multiple-visit group.

Table 1: Summary of several studies regarding the relationship between postoperative pain after root canal treatment and number of visits.

Conclusions

Postoperative pain after root canal treatment ranges from mild to moderate pain, and it can occur even after high-standard root canal treatment. New advances in endodontic treatment reduce postoperative pain. Proper management of postoperative pain is often considered a sign of clinical excellence; it requires a thorough understanding of the physiology of pain and the mechanisms by which drugs and therapies offer relief.

Bibliography

- Freeman R. "The role of memory on the dentally anxious patient's response to dental treatment". *Irish Journal of Psychological Medicine* 8.2 (1991): 110-115.
- Rosenberg PA. "Diagnosis, in Endodontic pain diagnosis, causes, prevention and treatment, P.A. Rosenberg, Editor". *Springer: New York* (2014) 1.
- Pak JG and SN White. "Pain prevalence and severity before, during, and after root canal treatment: a systematic review". *Journal of endodontics* 37.4 (2011): 429-438.
- Görduysus MÖ and M Görduysus. "Endodontic patient profile of Hacettepe University, Faculty of Dentistry in Ankara, Turkey". *International Dental Journal* 50.5 (2000): 274-278.
- Watkins CA., et al. "Anticipated and experienced pain associated with endodontic therapy". *The Journal of the American Dental Association* 133.1 (2002): 45-54.
- Gondim E., et al. "Postoperative pain after the application of two different irrigation devices in a prospective randomized clinical trial". *Journal of Endodontics* 36.8 (2010): 1295-1301.

7. Sathorn C., et al. "The prevalence of postoperative pain and flare-up in single-and multiple-visit endodontic treatment: a systematic review". *International Endodontic Journal* 41.2 (2008): 91-99.
8. Torabinejad M., et al. "Effectiveness of various medications on postoperative pain following complete instrumentation". *Journal of Endodontics* 20.7 (1994): 345-354.
9. Albashaireh Z and A Alnegrish. "Postobturation pain after single-and multiple-visit endodontic therapy. A prospective study". *Journal of Dentistry* 26.3 (1998): 227-232.
10. Rogers MJ., et al. "Comparison of effect of intracanal use of ketorolac tromethamine and dexamethasone with oral ibuprofen on post treatment endodontic pain". *Journal of Endodontics* 25.5 (1999): 381-384.
11. DiRenzo A., et al. "Postoperative pain after 1- and 2-visit root canal therapy". *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology* 93.5 (2002): 605-610.
12. Gotler M., et al. "Postoperative pain after root canal treatment: a prospective cohort study". *International Journal of Dentistry* (2012).
13. Wong AW., et al. "Incidence of post-obturation pain after single-visit versus multiple-visit non-surgical endodontic treatments". *BMC Oral Health* 15.1 (2015): 96.
14. Ng YL., et al. "Prevalence of and factors affecting post-obturation pain in patients undergoing root canal treatment". *International Endodontic Journal* 37.6 (2004): 381-391.
15. Siqueira JF., et al. "Incidence of postoperative pain after intracanal procedures based on an antimicrobial strategy". *Journal of Endodontics* 28.6 (2002): 457-460.
16. Harrison JW., et al. "Incidence of pain associated with clinical factors during and after root canal therapy. Part 2. Postobturation pain". *Journal of Endodontics* 9.10 (1983): 434-438.
17. de Oliveira Alves V. "Endodontic flare-ups: a prospective study". *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology* 110.5 (2010): e68-e72.
18. Ince B., et al. "Incidence of postoperative pain after single-and multi-visit endodontic treatment in teeth with vital and non-vital pulp". *European Journal of Dentistry* 3.4 (2009): 273-279.
19. Arias A., et al. "Predictive models of pain following root canal treatment: a prospective clinical study". *International Endodontic Journal* 46.8 (2013): 784-793.
20. El Mubarak AHH., et al. "Postoperative pain in multiple-visit and single-visit root canal treatment". *Journal of Endodontics* 36.1 (2010): 36-39.
21. Ali SG., et al. "Prevalence of and factors affecting post-obturation pain following single visit root canal treatment in Indian population: A prospective, randomized clinical trial". *Contemporary Clinical Dentistry* 3.4 (2012): 459-463.
22. Sadaf D and MZ Ahmad. "Factors associated with postoperative pain in endodontic therapy". *International journal of biomedical science: IJBS* 10.4 (2014): 243-247.
23. Segura-Egea J., et al. "Pain associated with root canal treatment". *International Endodontic Journal* 42.7 (2009): 614-620.
24. Pasqualini D., et al. "Postoperative pain after manual and mechanical glide path: a randomized clinical trial". *Journal of Endodontics* 38.1 (2012): 32-36.
25. Martín-González J., et al. "Influence of root canal instrumentation and obturation techniques on intra-operative pain during endodontic therapy". *Medicina Oral, Patología Oral Y Cirugía Bucal* 17.5 (2012): e912.
26. Edionwe JO., et al. "Single visit root canal treatment: a prospective study". *Nigerian Journal of Clinical Practice* 17.3 (2014): 276-281.
27. Al-Negrish ARS. and R Hababbeh. "Flare up rate related to root canal treatment of asymptomatic pulpally necrotic central incisor teeth in patients attending a military hospital". *Journal of Dentistry* 34.9 (2006): 635-640.
28. Alí A., et al. "Influence of preoperative pain intensity on postoperative pain after root canal treatment: A prospective clinical study". *Journal of Dentistry* 45 (2015): 39-42.
29. Marcus DA. "Interrelationships of neurochemicals, estrogen, and recurring headache". *Pain* 62.2 (1995): 129-139.
30. Dao T., et al. "Modulation of myofascial pain by the reproductive hormones: a preliminary report". *The Journal of Prosthetic Dentistry* 79.6 (1998): 663-670.
31. Walton R and A Fouad. "Endodontic interappointment flare-ups: a prospective study of incidence and related factors". *Journal of Endodontics* 18.4 (1992): 172-177.
32. Mehrvarzfar P., et al. "Effect of supraperiosteal injection of dexamethasone on postoperative pain". *Australian Endodontic Journal* 34.1 (2008): 25-29.
33. Greenspan JD., et al. "Studying sex and gender differences in pain and analgesia: a consensus report". *Pain* 132 (2007): S26-S45.

34. Paller CJ, *et al.* "Sex-based differences in pain perception and treatment". *Pain Medicine* 10.2 (2009): 289-299.
35. Genet J, *et al.* "Preoperative and operative factors associated with pain after the first endodontic visit". *International Endodontic Journal* 20.2 (1987): 53-64.
36. Cleghorn BM, *et al.* "Root and root canal morphology of the human permanent maxillary first molar: a literature review". *Journal of Endodontics* 32.9 (2006): 813-821.
37. Ahmed M, *et al.* "Root canal treatment in general practice in Sudan". *International Endodontic Journal* 33.4 (2000): 316-319.
38. Eleazer PD and KR Eleazer. "Flare-up rate in pulpally necrotic molars in one-visit versus two-visit endodontic treatment". *Journal of Endodontics* 24.9 (1998): 614-616.
39. Qualtrough A, *et al.* "Preclinical endodontology: an international comparison". *International Endodontic Journal* 32.5 (1999): 406-414.
40. Oginni AO and CI Udoye. "Endodontic flare-ups: comparison of incidence between single and multiple visit procedures in patients attending a Nigerian teaching hospital". *BMC Oral Health* 4.1 (2004): 4.
41. Mohammadi Z, *et al.* "One-visit versus multiple-visit endodontic therapy-a review". *International Dental Journal* 56.5 (2006): 289-293.
42. Su Y, *et al.* "Healing rate and post-obturation pain of single-versus multiple-visit endodontic treatment for infected root canals: a systematic review". *Journal of Endodontics* 37.2 (2011): 125-132.
43. Law A, *et al.* "Predicting severe pain after root canal therapy in the national dental PBRN". *Journal of Dental Research* (2014).
44. Raju T, *et al.* "Evaluation of pain in single and multi rooted teeth treated in single visit endodontic therapy". *Journal of International Oral Health: JIOH* 6.1 (2014): 27.
45. Singh S and A Garg. "Incidence of post-operative pain after single visit and multiple visit root canal treatment: a randomized controlled trial". *Journal of Conservative Dentistry* 15.4 (2012): 323-327.
46. Wang C, *et al.* "Comparison of post-obturation pain experience following one-visit and two-visit root canal treatment on teeth with vital pulps: a randomized controlled trial". *International Endodontic Journal* 43.8 (2010): 692-697.
47. Onay EO, *et al.* "The evaluation of endodontic flare-ups and their relationship to various risk factors". *BMC oral Health* 15.1 (2015): 142.
48. Alrahabi M and MS Ahmad. "Knowledge regarding technical aspects of non-surgical root canal treatment in Al-Madinah Al-Munawarah private dental centers". *Saudi Endodontic Journal* 5.3 (2015): 155.
49. Fleming CH, *et al.* "Comparison of classic endodontic techniques versus contemporary techniques on endodontic treatment success". *Journal of Endodontics* 36.3 (2010): 414-418.
50. Zarrabi MH, *et al.* "An in vitro comparative study of apically extruded debris resulting from conventional and three rotary (Profile, Race, FlexMaster) instrumentation techniques". *Journal of Oral Science* 48.2 (2006): 85-88.
51. Vaudt J, *et al.* "Ex vivo study on root canal instrumentation of two rotary nickel-titanium systems in comparison to stainless steel hand instruments". *International Endodontic Journal* 42.1 (2009): 22-33.
52. Madhusudhana K, *et al.* "Apical extrusion of debris and irrigants using hand and three rotary instrumentation systems-An in vitro study". *Contemporary Clinical Dentistry* 1.4 (2010): 234.
53. Nekoofar MH, *et al.* "Comparison of the effect of root canal preparation by using WaveOne and ProTaper on postoperative pain: a randomized clinical trial". *Journal of Endodontics* 41.5 (2015): 575-578.
54. Ramamoorthi S, *et al.* "Comparative evaluation of postoperative pain after using endodontic needle and EndoActivator during root canal irrigation: A randomised controlled trial". *Australian Endodontic Journal* 41.2 (2015): 78-87.
55. Van Wijk A and J Hoogstraten. "Reducing fear of pain associated with endodontic therapy". *International Endodontic Journal* 39.5 (2006): 384-388.
56. Van Wijk AJ, *et al.* "The effect of written information on pain experience during periodontal probing". *Journal of Clinical Periodontology* 31.4 (2004): 282-285.
57. Holstein A, *et al.* "Evaluation of NSAIDs for treating post-endodontic pain". *Endodontic Topics* 3.1 (2002): 3-13.
58. Doroschak AM, *et al.* "Evaluation of the combination of flurbiprofen and tramadol for management of endodontic pain". *Journal of Endodontics* 25.10 (1999): 660-663.

59. Penniston SG and KM Hargreaves. "Evaluation of periapical injection of ketorolac for management of endodontic pain". *Journal of Endodontics* 22.2 (1996): 55-59.
60. Mehrvarzfar P, *et al.* "Effects of three oral analgesics on post-operative pain following root canal preparation: a controlled clinical trial". *International Endodontic Journal* 45.1 (2012): 76-82.
61. Comfort M, *et al.* "A study of the comparative efficacy of three common analgesics in the control of pain after third molar surgery under local anaesthesia". *Australian Dental Journal* 47.4 (2002): 327-330.
62. Breivik EK, *et al.* "Combining diclofenac with acetaminophen or acetaminophen-codeine after oral surgery: a randomized, double-blind single-dose study". *Clinical Pharmacology and Therapeutics* 66.6 (1999): 625-635.
63. Wideman GL, *et al.* "Analgesic efficacy of a combination of hydrocodone with ibuprofen in postoperative pain". *Clinical Pharmacology and Therapeutics* 65.1 (1999): 66-76.
64. Thomas R. "How pain is controlled in endodontic therapy". *SAAD Digest* 31 (2015): 32-37.

Volume 2 Issue 9 September 2018

© All rights are reserved by Mothanna Al Rahabi.