



## Plasma-Rich-Fibrin Role in Dentistry

Ermira Biu<sup>1</sup>, Lordian Nunci<sup>2</sup>, Silvana Beraj<sup>1\*</sup>, Vjollca Shpata<sup>5</sup>, Bledar Lufi<sup>1</sup>, Gentian Vyshka<sup>4</sup> and Tatjana Cina<sup>3</sup>

<sup>1</sup>Department of Para clinical Sciences, Faculty of Technical Medical Sciences, UMT, Lahore, Pakistan

<sup>2</sup>Central Anesthesiology and Intensive Care Service of UHCT "Mother Theresa", Tirana, Albania

<sup>3</sup>Professor, Dean of the Faculty of Technical Medical Sciences, National Blood Transfusion Centre, Tirana, Albania

<sup>4</sup>Faculty of Medicine, UMT, Lahore, Pakistan

<sup>5</sup>Faculty of Technical Medical Sciences, UMT, Lahore, Pakistan

\*Corresponding Author: Silvana Beraj, Department of Para clinical Sciences, Faculty of Technical Medical Sciences, UMT, Lahore, Pakistan.

Received: March 09, 2018; Published: September 19, 2018

### Abstract

Platelet rich fibrin (PRF) is an improved version of Platelet Rich Plasma (PRP), a fibrin matrix which consists of growth factors and cytokine, can serve as a resorbable membrane facilitating wound healing. It is a second-generation platelet concentrate which is prepared from the patient's own blood free of any anticoagulant. This article enriches the benefits and role of plasma-rich fibrin in dentistry.

**Keywords:** Platelet Growth Factor; Maxillofacial Surgery; Graft Material; Periodontal Lesions

### Introduction

Platelets regenerative abilities were first described in 1974 [1] where the growth factors play an important role to wound healing. In a study of Gassling, *et al.* PRF membranes resulted to be better to collagen membranes (Bio-Gide) in proliferation and suitable for serving as a scaffold for periosteal tissue engineering [2]. Platelets play an important role not only in the regeneration of the wound, thanks to fibrin matrix which works as healing material, but also in hemostasis [3]. These growth factors stimulate the stem cells to produce new host tissue very quickly and this is why platelet rich fibrin is so effective in the post-treatment healing process and plays a key role in tissue regeneration.

In transfusion medicine, platelet concentrates are used to prevent serious bleeding due to severe thrombocytopenia or its prophylactic usage for patients undergoing coronary bypass grafting in cardiovascular services [4].

PRF (Platelet Rich Fibrin) was first used in 2001 by Choukroun, *et al.* [12] in oral and maxillofacial surgery and is currently

considered as a new generation of platelet concentrate. It's a natural fibrin matrix from autologous blood and shows high cicatricial capacity. Its membranes favor and support the healing process due to slow polymerization mode [8].

PRF is the new generation of PRP (or platelet rich plasma) which creates a fibrin matrix or scaffolds for new tissue construction rich in platelet-derived growth factors, but also many other vascular proteins and enzymes part of this PRF mesh such as mesenchymal stem cells, VEGF (vascular endothelial growth factors), leukocytes, TGF  $\beta$ 1 (transforming growth factor  $\beta$ 1) etc [5,12]. PRF accelerate the healing process and also optimizes bone grafting results [6,7].

### The PRF Procedure

The PRF procedure begins drawing the patient's own peripheral blood. Then the blood is placed in a centrifuge for 10 minutes (low speed centrifugation system) without anticoagulant in a 10 ml vacutainer. During the centrifuge process, the blood coagulates and is separated into three distinct layers: the upper straw-colored

acellular plasma, middle portion containing the fibrin clot, and red-colored lower portion containing red blood cells. Only the middle portion is collected, which is the PRF. The upper layer acellular plasma and the bottom layer of RBC) are removed and discarded. While, the middle layer is the fibrin mesh which contains platelets [8,9].

The mechanism involved here is the fibrinogen concentrated in the upper part of the tube combines with the circulating thrombin when centrifuged to form fibrin. So, fibrin clot stays between the upper layer of acellular plasma and the lower one containing red blood cells. The time between blood collection to its transfer for centrifuging should be short, in order to avoid its coagulation before the process, especially when it is taken without any anticoagulant [10] Once separated from the clot, the PRF may be withdrawer and we can put this layer into a plug, or use it as a membrane depending on how and in what treatment do we need it [11].

### PRF Role in Dentistry

PRF is basically used in various disciplines of dentistry to repair various lesions and regenerate dental and oral tissue. In recent times a lot of research has been done on PRF and numerous cases have been reported regarding the use of PRF clot and membranes: in oral surgery for bone augmentation, sinus lifts etc. and in periodontics to correct gingival recession, guided bone regeneration, periapical lesions etc. It has also been used for regeneration in open apex, regenerative pulpotomies, periapical surgeries etc [13].

PRF can be used in maxillofacial surgery as a filling material in extraction sockets acting as a stable blood clot for neovascularization and accelerated tissue regeneration. This can be used to improve wound healing in immunocompromised and diabetic patients. PRF stimulates coagulation (with thrombospondin) and wound closure, it can be used as an adjuvant in patients on anticoagulant therapy [14].

According to a study of Simonpieri., et al. PRF can be used successfully as the sole filling material during sinus lift and implantation and seems to be a reliable surgical option promoting natural bone regeneration [26].

PRF maybe used also in combination with beta tricalcium phosphate without bone graft in chronic periodontal lesions and sinus lift procedures [27].

A combination of PRF with bone substitutes and other adjuncts may be necessary in residual defects where one or several walls are missing or damaged in order to provide an adequate reconstruction of bone volume. Since PRF is an optimized blood clot, the fibrin increases the bond between the graft materials acting as a 'sticker' between them and serves as a physiological linkage between the wound tissues thanks to its healing properties [14]. The fibrin mesh promotes cellular proliferation, migration creating biologically links between tissues, remodeling their structure.

The PRF fibrin membrane, thanks to its elasticity and strength, can be used as a membrane for guided bone regeneration (GBR). Its matrix architecture covers and stabilizes bone graft material and operative site in general [13]. Simonpieri., et al. [25] suggested using a mix of PRF with a bone graft, placing in bone defects, or, in cases of immediate implants, covering it with several PRF layers, noting good clinical results.

PRF membranes acts as a "biological joining material" between grafts and surgical site acting as a matrix which leads to the process of angiogenesis and migration of osteoprogenitor cells to the graft [28].

While in periodontics, PRF has been used to treat gingival recession, intra-bony defects and periapical lesions. Some studies show the use of PRF gel and PRF membrane in combination with a bone graft for treating a tooth with a combined periodontics-endodontic lesion [15]. To cover the defect are used two layers of PRF membrane because are very thin, inhomogeneous and leucocytes and platelet aggregates are believed to be concentrated in end of the membrane. So, it can be used two layers of membrane in opposite sense in order to prevent the resorption of the membrane and to allow the entire surgical area to be exposed to leucocytes and platelet aggregates [15]. The rapid healing and regenerative properties of growth factors of PRF membrane can be used successfully in periodontal therapy, in gingival recession as a coverage in exposed root surfaces in mandibular process with a flap that covers this membrane over the exposed surface [16].

PRF can stimulate the osteogenic differentiation of the human dental pulp cells by up regulating osteoprotegerin and alkaline phosphatase expression promoting the healing of bone processes defects. PRF also releases growth factors such as platelet-derived growth factor and transforming growth factor which promote periodontal regeneration. These facts emphasize the role of PRF in reparative dentin formation [17].

Studies have shown that PRF can be used on periapical inflammatory lesion in cases of pulp tissue necrosis and can be used as a membrane for the treatment of this lesion inducing dental regeneration and as scaffolds accelerating bone repair [15]. Use of PRF in regenerative pulpotomy procedures can be done where coronal pulp is removed and the pulp wound is covered by PRF followed by sealing it with MTA and GIC [18]. Also, some case reports show that the PRF membrane usage as a matrix shows to speed up the process of periapical healing lesions. Use of a membrane can prevent the extrusion of material [18]. According to Keswani, *et al*, PRF can also be used in cases of big periapical lesions after surgeries to repair bone defects by filling them, as matrix in tissue ingrowth guiding proliferation, migration of cells and as scaffolds in order to enhance the microvascularization of necrotic pulp [19].

#### Advantages and disadvantages of its uses

Some studies have demonstrated that PRF is a healing biomaterial with a great potential for bone and soft tissue regeneration, without inflammatory reactions and may be used alone or in combination with bone grafts, promoting hemostasis, bone growth, and maturation [20]. This autologous matrix demonstrated in the *in vitro* studies a great potential to increase cell attachment [7]. Simonpieri, *et al.* reviewed advantages of the use of PRF as it acts as a stabilizing sheath and offers mechanical sustenance, such as the regeneration through PRF membranes both the bone volume and gingival tissue. They also reported satisfactory clinical results related to reshaping the whole alveolar bone and the restoration of gingival volume and peri-implant bone, achieving adequate mechanical and aesthetic properties [25].

Comparing to PRP it has several advantages over it: it's a minimized autologous blood manipulation [29,30] including easier preparation. This entire process is natural, without any external manipulation leading to the absence of any immunological reaction [8,29]. So, it's not required a chemical manipulation of the blood, which makes it strictly an autologous leukocyte-platelet-rich fibrin matrix [8,22]. which acts as a biodegradable scaffold [23] that

favors the development of microvascularization and is able to guide epithelial cell proliferation and migration to its surface [8].

It has a natural fibrin mesh with growth factors within that may keep their activity for a relatively longer period and stimulate tissue regeneration [7]. Used as a membrane, it avoids a donor site surgical procedure and results in a reduction in patient discomfort during the early wound-healing period [32]. It is an economical and quick option compared with recombinant growth factors when used in conjunction with bone grafts [31] It can be used in combination with bone grafts or as one layer, depending on the manipulation [21,25]. Comparing to PRP, it's more efficient and shows better clinical results [33].

While talking about disadvantages: The success of the PRF protocol depends on blood collection time and its way on to the centrifuge [8]. The final amount available is low because it is an autologous blood [30] and for the process is needed a glass-coated tube to achieve clot polymerization [34]. It exists possible refusal of treatment by the puncture required for blood collection (Wani 2014).

#### Conclusion

PRF is the new generation of platelet concentrates and has potential applications in medicine and dentistry too. It can be used alone or in combination with grafts or other biomaterials. Although the mechanisms how growth factors work is not clearly understood yet, PRF shows to have beneficial outcomes and satisfactory clinical results giving new perspectives of treatment in dentistry field.

#### Bibliography

1. Ross R, *et al.* "A platelet-dependent serum factor that stimulates the proliferation of arterial smooth muscle cells *in vitro*". *Proceedings of the National Academy of Sciences of the United States of America* 71 (1974): 1207-1210.
2. Gassling VL, *et al.* "Platelet-rich fibrin membranes as scaffolds for periosteal tissue engineering". *Clinical Oral Implants Research* 21 (2010): 543-549.
3. Gassling VL, *et al.* "Platelet-rich plasma and platelet-rich fibrin in human cell culture". *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology* 108 (2009): 48-55.
4. David M, *et al.* "Classification of platelet concentrates: from pure platelet rich plasma(P-PRP) to leucocyte and platelet rich fibrin(L-PRF)". *Trends Biotechnology* 27.3 (2008): 158-167.

5. Eren G., *et al.* "Effect of centrifugation time on growth factor and MMP release of an experimental platelet-rich fibrin-type product". *Platelets* 27.5 (2016): 427-432.
6. Kawase T., *et al.* "The heat-compression technique for the conversion of platelet-rich fibrin preparation to a barrier membrane Platelet-rich fibrin". *International Journal of Clinical and Experimental Medicine* 8 (2015): 7922-7929.
7. Wu CL., *et al.* "Platelet-rich fibrin increases cell attachment, proliferation and collagen-related protein expression of human osteoblasts". *Australian Dental Journal* 57 (2012): 207-212.
8. Dohan DM., *et al.* "Platelet-rich fibrin (PRF): a second-generation platelet concentrates. Part I: technological concepts and evolution". *Oral Surgery, Oral Medicine, Oral Pathology, and Oral Radiology* 101.3 (2006): e37-44.
9. Kumar KR., *et al.* "Role of plasma-rich fibrin in oral surgery". *Journal of Pharmacy and Bioallied Sciences* 8 (2016): S36-S38.
10. Naik B., *et al.* "Role of Platelet rich fibrin in wound healing: A critical review". *Journal of Conservative Dentistry* 16.4 (2013): 284-293.
11. Mazor Z., *et al.* "Sinus floor augmentation with simultaneous implant placement using Choukroun's platelet-rich fibrin as the sole grafting material: A radiologic and histologic study at 6 months". *Journal of Periodontology* 80 (2009): 2056-2064.
12. Choukroun J., *et al.* "Une opportunit  en paro-implantologie: le PRF". *Implantodontie* 42 (2000): 55-62.
13. Borie E., *et al.* "Fuentes R Platelet-rich fibrin application in dentistry: a literature review". *International Journal of Clinical and Experimental Medicine* 8.5 (2015): 7922-7929.
14. Corso MD., *et al.* "Use of autologous leukocyte and platelet rich fibrin (L-PRF) membrane in post avulsion sites: an overview of Choukroun's PRF". *The journal of implant and advanced clinical dentistry* 1.9 (2010): 27-35.
15. Shivashankar VY., *et al.* "Combination of platelet rich fibrin, hydroxyapatite and PRF membrane in the management of large inflammatory periapical lesion". *Journal of Conservative Dentistry* 16.3 (2013): 261-264
16. Anilkumar K., *et al.* "Platelet-rich-fibrin: A novel root coverage approach". *Journal of Indian Society of Periodontology* 13.1 (2009): 50-54.
17. Huang FM., *et al.* "Platelet-rich fibrin increases proliferation and differentiation of human dental pulp cells". *Journal of Endodontics* 36.10 (2010): 1628-1632.
18. Geeta IB., *et al.* "Natural Meliorate: Revolutionary Tissue Engineering in Endodontics". *Journal of Clinical and Diagnostic Research* 7.11 (2013): 2644-2646.
19. Keswani D and Pandey RK. "Revascularization of an immature tooth with a necrotic pulp using platelet-rich fibrin: a case report". *International Endodontic Journal* 46 (2013): 1096-1104.
20. Kim TH., *et al.* "Comparison of platelet-rich plasma (PRP), platelet-rich fibrin (PRF), and concentrated growth factor (CGF) in rabbit-skull defect healing". *Archives of Oral Biology* 59 (2014): 550-558.
21. Choukroun J., *et al.* "Platelet-rich fibrin (PRF): a second-generation platelet concentrates. Part V: histologic evaluations of PRF effects on bone allograft maturation in sinus lift". *Oral Surgery, Oral Medicine, Oral Pathology, and Oral Radiology* 101 (2006): 299-303.
22. Gupta V., *et al.* "Regenerative potential of platelet rich fibrin in dentistry: literature review". *Asian Journal of Oral Health and Allied Sciences* 1 (2011): 22-28.
23. Li Q., *et al.* "Platelet-rich fibrin promotes periodontal regeneration and enhances alveolar bone augmentation". *BioMed Research International* (2013): 638043.
24. Cortese A., *et al.* "Platelet-rich fibrin (PRF) in implant dentistry in combination with new bone regenerative technique in elderly patients". *International Journal of Surgery Case Reports* 28 (2016): 52-56.
25. Simonpieri A., *et al.* "Current knowledge and perspectives for the use of platelet-rich plasma (PRP) and platelet-rich fibrin (PRF) in oral and maxillofacial surgery part 2: Bone graft, implant and reconstructive surgery". *Current Pharmaceutical Biotechnology* 13 (2012): 1231-1256.
26. Simonpieri A., *et al.* "Simultaneous sinus-lift and implantation using microthreaded implants and leukocyte- and platelet-rich fibrin as sole grafting material: a six-year experience". *Implant Dentistry* 20.1 (2011): 2-12.
27. Jayalakshmi KB., *et al.* "Platelet-Rich Fibrin with  $\beta$ -Tricalcium Phosphate—A Novel Approach for Bone Augmentation in Chronic Periapical Lesion: A Case Report". (2012).
28. Simonpieri A., *et al.* "The relevance of Choukroun's platelet-rich fibrin and metronidazole during complex maxillary rehabilitations using bone allograft. Part II: Implant surgery, prosthodontics, and survival". *Implant Dentistry* 18 (2009): 220-229.

29. Kang YH., *et al.* "Platelet-rich fibrin is a Bioscaffold and reservoir of growth factors for tissue regeneration". *Tissue Engineering Part A* 17 (2011): 349-359.
30. Choukroun J., *et al.* "Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part IV: clinical effects on tissue healing". *Oral Surgery, Oral Medicine, Oral Pathology, and Oral Radiology* 101 (2006): e56-60.
31. Girish Rao S., *et al.* "Bone regeneration in extraction sockets with autologous platelet rich fibrin gel". *Journal of Oral and Maxillofacial Surgery* 12 (2013): 11-16.
32. Jankovic S., *et al.* "Use of platelet-rich fibrin membrane following treatment of gingival recession: a randomized clinical trial". *The International Journal of Periodontics and Restorative Dentistry* 32 (2012): e41-50.
33. Saluja H., *et al.* "Platelet-Rich fibrin: A second generation platelet concentrate and a new friend of oral and maxillofacial surgeons". *Annals Maxillofacial Surgery* 1 (2011): 53-57.
34. Dohan DM., *et al.* "Cytotoxicity analyses of Choukroun's platelet-rich fibrin (PRF) on a wide range of human cells: The answer to a commercial controversy". *Oral Surgery, Oral Medicine, Oral Pathology, and Oral Radiology* 103 (2007): 587-583.

**Volume 2 Issue 10 October 2018**

**© All rights are reserved by Silvana Beraj., et al.**