



## Evaluation of Vertical Root Fractures Using Cone-Beam Computed Tomography: A Mini Review

**Helen Rushdi Ayoubi\***

Assistant Professor, Department of Endodontics and Operative Dentistry, Faculty of Dental Medicine, Damascus University, Damascus, Syrian Arab Republic

**\*Corresponding Author:** Helen Rushdi Ayoubi, Assistant Professor, Department of Endodontics and Operative Dentistry, Faculty of Dental Medicine, Damascus University, Damascus, Syrian Arab Republic.

**DOI:** 10.31080/ASDS.2022.06.1457

**Received:** August 18, 2022

**Published:** August 26, 2022

© All rights are reserved by **Helen Rushdi Ayoubi**.

### Abstract

There are inadequate data or reviews to confirm the reliability and efficacy of cone-beam computed tomography (CBCT) in evaluation of vertical root fractures (VRFs). Thus, the aim of this mini review is to discover the reliability and efficacy of CBCT in the evaluation of VRFs and prove CBCT as a picturing addition in the different aspects of VRFs.

**Keywords:** Diagnosis; Vertical Root Fractures (VRFs); Cone-Beam Computed Tomography (CBCT); Teeth Fractures

### Introduction

The inflammatory practicability can occur due to the vertical root fracture (VRF) which is the strictest form of longitudinal tooth fracture [1]. Therefore, it can lead to the resorption of bone and formation of granulation tissue [2]. The radiographic signs, clinical signs and symptoms of vertical root fractures (VRFs) can be different, indefinite, and take a similar appearance to periodontal lesions and periapical lesions of dental origin [3-5].

For the time being, a good diagnosis and evaluation of longitudinal root fractures can be obtained through the patient's dental history, by asking him about the symptoms, the nature and aspect of the pain, the presence of swelling, as well as the formation of deep and shallow periodontal pockets, in order to reach an accurate diagnosis [6].

The presence of different types of longitudinal root fractures makes the radiographic evaluation process somewhat difficult, in addition to its lack of clarity on the radiographs in many situations [7]. One of the disadvantages of traditional radiographs is that it can show only one third of the VRFs [8-10]. Using periapical displacement radiography (PDR), it is possible to discover the VRFs

at the level of the X-ray beam [8,11] but in such situations, the anatomical details can appear on the radiographs indistinctly [12,13]. It is known that the traditional radiography shows radiographs in only two-dimensions (2D), those are the length and width without showing depth that needed for illustrating the anatomical structures and may be obtained using three-dimensions (3D) radiographs [14]. Several three-dimensional radiography methods and devices such as conventional computed tomography (CT), cone-beam computed tomography (CBCT), and multi-detector computed tomography have been developed to overcome the limitations of using two-dimensional intraoral radiographs. Recently, a lot of research has been published to confirm the reliability and effectiveness of these devices. The results of many studies were in favor of CT devices, as they provided very great credibility and effectiveness in evaluating root fractures [3-10,15-18]. In any case, there are many disadvantages related to CT, including: the large amount of radiation exposure, the factitious product, and the space dissolution [16]. As a result, companies have produced CBCT devices to eliminate these disadvantages, and to be the perfect solution for diagnosing many complex cases [3].

Many researchers have explained the usefulness and importance of CBCT in the diagnosis and arrangement in identified as-


pects of dentoalveolar trauma, especially root fractures [19-22], luxation and/or displacement, and alveolar fracture [20].

**Previous studies**


Many *in vitro* studies have been conducted regarding the use of CBCT, and it has shown its credibility and great effectiveness for discovering cases similar to the VRFs [3,23]. There are special uses of CBCT for the evaluation of root fractures. Hassan., *et al.* [3] showed in their *in vitro* study to discover the VRFs, that there is an effect of root canal filling on fracture clarity, by evaluating radiographs taken using CBCT, and periapical radiography (PR). They concluded that the radiographs taken using CBCT are more reliable (0.86) than the PR (0.66) for discovering the VRFs. However, these images are less accurate in the root canals filled with opaque material. Moudi., *et al.* [24] concluded that the accuracy of CBCT does not reduce in the presence of root canals filled with gutta-percha, and therefore, CBCT has a great reliability. However, the presence of prefabricated posts may have a slight impact on the reliability of these devices, but without statistically significant differences (Figures 1,2). There is a contrast between the researches [25], especially with regard to the method of creating the root fracture, which can lead to the formation of many broken pieces, and this in turn can be discovered clinically within (0.2 mm-0.4 mm), but this is still controversial [3,10]. As a solution to that, root fractures can be formed by applying a large compressive load to the posts within roots [12,13]. Several machines have been used in these researches to test the development and extension of these fractures. The overall reliability of the CBCT for discovering two fracture patterns was 0.87 for the non-filled root canals, and 0.45 for the filled root canals. The evaluative reliability of the PR was 0.63 for the non-filled root canals, and 0.53 for the filled root canals [12,13].

Discoverability also reveals to be CBCT scanner-specific [26,27]. The observation of discoverer, as well as the parameters used and specified in each device, can affect the reliability of the CBCT [25,28,29].


The hardened beam resulting from the root canal filling reduces the reliability and effectiveness of complete or incomplete fractures discovery in the filled root canals (Figures 3-5) [5,30-32]. Finally, there are many errors in the evaluation of root fractures [33], which may result in unsuitable treatments of these cases [34]. Disarrange from radiopaque materials may be misinterpreted as fracture lines [3,34].




**Figure 1:** An axial CBCT picture shows the molar and premolar with VRF [24].




**Figure 2:** Cross sectional pictures of molar and premolar with VRF. (A). A molar with VRF. (B). A premolar with prefabricated post and gutta-percha has VRF. (C). A premolar with gutta-percha shows VRF [24].



**Figure 3:** CBCT, axial view; (Orange arrow): VRF in gutta-percha filled root canals [32].



**Figure 4:** CBCT, axial view; (Orange lines): VRF in non-filled root canals (without gutta-percha) [32].



**Figure 5:** Effect of hardened beam on producing artifacts similar to VRF in gutta-percha filled root canals (Orange lines) [32].

When using CBCT, the presence of the hardened beam reduces the evaluative result in the discovery of root fractures when the metallic posts are present [35]. In contrast, the fiber posts do not seem to interpose with the reliability and efficacy of CBCT for discovery of VRF [31,36].

On the evaluation of ninety-five root fractures, Wang, *et al.* [37] showed that the CBCT has greater reliability than RP. Nevertheless, The disarranging of rays due to the presence of radiopaque root canal filling material reduces the reliability of CBCT [37]. Long, *et al.* [38] suggested that the CBCT has a great evaluative result for discovery of tooth fractures, with a reported reliability and efficacy of 0.92 and 0.85, respectively.

Chavda, *et al.* [39] compared several radiographs taken using PR or CBCT of atraumatically extracted teeth to confirm the presence or absence of root fractures. They found a reliability of 0.16 for PR, and 0.27 for CBCT, and a great efficacy of 0.92 for PR and 0.83 for CBCT.

A systematic review confirmed the existence of contrast between the available results, as well as the bias of many researchers to the inefficacy of CBCT in discovery VRFs in filled root canals [40]. This is also consistent with systematic reviews in other researches [41,42].

### Conclusion

There is inadequate proof to support the use of CBCT for discovery of VRF. Nevertheless, CBCT can be used when longitudinal root fractures are suspected with the patient's absence or fuzzy of signs and symptoms. So, CBCT can show that there is a periradicular bone resorption, and this is evidence of a VRF in the accompanying root.

### Bibliography

1. Tamse A., *et al.* "Radiographic features of vertically fractured, endodontically treated maxillary premolars". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 88 (1999): 348-352.
2. Tamse A. "Vertical root fractures in endodontically treated teeth: diagnostic signs and clinical management". *Endodontic Topics* 13 (2006): 84-94.
3. Hassan B., *et al.* "Detection of vertical root fractures in endodontically treated teeth by a cone beam computed tomography scan". *Journal of Endodontics* 35 (2009): 719-722.
4. Mora MA., *et al.* "In vitro assessment of local computed tomography for the detection of longitudinal tooth fractures".

- Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 103 (2007): 825-829.
5. Khedmat S., et al. "Evaluation of three imaging techniques for the detection of vertical root fractures in the absence and presence of gutta-percha root fillings". *International Endodontic Journal* 45 (2012): 1004-1009.
  6. Edlund M., et al. "Detection of vertical root fractures by using cone-beam computed tomography: a clinical study". *Journal of Endodontics* 37 (2011): 768-772.
  7. Melo SL., et al. "Diagnostic ability of a cone-beam computed tomography scan to assess longitudinal root fractures in prosthetically treated teeth". *Journal of Endodontics* 36 (2010): 1879-1882.
  8. Rud J and Omnell KA. "Root fractures due to corrosion. Diagnostic aspects". *Scandinavian Journal of Dental Research* 78 (1907): 397-403.
  9. Youssefzadeh S., et al. "Dental vertical root fractures: value of CT in detection". *Radiology* 210 (1999): 545-549.
  10. Kamburoğlu K., et al. "Detection of vertical root fracture using cone-beam computerized tomography: an in vitro assessment". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 109 (2010): e74-81.
  11. Meister F., et al. "Diagnosis and possible causes of vertical root fracture". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 49 (1980): 243-253.
  12. Patel S., et al. "The detection of vertical root fractures in root filled teeth with periapical radiographs and CBCT scans". *International Endodontic Journal* 46 (2013): 1140-1152.
  13. Brady E., et al. "A comparison of cone beam computed tomography and periapical radiography for the detection of vertical root fractures in nonendodontically treated teeth". *International Endodontic Journal* 47 (2014): 735-746.
  14. Scarfe WC., et al. "Use of cone beam computed tomography in endodontics". *International Journal of Dentistry* 2009 (2009): 634567.
  15. Kamburoğlu K., et al. "Effectiveness of limited cone-beam computed tomography in the detection of horizontal root fracture". *Dental Traumatology* 25 (2009): 256-261.
  16. Bernardes RA., et al. "Use of cone-beam volumetric tomography in the diagnosis of root fractures". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 108 (2009): 270-277.
  17. Bornstein MM., et al. "Comparison of intraoral radiography and limited cone beam computed tomography for the assessment of root-fractured permanent teeth". *Dental Traumatology* 25 (2009): 571-577.
  18. Ozer S. "Detection of vertical root fractures of different thicknesses in endodontically enlarged teeth by cone beam computed tomography versus digital radiography". *Journal of Endodontics* 36 (2010): 1245-1249.
  19. Tyndall DA and Rathore S. "Cone-beam CT diagnostic applications: caries, periodontal bone assessment, and endodontic applications". *Dental Clinics of North America* 52 (2008): 825-841.
  20. Cohenca N., et al. "Clinical indications for digital imaging in dento-alveolar trauma-part 1: traumatic injuries". *Dental Traumatology* 23 (2007): 95-104.
  21. Ilg'uy D., et al. "Detection of jaw and root fractures using cone beam computed tomography: a case report". *Dentomaxillofacial Radiology* 38 (2009): 169-173.
  22. Patel S. "New dimensions in endodontic imaging-part 2: cone beam computed tomography". *International Endodontic Journal* 42 (2009): 463-475.
  23. Tsesis I., et al. "Comparison of digital with conventional radiography in detection of vertical root fractures in endodontically treated maxillary premolars: an *ex vivo* study". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 106 (2008): 124-128.
  24. Moudi E., et al. "Assessment of vertical root fracture using cone-beam computed tomography". *Imaging Science in Dentistry* 44 (2014): 37-41.

25. Talwar S, et al. "Role of cone-beam computed tomography in diagnosis of vertical root fractures: a systematic review and meta-analysis". *Journal of Endodontics* 42 (2016): 12-24.
26. Elsaltani M, et al. "Detection of simulated vertical root fractures: which Cone-beam Computed Tomographic System Is the Most Accurate?" *Journal of Endodontics* 42 (2016): 972-977.
27. Tiepo M, et al. "Evaluation of root fracture in endodontically treated Teeth using Cone Beam Computed Tomography". *The Journal of Contemporary Dental Practice* 18 (2017): 94-99.
28. Bechara B, et al. "Number of basis images effect on detection of root fractures in endodontically treated teeth using a cone beam computed tomography machine: an *in vitro* study". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 115 (2013a): 676-681.
29. Bezerra IS, et al. "Influence of the artifact reduction algorithm of Picasso Trio CBCT system on the diagnosis of vertical root fractures in teeth with metal posts". *Dentomaxillofacial Radiology* 44 (2015): 20140428.
30. Schulze R, et al. "Artifacts in CBCT: a review". *Dentomaxillofacial Radiology* 40 (2011): 265-273.
31. Neves F, et al. "Oblique or orthoradial CBCT slices for preoperative implant planning: which one is more accurate?" *Brazilian Journal of Oral Sciences* 13 (2014): 104-108.
32. Hekmatian E, et al. "Detection of vertical root fractures using cone-beam computed tomography in the presence and absence of gutta-percha". *The Scientific World Journal* 2018 (2018): 1920946.
33. Bechara B, et al. "Comparison of cone beam CT scans with enhanced photostimulated phosphor plate images in the detection of root fracture of endodontically treated teeth". *Dentomaxillofacial Radiology* 42 (2013b): 20120404.
34. Kajan Z and Taromsari M. "Value of cone beam CT in detection of dental root fractures". *Dentomaxillofacial Radiology* 41 (2012): 3-10.
35. Costa FF, et al. "Detection of horizontal root fracture with small-volume cone-beam computed tomography in the presence and absence of intracanal metallic post". *Journal of Endodontics* 37 (2011): 1456-1459.
36. Pinto M, et al. "Influence of exposure parameters on the detection of simulated root fractures in the presence of various intracanal materials". *International Endodontic Journal* 50 (2017): 586-594.
37. Wang P, et al. "Detection of dental root fractures by using cone-beam computed tomography". *Dentomaxillofacial Radiology* 40 (2011): 290-298.
38. Long H, et al. "Diagnostic accuracy of CBCT for tooth fractures: a meta-analysis". *Journal of Dentistry* 42 (2014): 240-248.
39. Chavda R, et al. "Comparing the *in vivo* diagnostic accuracy of digital periapical radiography with cone-beam computed tomography for the detection of vertical root fracture". *Journal of Endodontics* 40 (2014): 1524-1529.
40. Chang E, et al. "Cone-beam Computed Tomography for Detecting Vertical Root Fractures in Endodontically Treated Teeth: a Systematic Review". *Journal of Endodontics* 42 (2016): 177-185.
41. Corbella S, et al. "Cone beam computed tomography for the diagnosis of vertical root fractures: a systematic review of the literature and meta-analysis". *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* 118 (2014): 593-602.
42. Rosen E, et al. "The Diagnostic efficacy of cone-beam computed tomography in endodontics: a Systematic review and analysis by a hierarchical model of efficacy". *Journal of Endodontics* 41 (2015): 1008-1014.

**Volume 6 Issue 9 September 2022**  
 © All rights are reserved by Helen Rushdi Ayoubi.