



Holistic Approach to Obesity Management without Strenuous Physical Activity

Jain Richa*, Sharma S and Jagmohan Duggal

Institute of Medical Education and Research, India

*Corresponding Author: Jain Richa, Institute of Medical Education and Research, India.

Received: August 01, 2019; Published: August 19, 2019

DOI: 10.31080/ASGIS.2019.02.0073

Abstract

Obesity is epidemic and the underlying cause of various lifestyle related disorders that are the scourge of mankind today. Using integrated medicine approach that combines best practices in Ayurveda and Allopathy to an method that has yielded amazing results of weight reduction coupled with health restoration. The approach involves powerful effects of Oral Food Chelation therapy and has been used in the treatment of 100 patients. What emerged as the primary preventive and curative therapy Diet control along with body mind healing.

Keywords: Obesity; Epidemic; Lifestyle

This method promotes balance and integrates body- mind-soul to provide phenomenal results right from achieving ideal weight with natural ease and maintaining it, to healing emotional pain underlying weight issues. All this and more with absolutely no side effects, supplements and reduced dependency on medicines.

Worldwide, more than 2 billion adults have been reported overweight 700 million are obese and the numbers are rapidly increasing. More than 3 billion deaths have been linked to underline causes of obesity. Unhealthy food habits, sedentary lifestyle, consumption of carbohydrate/fat-rich foods (fast food) is worsening the risk of ischemic heart disease, diabetes, etc. which is turning into an epidemic in developing countries. Incidents of obesity with increasing prevalence in women and children has doubled in the last ten years. Hormonal changes in middle years have been reported to be linked to weight gain. The problem of obesity is significantly linked with over and irregular eating patterns which can be prevented. It causes a lot financial burden on the government health care agencies Obesity has reached epidemic proportions. India is reportedly, as per recent studies, facing an epidemic of obesity with more than 135 million cases reported in a yearly basis. According to ICMR-INDIAB study 2015, prevalence rate of obesity and central obesity are varies from 11.8% to 31.3% and 16.9%-36.3% respectively. In India, the propensity of obesity varies with age, gender, geographical environment and socio-economic status with abdominal obesity being the major risk factor for myocardial infarction. Among women, obesity is significantly higher as compared to men due to genetic and lifestyle reasons.

Methodology

A pilot study of 100 patients divided into five groups was undertaken ranging from 15- 55years.

The research methodology was adopted to treat the patients by assessing the body constitution(prakriti) an ancient Ayurvedic method that helps to establish the root cause of the case. Once the body constitution is established the imbalance of the doshas (kapha pitha vata) is determined using Nadi Shodan. After this, we use dosha specific balancing herbs, traditional condiments and food preparation methods are provided to the patients. The patient is put on a diet therapy and use of natural herbs coupled with step by step instructions on food preparation so that nutritional value of the food ingredients is present.

A calm disposition stemming out relaxation techniques is used to enhance the absorption power of the digestive system by systemic relaxation using deep breathing from diaphragm. A focused treatment at a cellular level including a mind-body connect that is combination of lifestyle changes, detoxification through panchkarma cleansing.

Research findings

The studies reveals that the BMI which was showing a standard deviation of 5.52 was observed from a mean of 31.6 before the study and after the study mean values of 28.3 were observed. This means a percentage change of 27.3% reduction was obtained which is highly significant.

In group 1 (aged 15-25years) weight loss of 6.4kgs were obtained. In group 2(aged 26-35years) weight loss of 10.9kgs, group 3, 9.74kgs were obtained.

Discussion

We developed a new innovative practical approach for the assessment of the nutritional status of the patients. This combine with an improved understanding of the biochemical and metabolic process has led to focus treatment at the cellular level. This method introduces a mind body connect that helps by changing dietary patterns. This results in an emergence of new behavioral structure leading to management of chronic obesity. The results were evaluated using statistical methods and are presented in Annexure 1.

The results exhibit that an average weight loss of 9.3kgs in 78days was achieved in group 1. In the other age group of 46-55 years, they were able to shed 7.72kgs in the same time and maintain this weight loss. After the weight loss the patients were given suggestions for healthy eating to maintain this weight loss and they were monitored for 2 years.

The patients were able to overcome the compulsive food cravings that were the root cause for obesity using psychological diet counseling. Obesity is linked to maldigestion therefore the goal of the method was improving the digestive power by stimulating the mediator of the parasympathetic nervous system that travels from the brain to neatly all of the thoracic and abdominal organs and triggers a cascade it calming effects [1-8].

Conclusion

The research highlights the of Svelte that, 'Let food be your medicine lest medicine becomes your food. Billions have been spent on obesity prevention with no real cure in sight. Modern life-style that eulogizes a fast life coupled with fast food leads to obesity. Our unique methodology aims at lifestyle change that focuses on the root cause of the disease and tries to eliminate in a holistic manner. Another advantage of the method is the economic aspect as it involves no financial burden but uses traditional material already found in the kitchen. Further obesity is linked to a host of other lifestyle related chronic disorders that are intractable further research is required in this direction to see if this approach can be used to tackle them.

Initial		Final		t	df	Sig. (2-tailed)
Mean	SD	Mean	SD			
31.6	5.68	28.3	5.52	16.7	89	.000**

Table 1: Test between initial and final BMI.

** Significant at 0.01 level (t=2.58)

BMI	Initial	Final	% change BMI
Normal	1(1.1)	28(31.1)	27.3
OW	44(48.9)	37(41.1)	-0.2
OG1	26(28.9)	17(18.9)	-0.3
OG2	10(11.1)	5(5.6)	-0.5
OG3	9(10.0)	3(3.3)	-0.7
Total	90(100.0)	90(100.0)	

Table 2: Distribution of BMI.

AGE	Weight_Lose	Duration_M	Duration_W	Duration_D
15-25	6.4	1.9	7.9	55.5
26-35	10.9	3.1	13.1	91.5
36-45	9.74	2.4	10.2	71.4
46-55	7.72	2.6	11.2	78.1
>55	8.1	2.2	9.4	65.7
Total	9.3	2.6	11.2	78.4

Table 3: Distribution of average of weight loss according to age.

Final_BMI	Weight_Lose	Duration_M	Duration_W	Duration_D
Normal	11.4	3.0	12.7	88.9
OW	7.9	2.5	10.7	75.0
OG1	8.1	2.4	10.4	72.5
OG2	10.0	2.4	10.3	72.0
OG3	11.9	2.2	9.3	65.0
Total	9.3	2.6	11.2	78.4

Table 4: Distribution of average of weight loss according to Final BMI.

Group Statistics					
	Gender	N	Mean	Std. Deviation	Std. Error Mean
BMI_Diff	Male	14	-3.5136	1.78214	.47630
	Female	76	-3.2829	1.91861	.22008
Weight_Lose	Male	14	10.8318	5.34152	1.42758
	Female	76	8.9842	5.50033	.63093

Table 5: t test between male and female for BMI and weight loss.

	t-test for Equality of Means					95% Confidence Interval of the Difference	
	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	Lower	Upper
BMI_Diff	-.418	88	.677	-.23068	.55232	-1.32830	.86694
Weight_Lose	1.160	88	.249	1.84758	1.59296	-1.31810	5.01325

Table 6: t-test for weight loss.

Wt loses		t	df	Sig. (2-tailed)
Mean	SD			
9.3	5.48	16.0	89	.000**

Table 7: ** Significant at 0.01 level (t=2.58)

Acknowledement

I thank Prof Pawandeep Singh for contribution in manuscript preparation and editing

Bibliography

1. GE Mullin., *et al.* "Integrative Weight Management: A Guide for Clinicians". *Nutrition and Health*.
2. Sharron Dalton. "Integrative Weight management". (2014).
3. Dausch J. "Determining when obesity is a disease". *Journal of the American Dietetic Association* 101 (2001): 293.
4. World Health Organization. Obesity and overweight (2006).
5. Newburgh LH and Johnston MW. "Endogenous obesity—a misconception". *JAMA* 3 (1930): 815-825.
6. NIH, NHLBI. "Clinical guidelines for the identification, evaluation, and treatment of overweight and obesity in adults—the evidence report". *Obesity Research* 6 (1998): S51-209.
7. CMS. Decision memo for intensive behavioral therapy for obesity (CAG-00423N) (2011). <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx>
8. World Health Organization (2012).

Volume 2 Issue 7 September 2019

© All rights are reserved by Jain Richa., *et al.*