

## ACTA SCIENTIFIC GASTROINTESTINAL DISORDERS (ISSN: 2582-1091)

Volume 8 Issue 12 December 2025

Editorial

# Flood Syndrome

## Zain Majid1\* and Hassaan Yousuf2

<sup>1</sup>Department of Gastroenterology, Sindh Institute of Urology and Transplantation, SIUT, Pakistan <sup>2</sup>Sindh Medical College, Pakistan

\*Corresponding Author: Zain Majid, Department of Gastroenterology, Sindh Institute of Urology and Transplantation, SIUT, Pakistan.

Received: November 17, 2025

Published: November 20, 2025

© All rights are reserved by

Zain Majid and Hassaan Yousuf.

### **Abstract**

Flood Syndrome is a condition commonly seen in patients with decompensated cirrhosis; however, it currently lacks definitive treatment options. We through this editorial aim to raise awareness about this interesting yet serious condition.

Keywords: Chronic Liver Disease (CLD); Flood Syndrome

#### Introduction

Decompensated cirrhosis can have various presentations, with ascites being the most common manifestation. The presence of ascites is associated with an overall poor prognosis [1].

Umbilical hernias are observed in approximately 20% of patients with chronic liver disease (CLD) and ascites [2]. The spontaneous rupture of an umbilical hernia in a patient with CLD and ascites is known as Flood syndrome [2]. This syndrome was first described by Dr. Frank Flood in 1961 [3]. The name refers to the sudden gushing of ascitic fluid that occurs following rupture of the umbilical hernia in a CLD patient [4].

The main mechanism leading to Flood syndrome involves an abnormal increase in intra-abdominal pressure secondary to ascites, along with dilated periumbilical blood vessels that cause fascial defects in the abdominal wall [2]. This may potentially lead to infecti-

on and can be life-threatening if not managed promptly; therefore, surgical intervention is often required [2].

Currently, no definitive medical treatment exists for the management of this condition, and therapy primarily revolves around management of ascites and surgical intervention, that is repair of the hernia [5,6].

However, there are certain limitations to surgery, including patients having acute liver failure, acute viral hepatitis, coagulopathy, low platelet count and acute renal failure [7].

Early recognition and management of Flood syndrome are crucial, especially in patients awaiting liver transplantation. These patients should be examined on every opd visits to their general practitioners or hepatologists and counselled regarding their diet, advised to avoid lifting heavy weights and prevent constipation from occuring [6].

### **Bibliography**

- Perez-Lopez KP and Reyes-Zermeño MG. "Impact on survival of decompensated liver cirrhosis and large volume paracentesis: a retrospective cohort". Annals of Hepatology 30 (2025): 101804.
- 2. Jayaprakash A., *et al.* "Understanding the Spectrum of Flood Syndrome: A Case Series". *Cureus* 16.6 (2024): e62059.
- 3. Flood FB. "Spontaneous perforation of the umbilicus in Laennec's cirrhosis with massive ascites". *The New England Journal of Medicine* 264 (1961): 72-74.
- 4. DeLuca IJ and Grossman ME. "Flood syndrome". *JAAD Case Report* 1.1 (2014): 5-6.
- 5. Kumaraman S Srivastava. "A continuous leak from within: a case report on flood syndrome". *Journal of Surgical Case Reports* 2 (2022): rjac011.
- 6. Kingsnorth A and LeBlanc K. "Hernias: inguinal and incisional". *Lancet* 362 (2003): 1561-1571.
- Northup PG., et al. "AGA Clinical Practice Update on Surgical Risk Assessment and Perioperative Management in Cirrhosis: Expert Review". Clinical Gastroenterology and Hepatology 17 (2019): 595-606.