



## Cecal Worm Infection in Backyard Fowl of Kashmir Valley

ST Salam\*

Department of Zoology, Amar Singh College, Cluster University Srinagar, India

\*Corresponding Author: ST Salam, Department of Zoology, Amar Singh College, Cluster University Srinagar, India.

Received: March 15, 2018; Published: April 24, 2018

DOI: 10.31080/ASMI.2018.01.0053

### Abstract

Study was carried out for a period of two years on prevalence and pathology of cecal worm *Heterakis gallinarum* on a sample size of 478 domestic chicken (*Gallus g. domesticus* L., 1758), 243 males and 235 females, weighing between 1 - 2.5 kgs collected from the different localities covering almost entire Kashmir valley. The study revealed that prevalence rate of *Heterakis gallinarum* for 1<sup>st</sup> Year of study (2012) was 3.43% (8/233) and for the 2<sup>nd</sup> Year (2013) it was 5.3% (13/245) giving an overall prevalence rate for two year study (Jan 2012 to Dec 2013) to be 4.39% (21/478). Range of intensity of worms was found to be 02 - 55 with Mean intensity of infection observed to be  $20 \pm 2.1$ . Faecal Egg Count was found to be very low i.e <40 eggs per gram of faeces. Histological findings in the infected revealed presence of adult worms in the lumen of intestine along with cellular debris and the infected tissue revealed intense chronic diffuse inflammatory processes with mononuclear and polymorphonuclear (heterophils) leucocyte infiltrations extending up to submucosa. There was sloughing off of the epithelium and lumen was packed with fibrin, red blood cells and tissue debris.

**Keywords:** *Heterakis gallinarum*; Domestic Chicken; Kashmir Valley

### Introduction

Backyard fowl are freely roaming in search of food, thereby exposing themselves to various helminth larvae and eggs. One such common worm infecting chicken is the nematode *Heterakis gallinarum* because backyard chicken frequently feed on earthworms which act as intermediate hosts for this nematode. *Heterakis gallinarum* infection in chicken is usually subclinical but it may function as a vector for *Histomonas meleagridis* (black head) which induces severe pathological lesions in the gut and liver leading to high mortality rates in susceptible hosts [1-3]. The present study was designed to have an idea about the prevalence of the nematode *Heterakis gallinarum* in free ranging chicken of Kashmir Valley and to study the extent of pathology caused by it to the caecum to design some strategy in future to curb this fatal association of these two parasites which causes great economic losses to our backyard poultry industry.

### Materials and Methods

The present two year study was carried from January 2012 to December 2013 and for the study a sample size of 478 domestic chicken (*Gallus g. domesticus* L., 1758), 243 males and 235 females, weighing between 1 - 2.5 kg were selected. Chicken were collected from the different localities covering almost entire Kashmir valley. Individual clinical evaluation and euthanization was carried out according to Zander, *et al* [4]. Nematodes were collected, rinsed in normal saline (0.85%), fixed in hot 70% alcohol and then counted using a stereoscopic microscope. The nematodes were cleared in lactophenol, mounted in glycerin jelly, photographed and identified following Vicente, *et al* [5].

Prevalence was calculated as a percentage of the host population infected at a point in time [6]. Mean intensity was calculated as number of parasites per infested bird.

Faecal examination was carried out following Mc Master's flotation technique. Worm eggs were identified using the keys described by Thienpont, *et al* [7]. Faecal egg counts (FECs) were undertaken within 24 hours by a modification of the McMaster technique with a sensitivity of 50 eggs per gram of faeces [7].

Impression smears from scrapings of the caeca and intestine were stained with Giemsa and Gram's stain.


For pathological studies, fragments of the parasitized caecae and liver fixed in formalin and then routinely processed [8] for paraffin embedding. 5  $\mu$ m sections were cut and stained with hematoxylin and eosin (H&E). Mc Manus periodic acid Schiff (PAS) stain was used to demonstrate protozoan inclusions in the sections. Grocott's stain was used to differentiate with fungal elements. Micrographs were obtained using digital microscope model BX60F-3, Olympus Optical Co. Ltd. (Tokyo, Japan), fitted with the Olympus camera model DP12.

### Results and Discussion


#### Prevalence

*Heterakis gallinarum* is characterized by the presence of oesophageal bulb (Figure 1) and its eggs are more barrel shaped (Figure 2). Prevalence rate for 1<sup>st</sup> Year of study (2012) was found to be 3.43% (8/233) and that for the 2<sup>nd</sup> Year (2013) was found to be 5.3% (13/245) giving an overall prevalence rate for two year study (Jan 2012 to Dec 2013) to be 4.39% (21/478). Reported preva-

lence rates in chicken range from 10.2% to 72.5% in Europe [9,10], 1 to 84% in the USA [11] and 17.28% to 78.8% in Africa [12] but low prevalence in the present study can be attributed to either better adaptability of the nematode to chicks or to the more resistance of chicks which could be either innate or due to better nutrition. *Heterakis gallinarum* worm burden was slightly higher in backyard chickens with poor body conditions [13].




**Figure 1:** Month wise prevalence of *Heterakis gallinarum* in domestic fowl.




**Figure 2:** Season Wise Mean Intensity of infection of *Heterakis gallinarum*.

Range of intensity of worms was found to be 02 – 55 with Mean intensity of infection observed to be  $20 \pm 2.1$ . Faecal Egg Count was found to be very low i.e. < 40 eggs per gram of faeces.

Prevalence of nematode infection and Mean intensity of infection was found to be high during Summer and Autumn seasons of the study (Figure 3 and 4) which can be attributed to decreased resistance of chicks to infection and increased availability of intermediate host (Earthworms) due to high temperature and more rainfall in Summer and autumn (especially August – September).




**Figure 3:** *Heterakis gallinarum* recovered from the domestic fowl showing anterior end revealing mouth, lips and oesophageal bulb.



**Figure 4:** Barrel Shaped Egg of *Heterakis gallinarum*.


**Pathology**

No gross lesions were seen in the caeca during infection of *Heterakis gallinarum*. However histological findings in the caeca revealed presence of adult worms in the lumen of intestine along with cellular debris (Figure 5). Few sections show larvae penetrating the epithelium of cecum (Figure 6). The infected tissue revealed intense chronic diffuse inflammatory processes with mononuclear




**Figure 5:** Photomicrograph of caecum of domestic fowl revealing adult *Heterakis gallinarum* in the lumen, Note the mucosal denudation and cellular debris in lumen H&E 40X.

and polymorphonuclear (heterophils) leucocyte infiltrations extending up to submucosa. TS of the infected caeca showed mucosal erosion with parasites and cellular debris (Figure 5) as observed by other workers Toluidine blue staining of the infected sections clearly reveals the presence of mast cells. There was sloughing off of the epithelium and lumen was packed with fibrin, red blood cells and tissue debris (Figure 7). These observations are in line with the observations of other workers [14-23].



**Figure 6:** Photomicrograph of caecum of domestic fowl revealing adult *Heterakis gallinarum* in the lumen, Note the mucosal denudation and cellular debris in lumen H&E 40X.



**Figure 7:** Caeca of chicken heavily infected with *Heterakis gallinarum* showing plugging of lumen with cellular debris, inflammatory cells and fibrin (HE x40).

## Conclusions

*Heterakis gallinarum* is comparatively harmless nematode but with great potential to act as vector for blackhead causing severe pathological changes in the tissues during coinfection with *Histomonas meleagridis* especially in the warm and wet seasons. Investigations are thus advisable to know in detail local immunological responses of the chicken intestine to the mono infection and co infection *Heterakis gallinarum* with *Histomonas meleagridis* to devise the strategies such as recombinant vaccines and dietary immunomodulation to enhance gut immunity rather than relying on anti-helminthic drugs which contaminate meat and eggs showing great concerns in public.

## Bibliography

- Gibbs BJ. "The occurrence of the protozoan parasite *Histomonas meleagridis* in the adults and eggs of the cecal worm *Heterakis gallinae*". *Journal of Protozoology* 9 (1962): 288-293.
- Springer WT, et al. "Transmission of histomoniasis with male *Heterakis gallinarum* (nematoda)". *Parasitology* 59.2 (1969): 401-405.
- Lund EE and Chute AM. "The reproductive potential of *Heterakis gallinarum* in various species of galliform birds: implications for survival of *H. gallinarum* and *Histomonas meleagridis* to recent times". *International Journal for Parasitology* 4.5 (1974): 455-461.
- Zander DV, et al. "Principles of disease prevention: Diagnosis and control". In *Diseases of poultry*, 10<sup>th</sup> edition, BW Calnek, HJ Barnes, CW Beard, LR McDougald, and YM Saif (eds.). Iowa State University Press, Ames, Iowa (1997): 3-45.
- Vicente JJ, et al. "Nematoides do Brasil. Parte IV: Nematodes de aves". *Revista Brasileira de Zoologia* 12.1 (1995): 1-273.
- Thrusfield M. "Veterinary Epidemiology, 2<sup>nd</sup> edition". Blackwell Scientific, Oxford (1995).
- Thienpont D, et al. "Diagnosing of helminthiasis through coprological examination. 2<sup>nd</sup> edition". Janssen Research Foundation, Beerse, Belgium (1986).
- Bancroft JD and Gamble M. "Theory and Practice of Histological Techniques. 5<sup>th</sup> edition". Harcourt Publ. Ltd., London (2002): 125-131.
- Kokozidou M and Zafeires P. "Prevalence of gastrointestinal parasites in fowls". *Bulletin of the Hellenic Veterinary Medical Society* 47 (1996): 99-103.
- Permin A., et al. "Prevalence of gastrointestinal helminthes in different poultry production systems". *British Poultry Science* 40.4 (1999): 439-443.
- Waters CV, et al. "Status of commercial and non-commercial chickens as potential source of histomoniasis among wild turkeys". *Wildlife Society Bulletin* 22.1 (1994): 43-49.
- Permin A, et al. "A cross sectional study of helminthes in rural scavenging poultry in Tanzania in relation to season and climate". *Journal of Helminthology* 71.3 (1997): 233-240.
- Jansson D, et al. "Parasitforekomst hos tamhos I svenska hobbyflockar". *Sven Veterinartian* 56 (2004): 11-17.
- Kaushik RK and Deorani VPS. "Studies on tissue responses in primary and subsequent infections with *Heterakis gallinae* in chickens and on the process of formation of caecal nodules". *Journal of Helminthology* 43.1-2 (1969): 69-78.
- Riddell C and Gajadhar A. "Caecal and hepatic granulomas in chickens associated with *Heterakis gallinarum* infection". *Avian Diseases* 32.4 (1988): 836-838.
- Khan RW, et al. "Prevalence and gross pathology of helminth infection in domestic fowls of Hyderabad District". *Proceedings of Parasitology* 17 (1994): 4-7.

17. Choudury S and Das MR. "Studies on growth rate and pathological changes in chickens experimentally infected with *Heterakis gallinarum*". *Journal of Veterinary Parasitology* 7 (1993): 81-85.
18. Anna Schwarz., *et al.* "Pathobiology of *Heterakis gallinarum* mono-infection and co-infection with *Histomonas meleagridis* in layer chickens". *Avian Pathology* 40.3 (2011): 277-287.
19. Cortes PL., *et al.* "Histomoniasis in the bursa of Fabricius of Chickens". *Avian Diseases* 48 (2004): 711-715.
20. Eshetu Y., *et al.* "Study of gastro-intestinal scavenging chickens in four rural districts of Amhara region, Ethiopia". *Revue Scientifique et Technique* 20.3 (2001): 791-796.
21. Kemp RL and Reid WM. "Staining techniques for differential diagnosis of histomoniasis and mycosis in domestic poultry". *Avian Diseases* 10.3 (1966): 357-363.
22. McDougald LR. "Other protozoan diseases of the intestinal tract - histomoniasis (blackhead)". In: *Diseases of Poultry* 11<sup>th</sup> edition. Saif YM, Barnes HJ, Glisson JR, Fadly AM, McDougald LR, Swayne DE, (Eds.), Ames, Iowa State Press (2003): 1001-1006.
23. McDougald LR. "Blackhead disease (Histomoniasis) in poultry: A critical review". *Avian Diseases* 49.4 (2005): 462-476.

**Volume 1 Issue 5 May 2018**

**© All rights are reserved by ST Salam.**