



## Intestinal Helminthiasis in School Age Children of Kashmir Valley and the Need for Intervention: A Systematic Review

Showkat Ahmad Wani<sup>1\*</sup> and Sheikh Tanveer Salam<sup>2</sup>

<sup>1</sup>Department of Zoology, Government S. P. College, Cluster University, Srinagar, India

<sup>2</sup>Department of Zoology, Government A. S. College, Cluster University, Srinagar, India

\*Corresponding Author: Showkat Ahmad Wani, Senior Assistant Professor, Department of Zoology, Government S. P. College, Cluster University, Srinagar, J&K, India.

Received: March 30, 2020

Published: April 14, 2020

© All rights are reserved by Showkat Ahmad Wani and Sheikh Tanveer Salam.

### Abstract

Kashmir valley is the most densely populated part of Jammu and Kashmir, India. The enormous population, along with the absence of basic amenities, favourable climatic conditions and underdeveloped public health care system favour the transmission of intestinal helminthiasis. Present paper reviews the prevalence of intestinal helminth infections in the Kashmir Valley and need for control strategies. A survey of published literature by virtue of PUBMED was done. Also many other bibliographic databases were searched to retrieve the relevant articles. After excluding duplicate studies, 11 research papers were retained for this paper. Intestinal helminth infections like those of *Ascaris*, *Trichuris*, *Enterobius* and *Taenia saginata* in Kashmir Valley remain very high. Majority of the infected are young children between the age group of 5 and 14 years. The age group, rural or urban residence, type of water source, boiled or unboiled water, type of defecation site, level of personal hygiene and maternal education were associated with helminth infection. Since the World Health Organization (WHO) recommends chemotherapy for intestinal helminth infections among school-age children, there is no government policy for helminth control in Kashmir Valley. Present article stresses on regular school-based programs to deliver anthelmintics to all school age children so as to reduce the prevalence of helminth infections in Kashmir Valley.

**Keywords:** School-Based Control; Helminths; Chemotherapy; Children; Kashmir Valley

### Introduction

Helminth parasites are the common chronic infection among human beings. In developing countries like India, it is more common to be infected than not [1]. Infection by parasitic helminths thrives in areas which are in need of better housing, clean drinking water, better health care facilities, education and better personal hygiene [2]. This is also typical of almost all rural and urban areas of Kashmir Valley. Children growing up in these areas get soon infected after weaning or chances of them getting infected remain high continuously for the rest of their life [1]. Soil transmitted helminthiasis is a major contributor of diseases among children in developing countries [3]. This high infection rates indicate severe shortage in health care facilities, education and chronic poverty [2]. The increase in the level of school attendance in many developing countries in recent years has helped in the delivery of school-based health services. This is proving an excellent and cost-effective opportunity for educational, economic and developmental gains [4]. In many developing countries like India, Bangladesh, Pakistan etc. the children receive only primary school education,

and it is during this stage when chances of getting infected by helminth parasites are more. These infections could derail the efforts of governments to provide basic school education [5].

There is concrete evidence that school-based chemotherapy against intestinal nematode infections can be delivered at a very low cost [3,5] which can contribute in improving children's general health [6,7], nutritional status [8,9], cognitive ability [10] and school attendance [11]. This paper is based on published literature available on intestinal helminths in children of the Kashmir Valley and seeks to review current trend of infection which is needed in efforts for control.

### Materials and Methods

Information for this review came a survey of published literature by virtue of PUBMED and other bibliographic databases were searched. This study was conducted between January and March 2020, using the key words "School-based control", "Helminths", "Chemotherapy", "children" and "Kashmir Valley". The published literature available on-line were accessed. Full texts of some papers were also obtained from journals at the Iqbal Library of Kashmir

University and at the library of SK Institute of Medical sciences, Srinagar. Also searches were conducted on the basis of links from the cited papers this review was limited to published literature from 2007 to 2015. Websites of organizations like United Nations Children Emergency Fund (UNICEF) and the World Health Organization (WHO) were also thoroughly searched for any relevant data.

## Results and Discussion

After excluding duplicate studies, 11 articles were retained for this review. Prevalence of helminth parasites and other relevant information from the cited articles is summarized in table 1.

Epidemiology of soil-transmitted helminthic infections in Kashmir Valley, reported prevalence studies of soil transmitted hel-

S. No	Author/ Reference	Geographical area	Sample size	Sample setting	Parasites	Prevalence (%)	Comments
1	Wani SA., et al. 2007 [12]	Kupwara district	312	Rural	<i>Ascaris lumbricoides</i>	69.32	Infection highest in children 5-9 years old. Low public and personal hygiene, Multiple infections observed. Infection due to inadequate health education
					<i>Trichuris trichiura</i>	30.76	
					<i>Enterobius vermicularis</i>	7.69	
					<i>Taenia saginata</i>	7.69	
2	Wani SA., et al. 2007 [13]	Srinagar	514	Urban	<i>Ascaris lumbricoides</i>	28.4	Infection highest in children 5-9 years old. Low public and personal hygiene, Multiple infections observed. Infection due to inadequate health education
					<i>Trichuris trichiura</i>	4.9	
					<i>Taenia saginata</i>	3.7	
3	Wani SA., et al. 2007 [14]	Anantnag	432	Rural	<i>Ascaris lumbricoides</i>	77.7	Infection highest in children 5-9 years old. Low public and personal hygiene, Infection were more prevalent in children (10-12 years)
					<i>Trichuris trichiura</i>	31.0	
					<i>Taenia saginata</i>	13	
4	Wani SA., et al. 2007 [15]	Baramullah district	342	Rural	<i>Ascaris lumbricoides</i>	70.84	Infection due to poor disposal of human excreta, Health education and provision of toilet facilities suggested
					<i>Trichuris trichiura</i>	21.65	
					<i>Enterobius vermicularis</i>	12.80	
					<i>Taenia saginata</i>	3.01	
5	Wani SA., et al. 2007 [16]	Budgam district	425	Rural	<i>Ascaris lumbricoides</i>	56.7	Infection due to poor disposal of human excreta, School-based helminth control recommended
					<i>Trichuris trichiura</i>	38.2	
					<i>Enterobius vermicularis</i>	3.7	
6	Wani SA., et al. 2008 [17]	Kashmir Valley	2256	Rural/Urban	<i>Ascaris lumbricoides</i>	68.30	Infection due to poor disposal of human excreta, Health education and provision of toilet facilities suggested, Helminth infections related to level of environmental sanitation, socioeconomic status and water supply
					<i>Trichuris trichiura</i>	27.92	
					<i>Enterobius vermicularis</i>	12.67	
					<i>Taenia saginata</i>	4.60	
7	Wani SA., et al. 2008 [18]	Kashmir valley	382	Rural/Urban	<i>Ascaris lumbricoides and Trichuris trichiura</i>	78.27	Infection due to poor disposal of human excreta, Health education and provision of toilet facilities suggested
8	Wani SA., et al. 2009 [19]	Pulwama district	199	Rural	<i>Ascaris lumbricoides</i>	69.84	Infection due to poor disposal of human excreta, Multiple infections observed. Infection due to inadequate health education
					<i>Trichuris trichiura</i>	31.65	
					<i>Enterobius vermicularis</i>	16.80	
					<i>Taenia saginata</i>	3.01	
9	Baba AA., et al. 2009 [20]	Random	131	Radom	<i>Ascaris lumbricoides</i>	63.0	Infection highest in children 5-9 years old. Low public and personal hygiene, Multiple infections observed. Infection due to inadequate health education
10	Wani SA., et al. 2010 [21]	Gurez Valley	352	Rural	<i>Ascaris lumbricoides</i>	71.18	Infection highest in children 5-9 years old. Low public and personal hygiene
					<i>Trichuris trichiura</i>	26.42	
					<i>Enterobius vermicularis</i>	13.92	
					<i>Taenia saginata</i>	5.39	
11	Lone R., et al. 2011 [22]	Budgam district	396	Rural	<i>Ascaris lumbricoides</i>	54.9	Infection highest in children 5-9 years old. Low public and personal hygiene, Multiple infections observed. Infection due to inadequate health education
					<i>Trichuris trichiura</i>	32.49	
					<i>Enterobius vermicularis</i>	2.57	
					<i>Taenia saginata</i>	7.69	

**Table 1:** Prevalence estimates of intestinal helminthic infections in Kashmir Valley.

minths since the 2007 have indicated that the triad of *Ascaris lumbricoides*, *Trichuris trichiura*, *Enterobius vermicularis* and *Taenia saginata* species are common infections in Kashmir Valley (Table 1). These studies indicate that the prevalence of soil transmitted helminth infections mostly *Ascaris lumbricoides* has not changed in the past 30 years [23].

Most papers commented on the unhygienic practice of people defecating in open or in open pit latrines. The condition has not changed much unfortunately, as there is hardly any success in the construction of septic and hygienic latrines to rural Kashmir Valley. The drainage systems in Srinagar city of Kashmir valley unplanned and malfunctioned and is often blocked. This results in widespread dispersal of ova and larvae of these helminths especially during summer season. Nwosu's findings indicated a similar case scenario [24]. In an urban area, a survey of faecal samples collected from various sites showed that 96.3% of the samples contained eggs of *Ascaris lumbricoides* [25]. This playgrounds act as a main source of helminth infection for children [26,27]. Illiteracy and poor socioeconomic conditions of families has been related with higher helminth infestations among children [24,28]. Helminth infection rate among children and socioeconomic status of parents has a strong correlation and in Kashmir valley, helminthiasis is still a disease of poor. Adekunle, *et al.* (1986), also found the same correlation [29]. Some cultural practices favour spread of infection. Since water is used to clean the body after defecation and also there are religious beliefs that people should feed from common bowl, a common practice in many rural areas may also contribute for a high prevalence of intestinal helminth infections [26]. In most of the studies, bulk of infection has been found in children between the age groups of 5 - 14 year [24,29-32]. If this age group (5 - 14 years) is provided periodic anthelmintic drugs, it will definitely help in infection control in general population. Certainly, schools are the best places for its implementation [5].

#### Effects of helminth infections in school-aged children

Most of the children in helminth endemic areas seldom realize their full potential. The reasons of this underachievement are webbed [10], has been hard to prove [33]. However, many studies have recognized helminth infections as one of the causes, which affect the intellectual and physical development of children [10]. The morbidity is determined by the intensity of helminth infections [34,35], but many times clinical symptoms appear at light infections also [10]. Infection by multiple types of helminthes increases morbidity [29,30]. In some areas, people with multiple infections are more common than others which may have single or are uninfected [10]. The most common consequence of helminth infections

is that children seldom achieve their genetic potential for growth and physical development and suffer from malnutrition and iron deficiency anaemia [10]. Heavy *Trichuris* infection in children results in whip worm dysentery syndrome, which is recognized by retarded growth and anaemia [36]. Intense infections by both *Ascaris* and *Trichuris* are related with protein energy malnutrition [8]. The global public health importance of ascariasis and trichuriasis has been comprehensively reviewed [34-38]. The disability-adjusted life-years (DALYs) lost to intestinal helminth infections is very high, when compared with other infections [39] none of the studies by Savioli, *et al.* the main reason for high DALYs values have been attributed to intestinal helminth infections, as *Ascaris* infections lead to growth retardation and *Trichuris* causes decreased school performance [40]. If school children in heavily infected areas are provided periodic anthelmintic drugs, about 70% of the helminthiasis can be effectively managed [41]. In one of the studies in Kashmir valley, it has been proved beyond doubt that there occurs improvement in weight gain in children after deworming [42], suggesting that deworming for intestinal helminths may result in a period of growth in previously infected children. There are many studies from Kashmir valley, which show that *Ascaris* may lead to intestinal lumen blockage and inflammation of appendix, hepatic and bile ducts [20-22,43,44]. It is not easy to estimate mortality rate due to helminth infections in Kashmir Valley, because people go for widespread self-medication and public health care system is not effective. It is estimated that world over about 135000 direct deaths occur due to intestinal helminth infections annually [40].

#### Control strategies

Presently there are no control program for intestinal helminth infections in Kashmir Valley as school health services are poor. Researchers in parasite control had over time advocated for improved sanitation and health education so that environment doesn't get contaminated with helminth infective stages. It has also been advocated that children with heavy infections should be treated over a time, especially during the winter months, when transmission conditions are least favourable [13]. Unfortunately, these recommendations, have not been implemented and the prevalence of helminth infection still remain high across the Kashmir valley.

Chemotherapy can be introduced for intestinal helminth control at a very low cost in health care systems [3,10,33]. Drake and Bundy (2001) has suggested school health services as of paramount importance. These programs offer an opportunity to deliver anthelmintics to a largest number of children at a very low cost [10]. These programmes have therefore attracted the interest by policy makers in developing countries like India [46]. There are examples

that in developing countries, school health services such as providing anthelmintics can be provided at low cost and more conveniently [3,5,33]. For these reasons, de-worming has now become an essential component of school health programs in many developing countries [45]. School-based programs also reach children who are not enrolled in school, usually the most affected group [9] and serve to transmit health education to the whole population.

### Conclusion

Soil transmitted helminthiasis are still highly prevalent children in the age group of 5 - 14 years in Kashmir Valley and a primary cause of morbidity. Lack of personal hygiene, contaminated surroundings, lack of education, and absence of basic amenities, poverty and favourable climate are major reasons which sustain the transmission, but there has not been effort to control at government level. The effectiveness of intervention using anthelmintics at six monthly intervals has been proved to be cost-effective and feasible in other parts of the globe. This is the time for government agencies in Kashmir Valley to frame a policy regarding the school-based programs and relieve children of the burden of intestinal helminth infection, so that they can achieve their maximum potential.

### Conflict of Interest

No conflict of interest exists.

### Bibliography

1. Awasthi S., et al. "Helminth infections". *British Medical Journal* 327 (2003): 431-433.
2. Crompton DW. "How much human helminthiasis is there in the world?" *Journal of Parasitology* 85 (1999): 397-403.
3. World Bank. "World development report: Investing in Health". Oxford University Press, New York (1993).
4. Del Rosso JM., et al. "Class action, improving school performance in the developing world through better health and nutrition". The World Bank, Washington, DC (1996).
5. The Partnership for Child Development. "Better health, nutrition and education for the school-aged child". *Transactions of Royal Society of Tropical Medicine and Hygiene* 91 (1997): 1.
6. Stephenson LS., et al. "Treatment with a single dose of albendazole improves growth of Kenyan schoolchildren with hookworm, *Trichuris trichiura*, and *Ascaris lumbricoides* infections". *American Journal of Tropical Medicine and Hygiene* 41 (1989): 78-87.
7. Stoltzfus RJ., et al. "Effects of the Zanzibar school-based deworming program on iron status of children". *American Journal of Clinical Nutrition* 68 (1998): 179-186.
8. Stephenson L., et al. "Weight gain of Kenyan school children infected with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* is improved following once- or twice-yearly treatment with albendazole". *Journal of Nutrition* 123 (1993): 656-665.
9. Beasley NM., et al. "The health of enrolled and non enrolled children of school age in Tanga, Tanzania". *Acta Tropical* 76 (2000): 223-229.
10. Drake LJ., et al. "Multiple helminth infections in children: impact and control". *Parasitology* 122 (2001): 573-581.
11. Simeon DT., et al. "Treatment of *Trichuris trichiura* infections improves growth, spelling scores and school attendance in some children". *Journal of Nutrition* 125 (1995): 1875-1883.
12. Wani SA., et al. "Plasma Proteins in Children with Helminth Infections". *Journal of Parasitic Diseases* 31.2 (2007): 165-168.
13. Wani SA., et al. "Epidemiology of Gastrointestinal Helminths in the School Children of District Budgam Kashmir". In *Advances in Fish and Wildlife Ecology and Biology*, Edition. Kaul 4 (2007): 219-225.
14. Wani SA., et al. "Prevalence of Intestinal Parasites and Associated Risk Factors among School children in Srinagar City, Kashmir, India". *Journal of Parasitology* 93.6 (2007): 1541-1543.
15. Wani SA., et al. "Prevalence of Gastrointestinal Helminths in the Primary School Children of District Anantnag, Kashmir and their Effect on the Haemoglobin Status". In *Advances in Fish and Wildlife Ecology and Biology*, Edition. Kaul 4 (2007): 226-232.
16. Wani SA., et al. "Intestinal Helminth Infections among Children of District Baramulla of Kashmir Valley". *Oriental Sciences* 12.1 (2007): 49-54.
17. Wani SA., et al. "Soil-Transmitted Helminths in Relation to Haemoglobin Status among School Children of the Kashmir Valley". *Journal of Parasitology* 94.3 (2008): 591-593.
18. Wani SA., et al. "Intestinal Helminths in a population of children from Kashmir Valley India". *Journal of Helminthology* 82 (2008): 313-317.

19. Wani SA, et al. "Intestinal Helminths and Associated Risk Factors in Children of District Pulwama, Kashmir, India". *Indian Journal of Medical Microbiology* 27.1 (2009): 81-82.
20. Baba AA, et al. "Intestinal ascariasis: the commonest cause of bowel obstruction in children at a tertiary care center in Kashmir". *Pediatric Surgery International* 25.12 (2009): 1099-102.
21. Wani SA, et al. "Intestinal Helminthiasis in Children of Gurez Valley of Jammu and Kashmir State, India". *Journal of Global Infectious Diseases* 2.2 (2010): 91-94.
22. Lone R, et al. "Recent patterns and risk factors of intestinal helminthes infection among school children in Kashmir, India". *Archives of Clinical Microbiology* 2. 3 (2011): 2.
23. Khuroo MS. "Ascariasis". *Gastroenterology Clinics of North America* 25.3 (1996): 553-577.
24. Nwosu ABC. "The community ecology of soil-transmitted helminth infections of humans in a hyperendemic area of southern Nigeria". *Annals of Tropical Medicine and Parasitology* 75 (1981): 197-203.
25. Fashuyi SA. "The prevalence of helminth eggs in human feces deposited on the streets of Lagos, Nigeria". *West African Journal of Medicine* 2 (1983): 135-138.
26. Akogun OB. "Some social aspects of helminthiasis among the people of Gumau District, Bauchi State, Nigeria". *Transactions of Royal Society of Tropical Medicine and Hygiene* 92 (1989): 193-196.
27. Umeche N. "Helminth ova in soil from children's playgrounds in Calabar, Nigeria". *Central African Journal of Medicine* 35 (1989): 432-434.
28. Ayanwale FO, et al. "The epidemiology of human intestinal helminthiasis in Ibadan, South Western Nigeria". *International Journal of Zoonosis* 9 (1982): 69-72.
29. Adekunle LV, et al. "Family influence on incidence of intestinal parasites among Nigerian children". *Journal of Royal Society of Health* 106 (1986): 66-68.
30. Adeyeba OA, et al. "A survey of gastrointestinal parasites in a local government area of south-west Nigeria". *International Journal of Zoonosis* 11 (1984): 105-110.
31. Udonsi JK, et al. "Assessment of the effectiveness of primary health care interventions in the control of three intestinal nematode infections in rural communities". *Public Health* 107 (1993): 53-60.
32. Arinola O, et al. "Age and sex graded helminth infections in a Nigerian village". *East African Medical Journal* 72 (1995): 110-112.
33. De Silva NR. "Impact of mass chemotherapy on the morbidity due to soil-transmitted nematodes". *Acta Tropical* 86 (2003): 197-214.
34. Cooper ES, et al. "Trichuris is not trivial". *Parasitology Today* 4 (1988): 301-306.
35. Roche M, et al. "The nature and causes of "hookworm anemia". *American Journal of Tropical Medicine and Hygiene* 15 (1966): 1029-1102.
36. O'Lorcain P, et al. "The public health importance of *Ascaris lumbricoides*". *Parasitology* 121 (2000): 51-71.
37. Stephenson LS, et al. "The public health significance of *Trichuris trichiura*". *Parasitology* 121 (2000): 73-95.
38. Crompton DW. "The public health importance of hookworm disease". *Parasitology* 121 (2000): 39-50.
39. Chan MS. "The global burden of intestinal nematode infections-fifty years on". *Parasitology Today* 13 (1997): 438-443.
40. Savioli L, et al. "Progress in the prevention and control of schistosomiasis and soil-transmitted helminthiasis". *Parasitology International* 53 (2004): 103-113.
41. Asaolu SO, et al. "The role of health education and sanitation in the control of helminth infections". *Acta Tropical* 86 (2003): 283-294.
42. Wani SA, et al. "Soil Transmitted Helminth infection and its effect on nutritional status of children in Kashmir". *Oriental Science* 14 (2009): 105-111.
43. Shiekh KA, et al. "Mechanical small bowel obstruction in children at a tertiary care centre in Kashmir". *African Journal of Pediatric Surgery* 7.2 (2010): 81-85.
44. Javid G1, et al. "Ascaris-induced liver abscess". *World Journal of Surgery* 23.11 (1999): 1191-1194.

45. Savioli L, *et al.* "Intestinal parasitic infections: a soluble public health problem". *Transactions of Royal Society of Tropical Medicine and Hygiene* 8 (1992): 353-354.
46. Curtale F, *et al.* "The School Health Programme in Behera: an integrated helminth control programme at Governorate level in Egypt". *Acta Tropical* 86 (2003): 295-307.

#### Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

**Website:** <https://www.actascientific.com/>

**Submit Article:** <https://www.actascientific.com/submission.php>

**Email us:** [editor@actascientific.com](mailto:editor@actascientific.com)

**Contact us:** +91 9182824667