

## Study of Resistance Mechanisms and Evaluation of Biofilm Detection Tests in Clinical Isolates of *Pseudomonas aeruginosa* Circulating in Yaounde, Cameroon

Yves Le Grand Napa Tchuedji<sup>1,2\*</sup>, Hortense Gonsu Kamga<sup>3,4</sup>, Emilia Lyonga Mbamyah<sup>2,3</sup>, Anicette Chafa Betbeui<sup>3,4</sup>, Ibrahim Assana Raoul<sup>5</sup>, Giselle Fonyuy Bongnyang<sup>6</sup>, Etienne Guenou<sup>2</sup> and Francois-Xavier Etoa<sup>1</sup>

<sup>1</sup>Department of Microbiology, Laboratory of Microbiology, University of Yaoundé I, Cameroon

<sup>2</sup>Department of Public Health, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Cameroon

<sup>3</sup>Department of Microbiology, Haematology and Infectious Diseases, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Cameroon

<sup>4</sup>Laboratory of Bacteriology, University Teaching Hospital, Cameroon

<sup>5</sup>University and strategic institute of the estuary, Cameroon

<sup>6</sup>General Hospital of Yaoundé, Cameroon

\*Corresponding Author: Yves Le Grand Napa Tchuedji, Department of Microbiology, Laboratory of Microbiology, University of Yaoundé I, Cameroon.

Received: October 20, 2021

Published: November 29, 2021

© All rights are reserved by Yves Le Grand Napa Tchuedji, et al.

### Abstract

**Background:** Antibiotic resistance is a priority problem around the world. The occurrence of SARS-CoV-2 in a context where the resistance of *Pseudomonas aeruginosa* is increasing, could promote the cohabitation of these microorganisms and an increase in the risk of clinical damage and mortality in infected patients.

**Aim:** The aim of this study is to determine resistance strategies of clinical isolates of *Pseudomonas aeruginosa* that circulate in Yaoundé, Cameroon.

**Method:** A descriptive cross-sectional study was carried out from May 2019 to November 2020. A randomized sampling was conducted. The study of resistance phenotypes was carried out by the disc diffusion method and the double disc synergy method in Mueller-Hinton agar. The study of biofilm formation was performed using the tissue culture plate in Mueller-Hinton broth and Congo red agar.

**Results:** A total of 30 clinical isolates of *Pseudomonas aeruginosa* were collected from 300 patients. Pus presented the highest proportion of *Pseudomonas aeruginosa* followed by urine and blood with frequencies of 60%, 16.66% and 13.33% respectively. We detected 76.66% of multi-resistant *Pseudomonas aeruginosa* with high resistance frequencies to the majority of betalactams and the aminoglycoside, attributed to ertapenem (100%), cefotaxime (100%), ticarcillin (86.66%) and tobramycin (60%). These strains presented 10% and 3.33% of carbapenemases and Extended-spectrum beta lactamases (ESBLs) respectively. According to their ability to produce biofilm, 16.66% and 83.33% were non-biofilm formers with Plate culture method and culture method on Congo red agar respectively.

**Conclusion:** This study demonstrated that *P. aeruginosa* strains circulating in the Center region have several enzymatic mechanisms of resistance to antibiotics associated with a high production of biofilm. The study of biofilm formation shows a significant difference between the tissue culture technique and the Congo red agar culture technique.

**Keywords:** *Pseudomonas aeruginosa*; Resistance Phenotypes; Carbapenemases; ESBLs; Resistance Genes

## Abbreviations

ESBLs: Extended-spectrum Beta Lactamases; *P. aeruginosa* : *Pseudomonas aeruginosa*; SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus 2; MAR: Multiple Antibiotic Resistance; CAS-FM: Antibiogram Committee of the French Microbiology Society; OD: Optical Density

## Introduction

In the past, multi-resistant bacteria were first associated with nosocomial infections, but today they cause several infections in the community, causing considerable economic losses. Antibiotic resistance is present in all countries with an increased risk of clinical damage and mortality in patients exposed to resistant bacteria [1]. This situation is still very worrying in Africa, where several factors contribute to the emergence and transmission of resistant strains. *Pseudomonas aeruginosa* is known to be one of the major causes of opportunistic infections in immunocompromised patients associated with high mortality. In the world, and particularly in Cameroon, the situation of antimicrobial resistance is even more worrying, this can be attributed to reckless prescription of antibiotics by unqualified personnel, self-medication, the widespread use of counterfeit drugs whose dosage in active compounds is unknown, poor sanitation and hygiene in medical facilities and usage in subtherapeutic doses [2]. Many studies have shown that subminimal inhibitory concentrations of antibiotics could induce mutagenesis in clinical isolates of *Pseudomonas aeruginosa* [3].

*P. aeruginosa* is one of the multi-resistant bacteria which possess several virulence factors and which infect the respiratory tract. The occurrence of SARS-CoV-2 in a context where resistance to antibiotics is increasing, could promote the coexistence of these microorganisms in the respiratory tract. This coinfection can cause an amplification of the physiopathological mechanisms and accelerate the clinical and functional degradation of patients carrying SARS-CoV-2.

Very little data are available on the exhaustive list of the different strategies of resistance found in *Pseudomonas aeruginosa* in the Cameroonian population. This implies a surveillance of resistance focused on finding mechanisms of resistance of *P. aeruginosa* circulating in Cameroon. The aim of this study is to determine resistance strategies of *Pseudomonas aeruginosa* in Yaounde.

## Methods

### Isolation and handling of bacterial strains

The isolates of *Pseudomonas aeruginosa* from 300 patients attending the University Teaching Hospital of Yaounde, the Central Hospital of Yaounde, the General Hospital, and the Centre Pasteur of Cameroon were collected between May 2019 and October 2020. Bacteria isolates were conserved in Brain Heart infusion broth supplemented with glycerol (10%) and transported in ice to the bacteriology laboratory of the University Teaching Hospital.

### Identification

Presumptive identification of the isolates was performed in each collection site using the catalase, oxidase, mannitol and citrate Simmons agar test. The isolates were revived on Mueller-Hinton agar and further identified using the Hajna Kliger media to assess glucose, lactose fermentation and API 20 NE (BioMérieux, France) following the manufacturer's instructions.

### Antimicrobial susceptibility testing

Antimicrobial Susceptibility tests were conducted according to the recommendations of the Antibiogram committee of the French Microbiology Society using the disc diffusion method on Mueller-Hinton agar (CASFM) [4]. The bacterial inoculum adjusted to 0.5 Mc Farland was inoculated on Mueller-Hinton agar. Fifteen minutes later, thirteen antibiotic discs were deposited on the Mueller-Hinton agar. The following discs were used: tircacillin (75 µg), tircacillin/clavulanic acid (85 µg), piperacillin (30 µg), piperacillin/tazobactam (36 µg), ceftazidim (30 µg), cefotaxim (30 µg), imipenem (10 µg), meropenem (10 µg), ertapenem (10 µg), tobramycin (10 µg), ciprofloxacin (5 µg), netilmicin (30 µg), norfloxacin (10 µg). This was left at room temperature for 15 minutes and then incubated at 37°C for 24 hours. Inhibition zones were measured after incubation and scored as susceptible, intermediate or resistant. The multiple antibiotic resistance (MAR) index was determined by dividing the total number of resistances to antimicrobials for each isolate to the total number of tested antimicrobials.

### Detection of extended-spectrum beta lactamases

Extended-spectrum beta lactamases (ESBLs) are enzymes found in certain bacteria and are responsible for their resistance to antibiotics such as penicillins and cephalosporins. The double disc

synergy was used to screen all the isolates for ESBLs production as recommended by CASFM [4]. Antibiotic discs ceftazidime, cefotaxime, cefepime were placed 30 mm around ticarcillin/ clavulanic acid disc in pre-inoculated Mueller Hinton agar. The plates were incubated aerobically at 37 ° C for 24h. The production of heart-shaped clear zones that distorts the zone of inhibition indicates positive results for ESBLs production.

#### Detection of carbapenemase

The search for carbapenemases was carried out by the disc diffusion method according to the recommendation of CASFM [4]. The bacterial inoculum prepared according to 0.5 McFarland standard was inoculated on Mueller-Hinton Agar. After 15 minutes, imipenem disc (10 µg) was deposited on the plate. The whole is left to prediffused for 15 minutes and then incubated aerobically for 24 hours at 37°C. The observation of inhibition zones less than 28 mm testifies to a presence of carbapenemases.

#### Detection of cephalosporinases

The search for cephalosporinases was carried out by the disc diffusion test on solid medium according to the recommendation of CASFM [4].

The bacterial inoculum prepared according to 0.5 McFarland standard was inoculated on Mueller-Hinton agar. After 15 minutes cephalosporin discs, ticarcillin, ticarcillin/clavulanic acid was deposited on the surface of the agar plate. The whole is left prediffused for 15 minutes and then incubated aerobically for 24 hours at 37°C. The observation of resistance to these antibiotics confirm the presence of cephalosporinases.

#### Determination of biofilm formation by tissue culture plate method

The study of biofilm formation was carried out according to the method described by O'Toole, *et al.* [5]. To perform this, 150 µL of brain heart infusion broth was added to each well of a sterile microtitre plates. Then 10 µL of inoculum adjusted to 0.5 McFarland was inoculated into each well. Negative and positive controls were included, containing 150 µL of brain heart infusion broth and 150 µL of brain heart infusion broth containing strains producing biofilm respectively. The whole was incubated for 48 hours at 37 ° C. After incubation, each well was emptied gently using a syringe, avoiding contact with the wall of the well. The wells were then wa-

shed 3 times with distilled water and then stained with 1% crystal violet for 5 minutes. Then the wells were washed twice and then dried. Ethanol-acetone (75:25) was pipetted into each well in order to detach any biofilm which is on the wall of the well and the optical density read at 570 nm. The micro-well plates were agitated for one minute prior to measuring the turbidity. The tests were repeated three times. The categorization of isolates was done according to the recommendations of Christensen, *et al.* [6], which states:  $OD \leq OD_t$ : not producing biofilm;  $OD_t \times 2 \leq OD < OD_t \times 4$ : moderate production of biofilm;  $OD_t \times 4 \leq OD$ : highly biofilm producer.


#### Determination of biofilm formation by culture on Congo red agar method

The study of biofilm formation was carried out according to the method described by Hou, *et al.* [7]. For this, the pure colonies of *Pseudomonas aeruginosa* were inoculated on Congo red agar supplemented with saccharose 10 g/L. The Petri dishes were then incubated at 37 ° C for 24-48 hours. After incubation the presence of strains producing slimes give rough black colonies, non-producing strains give red colonies with a smooth surface, and strains with varying phenotypes give colonies with a red outline and black center [8].

## Results

### Frequency of *Pseudomonas aeruginosa*

Samples were taken from 300 patients. Thirty cultures were positive for *Pseudomonas aeruginosa* (10%). Among the 30 strains, 13 were isolated at the Centre Pasteur of Cameroon and 17 in hospital with 76.47% of hospitalized patient. Figure 1 shows the proportion of infected patients according to the unit of origin.



**Figure 1:** Distribution of infected patients according to the unit of origin.

As shown by figure 1, the Centre Pasteur of Cameroon have the highest frequency of *Pseudomonas aeruginosa* followed by the unit of medicine and intensive care with a frequency of 43.33%, 13.33%

and 13.33% respectively. The strains of *Pseudomonas aeruginosa* were isolated from different samples (Table 1).

	Urine	Wound	Bedsore	Blood	Lochia	Pus	Total
N <sub>o</sub> positive (%)	5 (16,66)	1 (3,33)	1 (3,33)	4 (13,33)	1(3,33)	18(60)	30 (100)

**Table 1:** Frequency of *Pseudomonas aeruginosa* by sample.


Pus presented the highest proportion of *Pseudomonas aeruginosa* (60%) followed by urine (16, 66%), blood (13, 33%) wounds (3.33%) bedsores (3.33%) and lochia (3.33%).

The multi-resistance index varied from 0.23 to 0.84 with 39.11% of the micro-organisms which having a multi-resistance index greater than 0.50.

**Antibiotic susceptibility profile**

The susceptibility profile (Figure 2) revealed a high resistance to the majority of betalactams and the aminoglycoside, attributed to ertapenem (100%), cefotaxime (100%), tircacillin (86.66%) and tobramycin (60%). We also noted resistance to netilmicin, imipenem and meropenem with respective frequency of 26.66%, 10% and 20%. High resistance to beta-lactam inhibitors was observed for piperacillin/tazobactam and tircacillin/clavulanic acid with frequencies of 86.66 and 50% respectively. Resistance to quinolones varying from 36.66 for norfloxacin to 40% for ciprofloxacin.

The distribution of *Pseudomonas aeruginosa* according to age group (Figure 3) shows a high frequency of multi-resistance in each group.



**Figure 3:** Frequency of *Pseudomonas aeruginosa* isolates according to age and multi-resistance.


Patients aged between [25-50[ years and [50-65[ years were the most infected. However, a high proportion of multi-resistant bacteria was observed between the ages of [25-50[, [50-65[ and [65-100[ years although the age group of [65-100[ years had the highest proportion of multi-resistant bacteria.

**Resistance phenotypes of *Pseudomonas aeruginosa***

Table 3 presents the resistance phenotypes detected in this study. The wild phenotypes were conserved in all the isolates. However, we detected the presence of ESBLs and carbapenemases with respective frequencies of 3.33% and 10%. No positive result was found for cephalosporinases.

**Study of biofilm formation by *Pseudomonas aeruginosa***

The study of biofilm formation was carried out using two methods described in previous studies. The results obtained by these



**Figure 2:** Antibiotic susceptibility profile of *Pseudomonas aeruginosa*.  
R: Resistant; I: Intermediate; S: Susceptible.

Among the 30 strains collected, 76.66% were multidrug resistant. The determination of the multi-resistance index did not consider natural resistance. Table 2 shows the frequency of *Pseudomonas aeruginosa* according to the multi-resistance index.

	Wild	ESBLs	Carbapenemases	Cephalosporinases
Frequency n(%)	100	3,33	10	0

**Table 3:** Resistance phenotypes of *Pseudomonas aeruginosa*.

two techniques show a difference in frequency of the strains producing biofilms. Table 4 shows the frequency of *Pseudomonas aeruginosa* according to the plate culture technique and the Congo red agar culture technique.

Plate culture method				Congo Red Agar culture method		
Biofilm formation	Strong bio-film formers	Moderate biofilm formers	Non-biofilm formers	biofilm formers	Variable biofilm formers	Non-biofilm formers
Frequency n (%)	9 (30%)	16 (53,33%)	5 (16,66%)	3 (10%)	2 (6,66%)	25 (83,33%)

**Table 4:** Frequency of *Pseudomonas aeruginosa* isolates according to their ability to produce Biofilm.

Thirty percent of the isolates of *Pseudomonas aeruginosa* were strong producers of biofilms and 53.33% carried out a moderate production of biofilm with plate culture technique, while 10% and 6,66% were biofilm formers and variable biofilm formers after culture on Congo red agar. The production of biofilm is one of the resistance strategies used by *Pseudomonas aeruginosa* to resist antibiotics. However, 16,66% and 83, 33% were non-biofilm formers with Plate culture method and culture method on Congo red agar respectively. This difference ( $P < 0.05$ ) observed between the two techniques on the frequency of bacteria which do not produce biofilms is statistically significant.

## Discussion

Among the 300 patients sampled, 10% were infected with *Pseudomonas aeruginosa*. These results are similar to those of Barbier and Wolff [9], in France who found a frequency of 10% and lower to that of Clotilde, *et al.* [10], who found a frequency of 15.4% in the city of Douala. This city is the economic capital of Cameroon where the majority of commercial and industrial activities are carried out leading to overcrowding of the city, unsanitary environment and non-observance of hygiene measure [11]. The Centre Pasteur of Cameroon have the highest frequency of *Pseudomonas aeruginosa*. This could be explained by the fact that this laboratory receives the majority of patients compared to laboratories of hospitals.

Concerning unit of origin in hospitals, medicine and intensive care unit had the highest frequency of *Pseudomonas aeruginosa*. Many studies show that medicine and intensive care have a high proportion of infection compared to other units. The work carried

out by Gonsu, *et al.* [12], also revealed that medicine and intensive care units had the highest proportion of infections compared to pediatrics, surgery, neonatology and ophthalmology. Pus presented the highest proportion of *Pseudomonas aeruginosa* (60%) followed by urine (16, 66%) and blood (13, 33%). These results are different from those of Ndip, *et al.* (2005) who found a high prevalence of *P. aeruginosa* in urine (30%) and wounds (28%) compared to pus (16%). This could be explained by its ubiquitous nature.

A high resistance to the majority of betalactams was observed with a frequency of 100% for ertapenem, 100% for cefotaxime, 86.66% for ticarcillin and 60% for tobramycin. This family of drugs has been affected by the constant increase in bacteria resistance [13]. We also noted 10% and 20% of resistance to imipenem and meropenem respectively. Carbapenem-resistant *P. aeruginosa* is one of the resistant ESKAPE pathogens in which there is a critical need for the development of new antibiotics to treat infections [14]. Resistance was noted to norfloxacin and ciprofloxacin with a frequency of 36,66% and 40% respectively. This bacteria has developed and maintained resistance to quinolones for many years [15]. The highest proportion of multi-resistant strains was observed in the age group of [65-100[ years. This could be explained by the immune failure in elderly patients which promotes the growth of bacteria and gives them time to develop resistance mechanisms.

We detected 10% of carbapenemase and 3,33% of ESBLs. The study of Madaha, *et al.* [16], shows that the double disc synergy test have a low capacity to detect ESBLs on *Pseudomonas aeruginosa* compared to the combined disc method. This constitute a limit



in the detection of resistance mechanisms because of the permanent unavailability of the antibiotic combined disc in developing country as Cameroon.

The difference observed between plate culture method and culture method on Congo red agar on the frequency of bacteria which do not biofilms is statistically significant. The work carried out by Madaha, *et al.* [16], revealed that under the same temperature, only 17.33% of isolates of *Pseudomonas aeruginosa* did not produce biofilms with plate culture method while the work of Kenge, *et al.* [17], showed a proportion of 42.85% of *Pseudomonas aeruginosa* strains which did not produce biofilm with the Congo red agar culture technique. This confirms the low proportion of biofilm detection by the Congo red agar culture method. It is therefore important to determine the performance of these tests in order to make an optimal choice for the detection of biofilm formation.

### Ethical Approval

This study was approved by the Ethical committee of the delegation of public health for the center Region of Cameroon under the approval number 0191AAR/MINSANTE/DRSPL/BCASS. Anonymity of participants and confidentiality of results were scrupulously respected.

### Conclusion

This study presented many resistance mechanisms of *Pseudomonas aeruginosa* circulating in Yaoundé Cameroon, such as enzymes that inactivate antibiotics and the development of biofilms. The Carbapenemases were the most represented. The difference observed between the tests for the detection of biofilm, proves a need to determine the performances, such as sensitivity and the specificity to make an optimal choice of these tests.

### Acknowledgment

We thank Rosanne NGOME who participated in the collection of strains at the Centre Pasteur of Cameroon, Michelle DJUIDJE, Melissa TSANGA and Ornella MINLONG who participated in the collection of strains at the Central Hospital of Yaoundé.

### Bibliography

1. Battraud Paul. "La résistance aux antibiotiques, un mythe ou une réalité?" Thèse pour le diplôme d'état de docteur en pharmacie (2017): 1-128.
2. Nko'o Amvene Samuel and Marie Claire Okomo Assoumou. "Antibiotiques et résistance bactérienne : un sujet de préoccupation". *Health Sciences and Disease* 14.4 (2013): 1-2.
3. Valencia Estela Ynés, *et al.* "Ciprofloxacin-mediated mutagenesis is suppressed by subinhibitory concentrations of amikacin in *Pseudomonas aeruginosa*". *Antimicrobial Agents and Chemotherapy* 61.3 (2017): e2107-e2116.
4. CA-SFM. Recommendations 1 (2020): 1-9.
5. O'Toole, *et al.* "Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development". *Molecular Microbiology* 30.2 (1998): 295-304.
6. Christensen, *et al.* "Adherence of coagulase-negative staphylococci to plastic tissue culture plates: a quantitative model for the adherence of staphylococci to medical devices". *Journal of Clinical Microbiology* 22.6 (1985): 996-1006.
7. Wenbo Hou, *et al.* "The Association for Research in Vision and Ophthalmology". *Investigative Ophthalmology and Visual Science* 53.9 (2012): 5624-5631.
8. Mathur, *et al.* "Detection of biofilm formation among the clinical isolates of Staphylococci: An evaluation of three different screening methods". *Indian Journal of Medical Microbiology* 24.1 (2006): 25-29.
9. François Barbier and Michel Wolff. "Multirésistance chez *Pseudomonas aeruginosa* Vers l'impasse thérapeutique?" *Medicine/Sciences* 26.11 (2010): 960-968.
10. Njall, pouth Clotilde, *et al.* "Écologie bactérienne de l'infection nosocomiale au service de réanimation de l'hôpital Laquintinie de Douala, Cameroun". *Pan African Medical Journal* 14.140 (2013): 1-8.
11. NAPA TCHUEDJI, *et al.* "Epidemiology of *Pseudomonas aeruginosa* and Its Resistance in Cameroon: Systematic Review and Meta Analysis". *Global Advanced Research Journal of Microbiology* 9.5 (2020): 069-072.
12. Gonsu Kamga, *et al.* "Caractérisation phénotypique des souches de *Pseudomonas aeruginosa* isolées dans la ville de Yaoundé (Cameroun)". *African Journal of Pathology and Microbiology* 4 (2015): 1-4.
13. Worthington Roberta and Melander Christian. "Overcoming Resistance to  $\beta$ -Lactam Antibiotics". *Journal of Organic Chemistry* 78.9 (2013): 4207-4213.
14. Founou Raspail Carrel, *et al.* "Clinical and economic impact of antibiotic resistance in developing countries: A systematic review and meta-analysis". *Plos One* 12.12 (2017): e0189621.

15. Robert Hancock Rew and Speert David. "Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and impact on treatment". *Drug Resistant Update* 3.4 (2000): 247-255.
16. Madaha Estelle Longla., *et al.* "Occurrence of blaTEM and blaCTXM Genes and Biofilm-Forming Ability among Clinical Isolates of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* in Yaoundé, Cameroon". *Microorganisms* 8.708 (2020): 1-22.
17. Kengne Landry. "Contribution à la lutte contre les contaminants microbiens des locaux et des collections des bibliothèques au moyen d'huiles essentielles". Thèse de Doctorat. Université de Yaoundé I. Cameroun (2019): 1-236.

**Volume 4 Issue 12 December 2021**

**© All rights are reserved by Yves Le Grand Napa Tchuedji, *et al.***