

## Role of Physical Activity in Severe COVID-19 Outcomes

Attapon Cheepsattayakorn<sup>1,2\*</sup>, Ruangrong Cheepsattayakorn<sup>3</sup> and Porntep Siriwanarangsun<sup>1</sup>

<sup>1</sup>Faculty of Medicine, Western University, Pathumtani Province, Thailand

<sup>2</sup>10<sup>th</sup> Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand

<sup>3</sup>Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

**\*Corresponding Author:** Attapon Cheepsattayakorn, 10<sup>th</sup> Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand.

**Received:** August 22, 2022

**Published:** September 01, 2022

© All rights are reserved by **Attapon Cheepsattayakorn, et al.**

Prevalence of obesity, a marker of evolution of the world-population lifestyle and global prevalence of physical inactivity were greater than the prevalence of the worldwide prevalence of tobacco smokers with potentially being a greater mortality cause than tobacco smoking [1]. All COVID-19 patient cohorts demonstrated increase in severe-COVID-19 risk among persons with obesity by the index of adiposity (Body-Mass Index (BMI)) [2]. Regular physical activity decreases the risk of developing chronic respiratory diseases, cardiovascular diseases, and type 2 diabetic mellitus [2]. A recent study on physical activity among 48,440 COVID-19-adult patients in the United States revealed that COVID-19 patients with consistently physical inactivity had a greater risk of hospital admission (Odds Ratio (OR): 2.26; 95% Confidential Interval (CI): 1.81-2.83), intensive-care-unit (ICU) admission (OR: 1.73; 95% CI: 1.18-2.55), and mortality (OR: 2.49; 95% CI: 1.33-4.67) that were related to COVID-19 than patients with consistently meeting physical activity guidelines [3].

In conclusion, efforts to promote physical activity following the guidelines should be the routine-medical-care's incorporation and public-health-agencies' prioritization. Tools for assessing and targeting physical activity in clinical practice are urgently needed.

### Bibliography

1. Wen CP and Wu X. "Stressing harms of physical inactivity to promote exercise". *Lancet* 380 (2012): 192-193.

2. Ross R., et al. "Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association". *Circulation* 134 (2016): e653-e699.
3. Sallis R., et al. "Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48,440 adult patients". *British Journal of Sports Medicine* 55 (2021): 1099-1105.