



Antibacterial Activity of Tuber Extracts of *Cyclamen rohlfsianum* against Some Human Pathogenic bacteria, *In vitro*

Asma A Elamari^{1*}, Wafa S Zubi¹, Tahani M Awin², Nawal M. Buzgaia² and Abduslam M Elmogassapi¹

¹Botany Department, Science Faculty, University of Benghazi, Benghazi, Libya

²Chemistry Department, Science Faculty, University of Benghazi, Benghazi, Libya

*Corresponding Author: Asma E Elamari, Botany Department, Science Faculty, University of Benghazi, Benghazi, Libya.

Received: August 19, 2024

Published: September 24, 2024

© All rights are reserved by José de Jesús Alba-Romero., et al.

Abstract

Cyclamen rohlfsianum is an endemic plant species to Al-jabal al-Akhdar, Libya. Lately, it got some attention in term of it's antimicrobial activity and use to treat the infections that cause by bacteria and candida. This study investigated the antibacterial activity of *Cyclamen rohlfsianum* tuber extracts against four species of human pathogenic bacteria (*Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter* sp., and *Escherichia coli*) using disc diffusion method. Four solvents were used in the extraction process; which were hexane, chloroform, ethyl acetate, and ethanol. The obtained results from all tuber extracts at concentration 0.01 mg/ml showed clear antibacterial activity against both *S. aureus* and *P. aeruginosa*. Ethanol tuber extract was the most effective against both *S. aureus* and *P. aeruginosa* (with inhibition zone 0.8 mm and 0.7 mm in diameter; respectively) compared to hexane, chloroform, and ethyl acetate tuber extracts. On the other hand, only chloroform plant extract inhibited the growth of *Enterobacter* sp. (with inhibition zone 0.5 mm in diameter). In addition, hexane and chloroform plant extract showed antibacterial activity against *E. coli* (with inhibition zone 0.3 and 0.5 mm in diameter, respectively) while the other applied plant extracts had no inhibitory activity against this bacterium. The study confirmed the efficacy of *Cyclamen rohlfsianum* tuber extracts as natural antibacterial and suggested that they may be used as a therapy to treat infectious diseases caused by the bacteria under study.

Keywords: Antibacterial Activity; *Cyclamen rohlfsianum*; Disc Diffusion Method

Introduction

For many decades, scientists in medical field have been concern about explore new antimicrobial effective substances from different sources such as medicinal plants [9]. Plants have been used as sources of medicinal and today a significant proportion of the population in the developed and developing countries depends on herbal medicines; especially in the areas where infectious diseases are endemic and modern health care facilities are unavailable.

In Libya, many people are consider traditional system of medicine as a first choice before visiting health care center that may be because of the fact that traditional medicine is safer and not costly [18]. That encourages the scientific community to incorporate the plants as very important part of the researches that concern about

human health. Many studies have been reported that plants contain many biologically active compounds that affect the growth of human pathogenic microorganisms [4]. That leads to discover newer, safer, and possibly more effective drugs against human pathogenic bacteria and fungi.

Cyclamen rohlfsianum is one of the medicinal plants that belong to the family *Primulaceae*. It is native to North Africa, and it is one of the tenderest cyclamen species. A tetraploid *Cyclamen rohlfsianum* is endemic to Al-Jabal Al-Akhdar [6,16], and it is locally known as "Racuf" [8,22]. It is well known that *Cyclamen rohlfsianum* is a poisonous plant because it contains cyclamen glycoside [24] but it still commonly use in the therapy of diabetes though it is a toxic plant [7]. Many reports indicated that *Cyclamen* sp. contains active com-

pounds such as alkaloid, flavinoid, phenolic compound, terpenoid, and saponins that may be attributed to its effect on the growth of human pathogenic bacteria and fungi [25, 26]. In addition, many studies have been showed that *Cyclamen spp.* had antimicrobial activity against wide variety of human pathogenic bacteria and candida [7,13]. indicated the antimicrobial effect of *Cyclamen persicum* tuber extracts against some of human pathogenic bacteria and candida. Another study showed that ethanol extracts of *Cyclamen mirabile* had a strong effect against *Staphylococcus aureus* [10]. Also there was a study showed that the aqueous extract of leaves and tubers of *Cyclamen rohlfsianum* had a strong inhibitory effect on the growth of some pathogenic bacteria (*Escherichia coli*, *Staphylococcus aureus*, *Proteus vulgaris*, and *Pseudomonas aeruginosa*) [2]. Therefore, the main goal of the current study is to evaluate the antimicrobial activity of tuber extracts of *Cyclamen rohlfsianum* against some human pathogenic bacteria.

Material and Methods

Collection plants and preparation of plant extracts

Fresh samples of *Cyclamen rohlfsianum* (Family- *Primulaceae*) were collected from Al-jabal al-Akhdar, Libya in March 2023. The plant was identified and classified at Department of Botany, University of Benghazi. The collected plants were transported to the laboratory and washed with water and dried in the shade for 14 days. The dried tubers were separated from each plant and then crushed into a fine powder using an electric blender [21]. The method was applied with some modification as following, 50g of the powder of *Cyclamen rohlfsianum* tubers was filled in the thimble and extracted with 100 ml of each solvent (hexane, chloroform, ethyl acetate, and ethanol; separately) using a Soxhlet apparatus under controlled temperature for 24 hours then all the extracts were evaporated [2]. Finally, all the crude extracts were dissolved in the 1% (v/v) dimethyl sulphoxide (DMSO). Only one concentration of the extracts was prepared (0.01 mg/ml) and stored at 4 °C in airtight bottles until further use.

Collection of bacteria

Bacterial species were obtained from the microbiology laboratory of Children Hospital, Benghazi, Libya. Four human pathogenic bacteria; which are *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter sp.* and *Escherichia coli* of clinical importance were used in this study. Bacteria were identified using BD Phoenix

for accurate identification. The isolates were cultured in nutrient agar (NA) at 37 °C and were maintained on nutrient agar slants at 4 °C.

Preparation of bacterial suspension

Bacterial stock cultures were sub-cultured onto nutrient agar (NA) plates and incubated overnight at 37°C. Three to four bacterial colonies were inoculated into 10 ml of Mueller Hinton Broth (MHB) and incubated at 37 °C then the overnight bacterial suspensions were diluted and adjusted [27].

Determination of antibacterial assay

For the detection of antibacterial activities of the *Cyclamen rohlfsianum* disc diffusion method (Kirby-Bauer antibiotic testing) was applied with modification [17]. Mueller-Hinton Agar (MHA) was used as basal medium for culture of bacteria (MHA was prepared by suspending 38 g in 1000 ml of distilled water). The medium was sterilized by autoclaving at 15 lbs pressure and 121°C for 15 minutes, cooled to 45-50°C and then 15 ml of the sterilized medium was poured into sterile petri plates. After solidification, 100 µl of each prepared isolate was inoculated on MHA plates using sterilized spreaders. Gentamycin (10 µg) was applied as a positive control, and DMSO was applied as negative control (DMSO would not affect the growth of bacteria) [11]. Sterile discs were filled with (20µl) of each extract (at the prepared concentration (0.01 mg/ml) and allowed to diffuse at room temperature for 1 hour then the plates were incubated at 37 °C for 24h. Triplets of the experiment were maintained for each bacterial species to ensure reliability. Finally after incubation time the diameter of the inhibitory zones formed around each disc were measured in mm and recorded.

Statistical analysis

The experiments were designed according to the complete random design. Statistical analysis was performed using Minitab 17 program and ANOVA variance analysis tables. The averages were compared using Tukey's test at P <0.05.

Results

The results from the experiments were showed in table (1) and figure (1). In this study, *Cyclamen rohlfsianum* was investigated to evaluate its antibacterial activity against four species of human pathogenic bacteria; which are *Staphylococcus aureus*, *Pseudomo-*

nas aeruginosa, *Enterobacter sp.* and *Escherichia coli* using a disc diffusion method. Gentamycin (10 µg) was applied as a positive control, and DMSO was applied as negative control for the antibacterial assays.

In general, the study showed that all used tuber extracts at concentration 0.01 mg/ml exhibited a varying degree of antimicrobial activity against all bacteria tested (table 1). Hexane, chloroform, ethyl acetate, and ethanol extracts of the tubers of *Cyclamen rohlfsianum* showed inhibitory effect against both *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Ethanol extract was the most effective plant extract against *Staphylococcus aureus* and *Pseudomonas aeruginosa* (with inhibition zone 0.8 and 0.7 mm in diameter; respectively) compared to the other extracts (hexane, chloroform, and ethyl acetate extracts). On the other hand, both ethanol and ethyl acetate extracts of the used plant had no effect on the growth on both *Enterobacter sp.* and *Escherichia coli*. Also, the re-

sults of this study showed that only chloroform tuber extract affect the growth of *Enterobacter sp.* (with inhibition zone 0.5 mm in diameter). In addition, both hexane and chloroform plant extract affect the growth of *Escherichia coli* (with inhibition zone 0.3 and 0.5 mm in diameter, respectively). However, ethyl acetate and ethanol plant extracts did not record any inhibitory activity against *E. coli*.

The results also showed the extent of an effect of the antibiotic used (Gentamycin) on the growth of the bacteria under study where it affected the growth of *S. aureus*, *P. aeruginosa*, *Enterobacter sp.* and *E.coli* (with inhibition zone 0.1, 0.1, 0.7, and 0.6 mm in diameter; respectively). It was clear that *E. coli* was the most resistant to both tuber extracts and gentamycin while *S. aureus* and *P. aeruginosa* were the most sensitive to both of them. In general, the comparison between the obtained results from tuber extracts and the antibiotic applied showed that the antibiotic had higher antibacterial activity than the tuber extracts against the growth of the tested bacteria but the plant extracts still affected the growth of

	The diameter of inhibition zone of bacteria (mm)			
	Concentration in mg/ml (0.01mg/ml)			
	Bacterial spp.			
Plant extracts	<i>S. aureus</i>	<i>P. aeruginosa</i>	<i>Enterobacter sp</i>	<i>E. coli</i>
Hexane extract	0.5 ± 0	0.6 ± 0.17	0.00	0.3 ± 0.57
Chloroform extract	0.5 ± 0.00	0.5 ± 0.00	0.5 ± 0.00	0.5 ± 0.00
Ethyl acetate extract	0.4 ± 0.15	0.5 ± 0.05	0.00	0.00
Ethanol extract	0.8 ± 0.11	0.7 ± 0.0	0.0	0.0
Gentamycin (Positive control)	1.0 ± 0.11	1.0 ± 0.36	0.7 ± 0.15	0.6 ± 0.1
DMSO (Negative control)	0.0	0.0	0.0	0.0

Table 1: Antibacterial activity of *Cyclamen rohlfsianum* tuber extracts against tested bacterial species.

(0.00): No inhibition zone

the bacteria. Finally, the results showed that there were significant differences of the effect of different tuber extracts (hexane, chloroform, ethyl acetate, and ethanol) on the growth of all used bacteria $P < 0.05$.

Discussion

Due to the medical importance of *Cyclamen rohlfsianum* and its commonly used in alternative medicine in *Libya*, researchers pay attention of this plant [13]. In *Libya*, there is lack of research on the biological activities of endemic plants; therefore, this study was

conducted. The results showed that *Cyclamen rohlfsianum* tubers extracts had inhibitory effect against the growth of both Gram-positive and Gram-negative bacteria under study and this result agreed with [2]. In addition, the results agreed with the study which indicated that *Cyclamen persicum* tuber extracts had inhibitory effect on the growth of some bacteria and candida species [13]. Moreover, another study where showed that aqueous extract of leaves and tubers of *Cyclamen rohlfsianum* affect the growth of *Escherichia coli*, *Staphylococcus aureus*, *Proteus vulgaris*, and *Pseudomonas aeruginosa*, and that agree with our results [2].




Figure 1: Effects of hexane, chloroform, ethyl acetate, and ethanol plant extracts against tested bacterial species.

The results also agreed with the study that indicated that aqueous extract of tubers of *Cyclamen purpurascens L.* had a potential effect on the growth of some types of human pathogenic bacteria [27]. In addition, several studies have shown that the extracts obtained from the *Cyclamen* species inhibit the growth of many species of bacteria and candida at various concentrations [2,3,29]. In general, many reports indicated that plants contain some active compounds such as flavonoids, aldehydes, ketones, saponins, alcohols, and phenolic compounds that inhibit the growth of microorganisms such as bacteria and candida [30,32,33, 34], and it is well known that these compounds are abundant in the *Primulaceae* family that *Cyclamen rohlfsianum* belongs to [3,7].

In common, our results showed that Gram-negative bacteria were more resistant to the tuber extracts, and it may be because the outer-membrane permeability barrier limits access of the antibacterial agents to their targets in the bacterial cell [9,12]. The active metabolic components of plant extracts such as terpenoid, alkaloid, and phenolics bind with enzymes and proteins found in bacterial cell membrane that lead to cell death or may inhibit the activity of the enzymes needed for amino acid biosynthesis [24,35]. The results showed that *Enterobacter sp.* was the most resistant to the plant extract and only chloroform plant extract inhibited the growth of this bacterium with inhibition zone 0.5 mm in diameter. In contrast, *S.aureus* was the most sensitive to both plant extracts and gentamycin. The Inhibitory effect of *Cyclamen rohlfsianum*

could be referred to present active metabolites such as alkaloid, flavinoid, phenolic compound, terpenoid, and saponins as what early said. Therefore, this research would recommend the importance of *Cyclamen rohlfsianum* for pharmacological use and to continue researches for purifying the active ingredient and to be subjected to furthermore standardized drug assays.

Conclusions

Cyclamen rohlfsianum tuber extracts inhibit the growth of some important human pathogenic bacteria, and it may be attributed to present of alkaloid, flavinoid, phenolic compound, terpenoid, and saponins. Therefore, this research recommends conducting more studies on the endemic Libyan plants in term of detection and purification of the active compounds and in order to incorporate these plants in the treatment of the infections caused by such bacteria.

Bibliography

1. AA Abdulraziq. "Antibacterial Activity of *Cyclamen rohlfsianum* against two Species of *Xanthomonas* 2.1 (2022): 71-75.
2. AA Abdulraziq and MS Salih. "Bio-effect of aqueous extract of *Cyclamen rohlfsianum* on some pathogenic bacteria". *Global Lib IOSR Journal of Pharmacy and Biological Sciences* 4.5 (2022): 6-8.

3. AA Abdulrazziq and MS Salih. "Effect of Aqueous Extracts of *Arum cyreniacum* on Some Negative and Positive Gram bacteria". *Al-Mukhtar Journal of Sciences* 35.1 (2020): 60-68.
4. AW Bauer, et al. "Antibiotic susceptibility testing by standardized single disk method". *American Journal of Clinical Pathology* 45 (1966): 493-496.
5. C Grey-Wilson. "Cyclamen. - Timber Press, Portland (2002).
6. FA Elabbar, et al. "Isolation and Identification Of Some Compounds From *Cyclamen Rohlfsianum* (Primulaceae) From Libya". *Scientific Reviews and Chemical Communications* 4.1 (2014): 1-10.
7. FM El-Mokasabi. "Floristic Composition and Traditional Uses of Plant Species at Wadi Alkuf, Al- Jabal Al-Akhder, Libya". *American-Eurasian Journal of Agricultural and Environmental Science* 14.8 (2014): 685-697.
8. GA Cordel. "Natural products in drug discovery - Creating a new vision 1 (2000): 261-273.
9. G Okmen, et al. "The antibacterial activities against mastitis pathogens of *Cyclamen mirabile* Hildebr, tubers and its non-enzymatic antioxidant activities". *European Journal of Experimental Biology* 4.2 (2014): 370-374.
10. JR Zgoda and JR Porter. "A Convenient Microdilution Method for Screening Natural Products against Bacteria and Fungi". *Pharmaceutical Research* 39.3 (2001): 211-225.
11. K Bayoub, et al. "Antibacterial activities of the crude ethanol extracts of medicinal plants against *Listeria monocytogenes* and some other pathogenic strains". *African Journal of Biotechnology* 9.27 (2010): 4251-4258.
12. M Al-zuabe, et al. "Antimicrobial Effect of *Cyclamen persicum* Tuber Extracts Against Bacteria and *Candida* Species". *Journal of Pure and Applied Microbiology* 13.1 (2019): 107-116.
13. MK Yousufi. "To Study Antibacterial Activity of *Allium Sativum*, *Zingiber Officinale* and *Allium Cepa* by Kirby-Bauer Method". *Journal of Pharmacy and Biological Sciences* 4 (2012): 06-08.
14. MM Miloud and NA Senuss. "Antibacterial activity of leaf extracts of *Silene succulent* Forsk. (Caryophyllaceae) against clinically important bacteria". *Academia Journal of Microbiology Research* 9.1 (2021): 013-020.
15. M Sibusawa N and K Ogawa. "Production of interspecific hybrids between *Cyclamen persicum* Mill. and *C. rohlfsianum* Aschers. or *C. persicum* and *C. libanoticum* Hirdebr". *Bull. Tokyo Agr. Exp. Stn* 27 (1997): 9-15.
16. M Yousufi. "To Study Antibacterial Activity of *Allium Sativum*, *Zingiber Officinale* and *Allium Cepa* by Kirby-Bauer Method". *Journal of Pharmacy and Biological Sciences* 4 (2012): 2278-3008.
17. NY Naguib, et al. "A comparative study on the productivity and chemical constituents of various sources and species of *Calendula* plants as affected by two foliar fertilizers". *The Journal of Applied Sciences Research* 1.2 (2005): 176-189.
18. R Puupponen-Pimiä, et al. "Antimicrobial properties of phenolic compounds from berries". *Journal of Applied Microbiology* 90.4 (2005): 494-507.
19. SL Sukanya, et al. "Antimicrobial activity of leaf extracts of Indian medicinal plants against clinical and phytopathogenic bacteria". *African Journal of Biotechnology* 8.23 (2009): 6677-6682.
20. S Mohammadi, et al. "Antimicrobial activity of methanolic root extracts of *Euphorbia condylocarpa* against pathogenic bacteria". *Advanced Studies in Biology* 7.2 (2015): 55-64.
21. SM El-Darier and FM El-Mogaspi. "Ethnobotany and relative importance of some endemic plant species at El-Jabal El-Akhdar Region (Libya)". *World Journal of Agricultural Sciences* 5.3 (2009): 353-360.
22. S Mohammadi, et al. "Antimicrobial activity of methanolic root extracts of *Euphorbia condylocarpa* against pathogenic bacteria". *Advanced Studies in Biology* 7.2 (2015): 55-64.
23. SR Chant. "Flowering Plants of the World, 1st, Oxford University Press, Oxford (1979).

24. AO Gill and RA Holley. "Disruption of *Escherichia coli*, *Listeria monocytogenes* and *Lactobacillus sakei* cellular membranes by plant oil aromatics". *International Journal of Food Microbiology* 108 (2006): 1-9.
25. A Mishra., *et al.* "*Bauhinia variegata* leaf extracts exhibit considerable antibacterial, antioxidant, and anticancer activities". *BioMed Research International* (2013): 915436.
26. L Wang., *et al.* "Fingerprint analysis and quality consistency evaluation of flavonoid compounds for fermented *Guava* leaf by combining high-performance liquid chromatography time-of-flight electrospray ionization mass spectrometry and chemometric methods". *Journal of Separation Science* 39 (2016): 3906-3916.
27. CH Teh., *et al.* "Determination of antibacterial activity and minimum inhibitory concentration of larval extract of fly-via resazurin-based turbidometric assay". *BMC Microbiology* 17.36 (2017): 1-8.
28. LP Stanojevic., *et al.* "Aqueous extract of wild cyclamen tubers (*Cyclamen Purpurascens* L.)-a potential source of natural antioxidants and antimicrobial agents". *Quality of Life* 16.1-2 (2018): 13-19.
29. A Abdi Ali., *et al.* "The Study of Synergistic Effects of n. butanolic *Cyclamen* coum Extract and Ciprofloxacin on inhibition of *Pseudomonas aeruginosa* biofilm formation". *Biological Journal of Microorganisms* 3.12 (2015): 25-32.
30. A Akgul. "Antimicrobial activity of black cummin (*Nigella sativa* L.) essential oil". *Gazi Universitesi Eczacilik Fakultesi Dergisi* 6.1 (1989): 63-68.
31. S Sindhu and S Manorama. "Screening of *Polycarpaea corymbosa* lam. (Caryophyllaceae) for its *in vitro* antioxidant activity". *Asian Journal of Pharmaceutical and Clinical Research* 5.4 (2012): 175-178.
32. S Amaral., *et al.* "Plant extracts with anti-inflammatory properties-A new approach for characterization of their bioactive compounds and establishment of structure-antioxidant activity relationships". *Bioorganic and Medicinal Chemistry* 17.5 (2009): 1876-1883.
33. SY Lin., *et al.* "Antioxidant, anti-semicarbazide-sensitive amine oxidase, and anti-hypertensive activities of geraniin isolated from *Phyllanthus urinaria*". *Food and Chemical Toxicology* 46.7 (2008): 2485-2492.
34. AJ Driscoll., *et al.* "Disk diffusion bioassays for the detection of antibiotic activity in body fluids: applications for the pneumonia etiology research for child health project". *Clinical Infectious Diseases* 54 (2012): S159.
35. S Burt. "Essential oils: their antibacterial properties and potential application in foods: A review". *International Journal of Food Microbiology* 94.3 (2004): 223-253.