



## Point of Care Diagnostics for Acute Organophosphorus Poisoning: An Emerging Need for Management

**Sheemona Chowdhary and Dibyajyoti Banerjee\***

*Department of Experimental Medicine and Biotechnology, Postgraduate Institute of Medical Education and Research, Chandigarh, India*

**\*Corresponding Author:** Dibyajyoti Banerjee, Department of Experimental Medicine and Biotechnology, Postgraduate Institute of Medical Education and Research, Chandigarh, India.

**Received:** June 26, 2018; **Published:** July 01, 2018

Poisoning with organophosphorus pesticides can occur due to suicidal or non-suicidal exposure [1]. It is a serious health concern, especially in the developing nations. According to WHO, in rural Asia majority of the deaths occur due to suicidal ingestion of organophosphorus pesticides as they are freely available in the market [2]. Toxicity due to organophosphates leads to depression of Acetylcholinesterase activity in the nervous system thereby causing excessive cholinergic stimulation. Measurement of AChE is recommended in all clinical cases of OP poisoning [1]. Diagnosis is usually based on the history of exposure and the clinical manifestations as reliable enzyme assay methods are not available. As the decrease in AChE level correlates with the severity of poisoning, therefore, a rapid and reliable enzyme assay method is required for early diagnosis and management of the poisoning [3, 4].

At present, the recommended method for the measurement of AChE activity in case of organophosphorus poisoning has certain pitfalls [5-9]. Researchers have tried to address such limitations, but the major problem is the wide inter-individual difference in the baseline value of acetylcholinesterase [4,10-13]. Till date, there is no established reference range of acetylcholinesterase which makes the interpretation of the results difficult especially in cases of less severe poisoning.

With the growing incidence of poisoning, there is an emerging need for the bedside assessment of acetylcholinesterase activity. This can be met by a point of care testing platform which can enable timely management of the poisoning. There is ongoing research for the development of point of care testing tools for the detection of acetylcholinesterase activity in the context of organophosphorus poisoning [14,15]. We have recently reported that AChE estimation is still important for understanding organophosphorus poisoning and feel we feel that AChE is the target biomarker for point of care testing tool development [16].

At present, there is no recommended point of care testing tool for the desired purpose. Therefore, we believe that with the advent of a cost effective, user-friendly and rapid point of care testing tool which can overcome the problem of baseline measurement of acetylcholinesterase activity can potentially reduce the interface between diagnosis and management of acute organophosphorus poisoning.

### Bibliography

1. Chowdhary S., *et al.* "Acute organophosphorus poisoning". *Clinica Chimica Acta* 431 (2014): 66-76.
2. World Health Organisation (WHO). "Clinical Management of Acute Pesticide Intoxication: Prevention of Suicidal Behaviours". (2008).
3. Namba T. "Cholinesterase inhibition by organophosphorus compounds and its clinical effects". *Bulletin of World Health Organisation* 44.1-3 (1971): 289-307.
4. Lotti M. "Cholinesterase inhibition: complexities in interpretation". *Clinical Chemistry* 41.12 (1995): 1814-1818.
5. Ellman GL., *et al.* "A new and rapid colorimetric determination of acetylcholinesterase activity". *Biochemical Pharmacology* 7.2 (1961): 88-95.
6. Simpson RJ., *et al.* "Estimation of free thiols and disulfide bonds using Ellman's reagent". *Cold Spring Harbor Protocols* 3 (2008): 1-2.
7. Riener CK., *et al.* "Quick measurement of protein sulfhydryls with Ellman's reagent and with 4,4'-dithiodipyridine". *Analytical and Bioanalytical Chemistry* 373.4-5 (2002): 266-276.
8. Sinko G., *et al.* "Limitation of the Ellman method: cholinesterase activity measurement in the presence of oximes". *Analytical Biochemistry* 370.2 (2007): 223-227.

9. Walmsley TA, *et al.* "Effect of daylight on the reaction of thiols with Ellman's reagent, 5,5'-dithiobis (2-nitrobenzoic acid)". *Clinical Chemistry* 33.10 (1987): 1928-1931.
10. George PM and Abernethy MH. "Improved Ellman procedure for erythrocyte cholinesterase". *Clinical Chemistry* 29.2 (1983) :365-368.
11. Wille T, *et al.* "Evaluation of 6,6'-dithionicotinic acid as alternative chromogen in a modified Ellman method-comparison in various species". *Toxicology Mechanisms and Methods* 21.7 (2011): 533-537.
12. Augustinsson KB, *et al.* "A new approach to determining cholinesterase activity in samples of whole blood". *Clinica Chimica Acta* 89.2 (1978): 239-252.
13. Lessenger JE and Reese BE. "Rational use of cholinesterase activity testing in pesticide poisoning". *Journal of American Board of Family Practice* 12.4 (1999): 307-314.
14. Seonhee L, *et al.* "Foldable paper-based analytical device for the detection of an acetylcholinesterase inhibitor using an angle-based readout". *Sensors and Actuators B: Chemical* 273 (2018): 322-327.
15. Shuo W, *et al.* "Gold nanoparticles dissolution based colorimetric method for highly sensitive detection of organophosphate pesticides". *Sensors and Actuators B: Chemical* 238 (2017): 427-433.
16. Chowdhary S and Banerjee D. "Whether lack of Measurement of Erythrocyte Cholinesterase or Acetyl Cholinesterase Activity providing us misleading Information in Organophosphorous Exposure?" *Indian Journal of Clinical Biochemistry* (2018).

**Volume 2 Issue 5 August 2018**

**© All rights are reserved by Sheemona Chowdhary and Dibyajyoti Banerjee.**