



## Goat Milk Fatty Acids on Brain Growth and Functions: A Systematic Review

**Salleh Noraini, Siti Norazah Said, Abdullah nurlaily, Nooriah Salleh, Nor Aripin Shamaan, Khairun NB Nor Aripin and Nor Azila Noh\***

*Department of Medical Science, Faculty of Medicine and Health Sciences, Universiti Sains Islam Malaysia, Kuala Lumpur, Malaysia*

**\*Corresponding Author:** Nor Azila Noh, Department of Medical Science I, Faculty of Medicine and Health Sciences, Universiti Sains Islam Malaysia, Kuala Lumpur, Malaysia.

**Received:** December 20, 2018; **Published:** February 08, 2019

### Abstract

A systematic review was conducted to elucidate the role of goat milk fatty acids on neural development and functions. The search was systematically done on six databases; SCOPUS, EMBASE, Cumulative Index to Nursing and Allied Health Literature (CINAHL), OVID, Proquest Education Journal and PUBMED. The review was conducted until June 2018. The search focused on any studies related to the neural developmental effect of goat milk in human and animal brain. The search strategies used keywords related to "goat", "milk" and "brain". There was no restriction on study designs, publication date or language. A total of 3185 abstracts were initially identified. After reviewing the titles, abstracts and full papers using pre-determined inclusion and exclusion criteria, no human-related study was found. However, four studies showed positive effects on brain growth and development and reflex maturation in animals. Therefore, more research should be done to elucidate the relationship between goat milk and brain physiology especially in human.

**Keywords:** Brain; Goat Milk; Linoleic Acid; Lipid; Systematic Review

### Abbreviations

CLA: Conjugated Linoleic Acid; CNS: Central Nervous System; CSD: Cortical Spreading Depression; DGW: Dried Goat Whey; DNA: Deoxyribonucleic Acid; EFA: Essential Fatty Acids; EPM: Elevated Plus Maze; NEU: Neuraminidase; ORT: Object Recognition Test; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; PUFA: Polyunsaturated Fatty Acid

### Introduction

The goat milk is the milk of choice by those who are allergic to cow and sheep milk [1]. A study showed that allergic reactions towards goat milk were not as severe as cow milk [2]. Studies conducted by researchers stated that weight gain, height, and blood serum contents of calcium, thiamin, vitamin A, riboflavin and hemoglobin in human and rats who had consumed goat milk surpassed those who consumed cow milk [4,5]. Biochemical composition of goat milk may varies according to the breed, type of food, lactating state and environmental conditions [6]. Due to these factors, the quantity and the quality of goat milk fat content such as omega-6 and omega-3 also varies [7].

Goat milk fat is one of the most complex natural fats comprising approximately 400-500 fatty acids. Omega-3 and omega-6 fatty acids are important for brain development [8]. Essential fatty acids (EFA) is one of the subsets of fatty acids involved in brain development especially in neonates, which are incapable of synthesizing their own EFA [9]. During pregnancy and lactation, polyunsaturated fatty acids are crucial for neonatal development where the supply of these fatty acids is derived from maternal nutrition [10-13].

Polyunsaturated fatty acids are initially catalysed by enzymes initially into two precursors: linoleic acid (18:2) and alpha-linolenic acid (18:3) [14-16]. Linoleic acid consists of a few geometrical isomers which are generally designated as conjugated linoleic acid (CLA) [17]. CLA is found in goat milk. Linoleic acid from the omega-6 group and linolenic acid from the omega-3 group are important for the development of the nervous system especially in the final trimester of intrauterine life. During this stage of pregnancy, brain development is accelerated [18].

The quality and the amount of fatty acids are important for the development of the neonatal nervous system. Malnutrition impairs

brain growth and the development by decreasing replication of the cell cycle, reducing brain total deoxyribonucleic acid (DNA), and restricting dendritic arborization [19]. Thus, these activities will reduce neuronal connectivity. Deprivation of the essential nutrients in the neonate and toddler will affect the development of the cerebellum, particularly on its synaptic connectivity [20,21]. Furthermore, the EFA will mainly affect brain structural composition and myelin sheaths [10]. These include alterations in the electroencephalographic activity, motor and cognitive development, visual and auditory evoked responses and social abilities [19].

Goat milk has been highlighted as one of the high quality raw materials for food manufacturing [22,23]. Most of the studies conducted were mainly on neurodegenerative diseases and microbiological effects of goat milk. However, the ability of goat milk to improve brain development and functions is still unknown. Based on many effects of goat milk fatty acid on neural development and functions, it is important to systematically review the present literature on the benefits of goat milk beneficial to brain development. This helps in explaining goat milk effects on brain development and function.

## Methods

A comprehensive and updated database was used to catalogue the literature on goat milk and its influence on brain development and function. The evidence database was assembled according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for systematic reviews (<http://www.prisma-statement.org/>). The method was adopted from Cochrane Reviewers' Handbook.

### Search methods for identification of studies

#### Computer search

A broad search for the literature was performed by incorporating both electronic and manual components. The electronic search was accomplished using SCOPUS, EMBASE, Cumulative Index to Nursing and Allied Health Literature (CINAHL), OVID, Proquest Education Journal and PUBMED, which were conducted until June 2018. Based on the databases, relevant papers via the main keywords: goat, milk and brain were identified. Further search terms are presented below. The study was accomplished through the screening steps by referring to the inclusion and exclusion criteria.

The search terms strategies include all terms related to main keywords as follows:

1. Goat\*: capra\*; caprine\*; hircus\*; OR hircine\*;
2. Milk\*: dairy\*; butter\*; cream\*; yogurt\*; cheese\*; OR whey\*;
3. Brain\*: nerv\*; cereb\*; cortex\*; cortical\*; OR neuro\*;

An asterisk after a term means that all terms that begin with that root were included in the search.

#### Step 1

All the identified and relevant abstracts were compiled in an individual library in Endnote (X7.0.1) software. All the duplicated data identified and removed from the library. Abstracts were reviewed based on the following inclusion criteria: 1) human and animal studies, 2) sources from goat milk and its derivative products and 3) studies related to brain and central nervous system (CNS). If the criteria were not met because of inadequate information, the abstract was set aside for further evaluation.

#### Step 2

The abstracts were reviewed independently by three investigators and were selected based on their consensus according to the same criteria used in Step 1. If the consensus was not reached, the abstract was set aside for further evaluation.

#### Step 3

Full-text articles of abstracts selected in Step 2 were retrieved and reviewed. Studies were included if they met the following criteria: 1) used transgenic mice, 2) involved goat milk's fatty acids and 3) showed measurements of a central nervous system and physical activity. The exclusion criteria were 1) studies that did not discuss the effectiveness of the goat milk relevant to the brain activity, 2) studies that involved drugs as intervention and 3) studies that involved goat gene as milk promoter.

#### Review of reference lists

The reference lists of all articles selected in Step 3 were reviewed, and the full texts of the potentially related studies were examined.

#### Data Extraction

For data extraction, the screened studies had to report the outcomes based on the following conditions: *In vitro* study, related to goat products consumption and neural developmental effect. There were no restrictions in the study design and publication date. The adapted data extraction form (Appendix A) include information about the authors, selection procedure, analyses, animals' characteristics, type of intervention(s), characteristics of methodological quality, and results.

| Heading               | Subheading                  | Description  | For completion by reviewer(s) |     |    |         |
|-----------------------|-----------------------------|--|-------------------------------|-----|----|---------|
| Bibliographic details | Title                       |  |                               |     |    |         |
|                       | Authors                     |  |                               |     |    |         |
|                       | Publication date            |  |                               |     |    |         |
|                       | Study aims                  | What were the study's aims and purpose?  |                               |     |    |         |
| Ethical standards     |                             | Was ethical committee approval obtained? Was informed consent obtained? Does the study address ethical issues adequately? Has confidentiality been maintained?   | Ethical approval              | Yes | No | Unclear |
|                       |                             |  | Informed consent              | Yes | No | Unclear |
|                       |                             |  | Ethical issues addressed      | Yes | No | Unclear |
|                       |                             |  | Confidentiality maintained    | Yes | No | Unclear |
| Context               | Aims                        | Are the aims and purpose of the study clearly stated?  |                               | Yes | No | Unclear |
| Setting               | Area and care setting       | What is the geographical and care setting for the study?   |                               |     |    |         |
|                       | Rationale                   | What is the rationale and appropriateness for this choice?   |                               |     |    |         |
|                       | Detail                      | Is there sufficient detail about setting?  |                               | Yes | No | Unclear |
|                       | Timing                      | Over what period did the data collection take place?   |                               |     |    |         |
| Sample                | Inclusion criteria          | Who was included in the study?   |                               |     |    |         |
|                       | Exclusion criteria          | Who was excluded in the study?   |                               |     |    |         |
|                       | Selection                   | How was the sample selected? Were there any factors that influenced how the sample was selected (e.g. access, timescale issues)?   |                               |     |    |         |
|                       | Size                        | What is the size of the sample and groups comprising the study?  |                               |     |    |         |
|                       | Appropriateness             | Is the sample appropriate in terms of its ability to meet the aims of the study, the depth of data that it is enables to be collected, and its breadth?  |                               | Yes | No | Unclear |
| Data collection       | Methods                     | What data collection methods were used?  |                               |     |    |         |
|                       |                             | Was the data adequately described and rigorously conducted?  |                               | Yes | No | Unclear |
|                       | Fieldwork                   | Is the process of fieldwork adequately described?  |                               | Yes | No | Unclear |
|                       | Data analysis               | How are the data analyzed? How adequate is the description of the data analysis? Is adequate evidence provided to support the analysis (e.g. use of original data, iterative analysis, efforts to establish validity and reliability)? Is the study set in context in terms of findings and relevant theory? |                               |     |    |         |
|                       | Researcher's potential bias | Are the researcher's /researchers' own position, assumptions and possible biases outlined? Indicate how they could affect the study in terms of analysis and interpretation of the data  |                               |     |    |         |
|                       | Reflexivity                 | Are the findings substantiated by the data and has consideration been given to any limitations of the methods or data that may have affected the results?  |                               | Yes | No | Unclear |
| Outcomes              | Outcomes                    | What outcome measures were adopted? What was the impact of the study for (a) service users (b) careers (c) practitioners (d) organization responsible for service  |                               |     |    |         |
| Findings              | Themes                      |  |                               |     |    |         |
|                       | Conclusion                  |  |                               |     |    |         |
|                       | Opinions                    | What this person argues?   |                               |     |    |         |

**Appendix A:** Data Extraction form (Supplementary data).

### Risk assessment


All the studies were screened to confirm that goat milk was given as supplementation as included in the methodology. Once the studies were confirmed to show the effectiveness of goat milk consumption on the central nervous system and physical activity, the studies were categorised as potential studies for further evaluation.

### Summary measurement

This study was conducted qualitatively to synthesise the effects of goat milk on neural development. Meta-analysis was not attempted due to the exploratory design of the study.

### Results

A flow diagram in Figure 1 has outlined the systematic review process of the research result selection and the reasons of exclusion. The search strategies initially identified 3185 abstracts from six databases. After duplicates abstracts were removed, there are 2702 abstracts remained. By screening the titles and abstracts, seven articles were selected for full-text review where among them, no human-study was retrieved and four of them are animal-related studies which focusing on physical and brain development and reflex maturation. One of the studies used goat milk gene promoter, hence it was excluded.



**Figure 1:** Flow diagram of study selection. Note: n = number of publications.

### Study characteristics

Three out of four studies that fit the criteria were conducted by the same group of researchers between the year of 2012 to 2014 [9,24,25]. The studies discussed the effect of conjugated linoleic acid of goat milk on the brain. Two of the studies compared goat milk fat (experimental; n = 83) and soybean oil (control; n = 27), while other studies compared dietary lipids from goat milk (n = 24), soybean (n = 18), and coconut oil (n = 12). Another similar study was conducted by Medeiros, *et al.* [26] where the study was conducted using dried goat whey. Table 1 summarises the findings of all four studies.

### Risk of bias across studies

All the studies were conducted in the same laboratory. The rat diet which exclude goat milk was not discussed thoroughly in the methodology. Therefore, it cannot be concluded that the goat milk supplementation was beneficial to the brain development.

### Study intervention

According to Soares, *et al.* [9,24,25], female Wistar rats that had been mated to obtain a few progenies for further analysis were used in the three studies. Each diet differs in the composition of the saturated-to-polyunsaturated fatty acid ratio [9,24,25]. The experimental group received a diet containing 7% fat from goat milk. In this study, the experimental rats were divided into three groups. The first group (n = 27) of rats were fed with goat milk during the gestation phase. The second group (n = 28) of rats were fed with goat milk during the lactation phase, and the third group (n = 28) of rats were fed with goat milk from gestation phase up to 35 to 45 days of lactation phase.

Medeiros *et al.* [26] tested only male pups in their study. This study focused on the effects of sialic acid (SA) content in goat whey on memory and cortical spreading depression (CSD). In this study, the experimental pups were divided into two major groups; litters with nine (L<sub>9</sub>) (n = 51) and fifteen (L<sub>15</sub>) (n = 52) pups, respectively. There are five subgroups in each group; dried goat whey (DGW), SA20 (20 mg/kg/day), SA100 (100 mg/kg/day), saline solution and naïve groups. The pups administered DGW and SA via gavage. No gavage was introduced to control groups; saline and naïve groups. The DGW received 17.45 g/kg/day dried goat whey (DGW), which equivalent to 20 mg of SA. The whole experiment required samples size of 51 L<sub>9</sub> and 51 L<sub>15</sub>.

| Author and Year               | Milk Composition      | Animals   | Experimental design   | Parameters  | Result  |
|-------------------------------|-----------------------|---|---|---|---|
| Soares, <i>et al.</i> 2012    | 7% fat from goat milk | Female Wistar rats<br>Aged 120-150 days of life<br>Weighing 300 ± 50 g were mated | 1 control group (soybean oil); n = 27<br>3 experimental groups (goat milk); n = 83    | Body weight<br>Cortical spreading depression (CSD) recording  | Body weight increased in <i>lactation</i> and <i>gestation+lactation</i> groups (p < 0.05)<br>Increase CSD phenomenon (p < 0.05)  |
| Soares, <i>et al.</i> 2013    | 7% fat from goat milk | Female Wistar rats<br>Aged 120-150 days of life<br>Weighing 250 ± 50 g were mated | 1 control group (soybean oil); n = 27<br>3 experimental groups (goat milk); n = 83    | Body weight<br>Anxiety behaviour<br>Reflex responses  | Pups in all groups showed higher body weight<br><i>Gestation</i> group indicated less anxiety than control group (p < 0.05)<br>In the <i>lactation</i> group, cliff avoidance and free-fall righting responses were respectively delayed and accelerated (p < 0.05) |
| Soares, <i>et al.</i> 2014    | 7% fat from goat milk | Female Wistar rats<br>Aged 120-150 days of life<br>Weighing 250 ± 50 g were mated | Soybean oil group; n = 18<br>Coconut oil group; n = 12<br>Goat milk fat group; n = 24 | Body weight<br>Reflex responses   | Lipids from goat milk increase body weight in all groups<br>Acceleration of reflex maturation (p < 0.05)  |
| Me-deiros, <i>et al.</i> 2016 |                       | Male pups Wistar rats   | 20 mg DGW; n = 19<br>SA20; n = 21<br>SA100; n = 20<br>Nv; n = 21<br>Sal; n = 20       | Body weight<br>Recognition task (Spatial position, shape and novel object)<br>Cortical spreading depression (CSD) recording | DGW, SA20 and SA100 showed increased in body weight (p < 0.05)<br>Longer exploration time was presented in DGW group (p < 0.05)<br>Increased CSD propagation velocities (p < 0.05)  |

**Table 1:** Summary of included studies (in chronological order).

## Outcomes

All studies examined the effect of fat (conjugated linoleic acid) and sialic acid contained in goat milk and dried goat whey, respectively, based on cortical activity. The effects of diet on reflex maturation, memory performance and anxiety behaviour were also discussed [9,24-26].

## Reflex response

The pup's reflexes ontogeny (maturation process) was investigated based on palm grasp, righting, free-fall righting, vibrissae placing, auditory startle response, negative geotaxis and cliff avoidance. Goat milk supplementation improved reflexes response and somatic maturation in pups. It was reported that cliff avoidance and free-fall righting responses were respectively delayed and accelerated in all three experimental groups especially in the lactation group [9,24].

## Cortical activity

The study by Soares, *et al.* [24] focused on the influence of the dietary lipids on neuronal excitability. The investigation was conducted using the electrophysiological activity known as cortical spreading depression (CSD). In both gestation and combined gestation-lactation groups of goat milk fat consumption, there is a significantly higher in number of multiple episodes occurred in these groups. Moreover, the propagation of CSD velocities in both lactation and combined gestation-lactation group were higher than the control and gestation groups. This is in line with a study where the experimental groups show a higher propagation of CSD velocities compared to the control groups [26]. Leo [27] had described that CSD is a fully reversible and excitability-related neural response. It is a slow propagation wave of depression of spontaneous neuronal activity from electrical, mechanical or chemical stimulation of one point on brain tissue, from which it spreads concentrically to remote cortical regions.

### Anxiety behavior

Elevated plus maze (EPM) was used as a model for testing the anxiety behaviour of pups [9]. This test was used to characterise the exploratory and anxiolytic effects based on the recorded time spent in the maze. The total time spent in the open and closed arms as well as in the central square were recorded and the number of head dipping was counted. The anxiety level was reduced in dams and pups supplemented with goat milk and the behavioural effect was minimised. The duration of time that the pups remain stationary at the open arm of the EPM was increased associated with the increased in the number of head dipping. However, the pups tend to spend less time at the centre of the maze.

### Memory and behavioural test

A specific test named object recognition test (ORT) is a behavioural assay that is commonly used in order to investigate animal's learning and memory. This test helps to evaluate the efficacy of an animal to recall identical objects after a new object was switched with one of the objects in the apparatus. The rat will spend more time on the new object except for the rat with a memory deficit. Longer exploration time was taken in DGW groups.

### Discussion

The systematic review was conducted to study the effect of goat milk on neural development and functions. The result revealed that three studies by the same author recorded the benefits of fatty acid in goat milk, which enhance brain development and functions [9,24,25]. The main results from all three studies showed an increase in body weight, and improvement in reflex response and brain development after a period of goat milk fat consumption. In addition, anxiety-like behaviour with an anxiolytic effect and cortical activity were also discussed. These experimental studies were conducted on rat dam and its progeny.

The fatty acids composition of both experimental and control diets revealed a different ratio between saturated and polyunsaturated fatty acids. The experimental diet of goat milk recorded CLA in fatty acids composition, whereas no CLA was detected in the standard diet [9,24,25]. Based on all three studies that have been conducted by Soares, *et al.* [9,24,25], chronologically from the year 2012 to 2014, CLA is crucial for the CNS development, especially during pregnancy and lactation period.

Another study revealed the benefit of sialic acid on pups memory after the consumption of dried goat whey [25]. Goat whey, which that rich is sialic acid is a by-product that was discarded after goat cheese manufacturing. The experimental groups showed a positive result in object recognition test (ORT). The rats took a longer time to explore the new object among old objects that were placed in the apparatus concluding that sialic acid is a good source for memory and brain excitability processes.

### Reflex response

The consumption of goat milk fat shows an acceleration in reflex maturation compared to control group, such as soybean oil and/or coconut oil [24,25]. CLA content in goat milk lipid might be responsible for the change in the acceleration and the reflex maturation of the pups. These findings were consistent with the previous studies, where rats fed with diet supplementation showed an acceleration of reflexes [14]. These findings demonstrate that the changes in maternal dietary fatty acids may facilitate the reflex ontogeny in the offspring.

During the postnatal period, free-fall righting responses can only be observed once the central nervous system has been fully developed [28]. Besides, vestibulospinal, visual and somesthetic afferents are also directly related to posture correction [29]. There are a few developmental events that are able to generate such reflex responses. The events include neuronal and glial proliferation and the establishment of synaptic contacts [30], as well as myelination [18], which are directly related to dietary lipids. Goat milk fat diet consumption showed that pups are able to turn in mid-air to land on its four paws [23,24]. Therefore, during the gestation period, the accumulation of fatty acids in the brain is crucial for fetal development [10]. Hence, the accumulation of fatty acids is very important during the rapid development of nervous system which was seen in the ontogeny reflex [31].

### Cortical activity

Leo [27] discovered that the initial phase of cortical spreading depression (CSD) was characterised by the depression of the spontaneous brain electrical activity and the electroencephalography depression spread slowly all over the brain cortical surface. The increased in CSD is associated with the pathophysiological events of ischemia, migraine and epilepsy in human [32,33]. By referring to the previous work published, the measurement of the physi-

cal parameters required an anaesthesia [25,26]. Therefore, at this stage, no changes were seen by the physical parameters in the experimental groups. In contrast, the CSD propagation in the experimental groups showed an activity in brain electrophysiological even though an absence of weight difference. Thus, this suggested that the CLA and sialic acid effects on body development were short lasting compared to CSD propagation, where long-lasting effects on brain electrophysiological were observed in pups growth.

The CLA would impair brain myelination, a development process in mammals to reach the maximal intensity during the lactation period. Modification in dietary lipids has been proven to affect on the myelination. A study reported that CSD propagations either accelerated or decelerated, were associated with demyelination and hypermyelination, respectively [34]. According to Borba, *et al.* [35] showed that the dam and progeny which received diet rich in saturated fatty acid caused an increased in myelination and decelerated the CSD. In contrast with Soares, *et al.* [24] and Medeiros *et al.* [26], an acceleration of CSD may be due to the diet used which probably impaired myelin synthesis in the progeny. It is a fact that CSD plays an important role in human neurological disorders like epilepsy [32]. An increase of epilepsy prevalence was noted among patients who have been diagnosed with multiple sclerosis [36] and demyelinating disease [37,38]. Moreover, the key enzyme that cleaves SA, known as neuraminidase (NEU), has been proven to have better anticonvulsant action [39]. This was in line with the study conducted by Isaeva, *et al.* [39,40]. Still, further experiment on brain myelin needs to be quantified to show the relationship of this hypothesis.

In the study by Soares, *et al.* [24], the *gestation* and the combined *gestation-lactation* groups of rats consuming fat from goat milk showed a significantly high multiple episodes of CSD. Normally, once potassium chloride (KCl) was stimulated, a single CSD episode will be elicited, but in this study, "multiple" CSD episodes occurred with a single potassium chloride (KCl) stimulation. As supported by Bourre [31], the brain that is susceptible to the CSD phenomenon could be observed from these "multiple" CSD episodes. This result showed that goat milk may increase CSD phenomenon, which is an adverse reaction of goat milk. However, the reason for the increased of CSD in their study may be due to the diet of goat milk that is deficient in both n-3 and n-6 fatty acid, which is important for an adequate brain development [15]. Moreover, the goat milk has high PUFA content, which protects the brain from damage by inducing robust dilatation of the cerebral vessels [41].

### Anxiety behaviour

Anxiety disorder can interfere in the learning process by impairing attention and memory. Thus, to measure the anxiety level or to characterise an exploratory and anxiolytic effect, elevated plus maze apparatus has been used. The elevated plus maze (EPM) has been designed to accomplish the anxiety test conducted on animals where the main target was to measure the time taken for an animal to be in different position.

Pups that consumed goat milk fat exhibited an increase and decrease of the time spent in the open arm and centre part of the maze, respectively, which were significantly different as compared to the control group. Head dipping recorded was also increased recorded by the goat milk fat consumed group. According to Soares, *et al.* [9], goat milk fat consumption especially n-3 fatty acid is associated with the decreased of anxiety-like behaviour in the plus maze test. Once the anxiety becomes pathological, it will impair the attention and memory which then disturbed the learning processed [42]. Similarly, this study discusses the work presented by Sharma, *et al.* [43] and Sivanathan, *et al.* [44], where the anxiety behaviour was also reduced when the rat consumed hypercaloric diet. This reinforces the idea that the changes in anxiety behaviour in the progeny were mainly caused by the consumption of goat milk fat which consequently affects the pups' nervous system. This is not due to the dams' nutritional deficiency. It is suggested that the CLA has been postulated to contribute a positive effect on the anxiety behaviour (less anxiety) as compared to the control group.

### Memory and behavioural test

Memory and learning impairment may interfere with the spatial cognitive ability. In order to measure the spatial cognitive ability, object recognition test (ORT) was applied using a circular arena apparatus with a diameter of 89 cm. This test was conducted on animals, where the main objective was to measure the time taken for an animal to recognise the novel object among other objects that had been placed in the same apparatus.

Pups that consumed dried goat whey (DGW) required more time during the recognition of the new object showing a significant difference as compared to the control group. Once memory impairment becomes pathological, it will impair the behavioural and electrophysiological actions. According to Medeiros, *et al.* [26], DGW contains sialic acid (SA) received by gavage, associated with the improvement of object recognition memory of an animal. Similarly,

this study discusses the work presented by Limon, *et al.* [45] and Wang, *et al.* [46], where memory and learning test were also demonstrated to perform better when the pups consumed with sialic acid. However, the effect was not caused only by sialic acid (SA), but other nutrients such as taurine [47,48] contained in DGW were necessary to support the memory and learning ability. This reinforces the idea that the changes in memory effects among the pups were mainly caused by the gavage of DGW, resulting in the direct action on the pups' memory and learning ability. It is suggested that the SA has been postulated to improve the spatial cognitive ability when compared to the control group.

The present study has several limitations. Although the search strategies are not limited to human and animal studies, there was lack of evidence on the effect of goat milk on brain activity. This may also due to the collected data from unpublished papers that were not indexed in the databases and those that are unable to be retrieved. However, the limited findings suggest a relationship between the goat milk fat consumption by the maternal rats with the physical development, cortical activity, and anxiety behaviour in the progeny. Due to the small number of studies on goat milk consumption related to brain activity, future research is required to provide the evidence that goat milk consumption could affect brain activity in the rat. The data currently gathered was limited because the work was mainly conducted by the same authors. The papers did not discuss clearly the methodology of the experimental studies, especially regarding the rat's diet. Thus, the information whether the dams and the progeny were given any other food was not clearly stated. To date, the recorded goat milk effects were all positive with respect to its body development (improvement in physical growth), the central nervous system (reduction in anxiety behaviour), and cortical activity (susceptible to the CSD phenomenon).

## Conclusion

In conclusion, there is a need to provide more evidence on the effects of goat milk consumption on brain activity among human and animal. More research should be conducted to strengthen the relationship between goat milk consumption and brain activity.

## Acknowledgment

This study is sponsored by Ministry of Higher Education Malaysia under the NRGs grant USIM/NRGs\_P8/FPSK/8404/52113

## Conflict of Interest

The authors declare that there are no conflicts of interest.

## Bibliography

1. Hodgkinson AJ, *et al.* "Allergic Responses Induced by Goat Milk As1-Casein in a Murine Model of Gastrointestinal Atopy". *Journal of Dairy Science* 95.1 (2012): 83-90.
2. Fiocchi Alessandro, *et al.* "World Allergy Organization (Wao) Diagnosis and Rationale for Action against Cow's Milk Allergy (Dracma) Guidelines". *World Allergy Organization Journal* 3.4 (2010): 57.
3. Mack PB. "A Preliminary Nutrition Study of the Value of Goat's Milk in the Diet of Children". *Yearbook of the American Goat Society* 1953 (1952).
4. Park Young W, *et al.* "Bioavailability of Iron in Goat Milk Compared with Cow Milk Fed to Anemic Rats 1, 2". *Journal of Dairy Science* 69.10 (1986): 2608-2615.
5. Ceballos Laura Sanz, *et al.* "Composition of Goat and Cow Milk Produced under Similar Conditions and Analyzed by Identical Methodology". *Journal of Food Composition and Analysis* 22.4 (2009): 322-29.
6. Raynal-Ljutovac K, *et al.* "Composition of Goat and Sheep Milk Products: An Update". *Small Ruminant Research* 79.1 (2008): 57-72.
7. Sampelayo MR Sanz, *et al.* "Influence of Type of Diet on the Fat Constituents of Goat and Sheep Milk". *Small Ruminant Research* 68.1 (2007): 42-63.
8. Barłowska J. "Nutritional and Pro-Health Properties of Milk Fat". *Medycyna Weterynaryjna* 65.3 (2009): 171-174.
9. Soares Juliana KB, *et al.* "Conjugated Linoleic Acid in the Maternal Diet Differentially Enhances Growth and Cortical Spreading Depression in the Rat Progeny". *Biochimica et Biophysica Acta (BBA)-General Subjects* 1820.10 (2012): 1490-1495.
10. Lenzi Almeida, *et al.* "Influence of Omega-3 Fatty Acids from the Flaxseed (*Linum Usitatissimum*) on the Brain Development of Newborn Rats". *Nutricion Hospitalaria* 26.5 (2011).

11. Bautista Claudia J., *et al.* "Protein Restriction in the Rat Negatively Impacts Long-Chain Polyunsaturated Fatty Acid Composition and Mammary Gland Development at the End of Gestation". *Archives of Medical Research* 44.6 (2013): 429-436.
12. Innis Sheila M. "Impact of Maternal Diet on Human Milk Composition and Neurological Development of Infants". *The American Journal of Clinical Nutrition* 99.3 (2014): 734S-741S.
13. Hadley Kevin B., *et al.* "The Essentiality of Arachidonic Acid in Infant Development". *Nutrients* 8.4 (2016): 216.
14. Santillán, María E, *et al.* "Developmental and Neurobehavioral Effects of Perinatal Exposure to Diets with Different  $\Omega$ -6:  $\Omega$ -3 Ratios in Mice". *Nutrition* 26.4 (2010): 423-431.
15. Schmitz Gerd and Josef Ecker. "The Opposing Effects of N-3 and N-6 Fatty Acids". *Progress in Lipid Research* 47.2 (2008): 147-155.
16. Zhu Haiyan., *et al.* "Dietary Fish Oil N-3 Polyunsaturated Fatty Acids and Alpha-Linolenic Acid Differently Affect Brain Accretion of Docosahexaenoic Acid and Expression of Desaturases and Sterol Regulatory Element-Binding Protein 1 in Mice". *The Journal of Nutritional Biochemistry* 21.10 (2010): 954-960.
17. Park Yeonhwa. "Conjugated Linoleic Acid (Cla): Good or Bad Trans Fat?". *Journal of Food Composition and Analysis* 22 (2009): S4-S12.
18. Nochera Carmen L., *et al.* "Consumption of Dha+ Epa by Low-Income Women During Pregnancy and Lactation". *Nutrition in Clinical Practice* 26.4 (2011): 445-450.
19. Uauy, Ricardo., *et al.* "Essential Fatty Acids in Visual and Brain Development". *Lipids* 36.9 (2001): 885-895.
20. Bazinet Richard P and Sophie Layé. "Polyunsaturated Fatty Acids and Their Metabolites in Brain Function and Disease". *Nature Reviews Neuroscience* (2014).
21. Igarashi., *et al.* "Impact of Maternal N-3 Polyunsaturated Fatty Acid Deficiency on Dendritic Arbor Morphology and Connectivity of Developing *Xenopus Laevis* Central Neurons in Vivo". *The Journal of Neuroscience* 35.15 (2015): 6079-6092.
22. Park Young W. "2.2 Goat Milk - Chemistry and Nutrition". *Handbook of Milk of Non-Bovine Mammals* (2008): 34.
23. Fox PF, *et al.* "Chemistry and Biochemistry of Fermented Milk Products". *Dairy Chemistry and Biochemistry*. Springer, (2015): 547-67.
24. Soares Juliana KB., *et al.* "Anxiety Behavior Is Reduced, and Physical Growth Is Improved in the Progeny of Rat Dams That Consumed Lipids from Goat Milk: An Elevated Plus Maze Analysis". *Neuroscience Letters* 552 (2013): 25-29.
25. Soares Juliana Kessia Barbosa., *et al.* "Acceleration of Reflex Maturation and Physical Development in Suckling Rats: Effects of a Maternal Diet Containing Lipids from Goat Milk". *Nutritional Neuroscience* 17.1 (2014): 1-6.
26. Medeiros Larissa de Brito., *et al.* "Neonatal Administration of Goat Whey Modulates Memory and Cortical Spreading Depression in Rats Previously Suckled under Different Litter Sizes: Possible Role of Sialic Acid". *Nutritional Neuroscience* (2016).
27. Leo Aristides AP. "Spreading Depression of Activity in the Cerebral Cortex". *Journal of Neurophysiology* 7.6 (1944): 359-390.
28. Falcão-Tebas Filipe., *et al.* "Maternal Low-Protein Diet-Induced Delayed Reflex Ontogeny Is Attenuated by Moderate Physical Training During Gestation in Rats". *British Journal of Nutrition* 107.03 (2012): 372-377.
29. Boyle R. "Vestibulospinal Control of Reflex and Voluntary Head Movement". *Annals of the New York Academy of Sciences* 942.1 (2001): 364-80.
30. Altman J and SA Bayer. "An Overview of the Postnatal Development of the Rat Cerebellum". *Development of the cerebellar system: in relation to its evolution, structure, and functions* (1997): 324-333.
31. Bourre Jean-Marie. "Effects of Nutrients (in Food) on the Structure and Function of the Nervous System: Update on Dietary Requirements for Brain. Part 2: Macronutrients". *Journal of Nutrition Health and Aging* 10.5 (2006): 386.
32. Rogawski Michael A. "Common Pathophysiologic Mechanisms in Migraine and Epilepsy". *Archives of Neurology* 65.6 (2008): 709-714.
33. Nozari Ala., *et al.* "Microemboli May Link Spreading Depression, Migraine Aura, and Patent Foramen Ovale". *Annals of Neurology* 67.2 (2010): 221-229.

34. Lehmann Michael L., *et al.* "Chronic Social Defeat Reduces Myelination in the Mouse Medial Prefrontal Cortex". *Scientific reports* 7 (2017): 46548.
35. Borba., *et al.* "Essential Fatty Acid Deficiency Reduces Cortical Spreading Depression Propagation in Rats: A Two-Generation Study". *Nutritional Neuroscience* 13.3 (2010): 144-150.
36. Poser Charles M and Vesna V Brinar. "Epilepsy and Multiple Sclerosis". *Epilepsy and Behavior* 4.1 (2003): 6-12.
37. Odier Céline., *et al.* "Central Pontine and Extrapontine Myelinolysis: From Epileptic and Other Manifestations to Cognitive Prognosis". *Journal of Neurology* 257.7 (2010): 1176-1180.
38. Mao Ling-Yan., *et al.* "Interictal Interleukin-17a Levels Are Elevated and Correlate with Seizure Severity of Epilepsy Patients". *Epilepsia* 54.9 (2013).
39. Isaeva Elena., *et al.* "Effect of Neuraminidase Treatment on Persistent Epileptiform Activity in the Rat Hippocampus". *Pharmacological Reports* 63.3 (2011): 840-844.
40. Isaev Dmytro., *et al.* "Neuroaminidase Reduces Interictal Spikes in a Rat Temporal Lobe Epilepsy Model". *Epilepsia* 52.3 (2011).
41. Blondeau N., *et al.* "Polyunsaturated Fatty Acids Induce Ischemic and Epileptic Tolerance". *Neuroscience* 109.2 (2002): 231-241.
42. Françolin-Silva AL and SS Almeida. "The Interaction of Housing Condition and Acute Immobilization Stress on the Elevated Plus-Maze Behaviors of Protein-Malnourished Rats". *Brazilian Journal of Medical and Biological Research* 37.7 (2004): 1035-1042.
43. Sharma S., *et al.* "Adaptations in Brain Reward Circuitry Underlie Palatable Food Cravings and Anxiety Induced by High-Fat Diet Withdrawal". *International Journal of Obesity* 37.9 (2013): 1183.
44. Sivanathan Shathveekan., *et al.* "Chronic High Fat Feeding Increases Anxiety-Like Behaviour and Reduces Transcript Abundance of Glucocorticoid Signalling Genes in the Hippocampus of Female Rats". *Behavioural Brain Research* 286 (2015): 265-270.
45. Limón Ilhuicamina Daniel., *et al.* "Alteration of the Sialylation Pattern and Memory Deficits by Injection of A $\beta$  (25-35) into the Hippocampus of Rats". *Neuroscience Letters* 495.1 (2011): 11-16.
46. Wang Bing. "Molecular Mechanism Underlying Sialic Acid as an Essential Nutrient for Brain Development and Cognition-3". *Advances in Nutrition* 3.3 (2012): 465S-72S.
47. Rutherford Shane M., *et al.* "Amino Acid Composition Determined Using Multiple Hydrolysis Times for Three Goat Milk Formulations". *International Journal of Food Sciences and Nutrition* 59.7-8 (2008): 679-690.
48. Wenting Lu., *et al.* "Therapeutic Effect of Taurine against Aluminum-Induced Impairment on Learning, Memory and Brain Neurotransmitters in Rats". *Neurological Sciences* 35.10 (2014): 1579-1584.

### Volume 3 Issue 3 March 2018

© All rights are reserved by Nor Azila Noh., *et al.*