

Physical Activity Level and Prevalence of Obesity among Adults' Population of Abeokuta, Ogun State, Nigeria

AS Adekoya^{1,2,3*}, CI Ojo³, OA Mafimidiwo^{3,4} and FO Oyeleke⁵

¹SAM Obesity Consult, Nigeria

²Marie Curie Academy, Abeokuta, Nigeria

³Federal Cooperative College, Ibadan, Nigeria

⁴Department of Human Nutrition, University of Ibadan, Nigeria

⁵Department of Human Nutrition and Dietetics, Kwara State College of Health

Technology, Offa, Nigeria

*Corresponding Author: AS Adekoya, SAM Obesity Consult, Nigeria.

Received: March 31, 2020

Published: June 22, 2020

© All rights are reserved by AS Adekoya,
et al.

Abstract

Objective: To examine the effect of physical activity level (PAL) on the prevalence of obesity among adults (20 - 64 yr) in Abeokuta, Ogun State, Nigeria.

Designs: 120 adult males and 120 adult females were randomly selected to participate in the study. Body mass index (BMI) was measured using cut-off $\geq 30 \text{ kg/m}^2$ and the physical activity assessed according to WHO/FAO work classification.

Results: Mean BMI was high above the cut-off (7.2% for rural male and 11% for urban male, $p = 0.169$) and (28.3% for urban female but lower than the cut-off among rural female 7.7%, $p = 0.006$). The pattern of physical activity significantly fell on the sedentary side, mostly among the urban subjects. There were positive changes (statistically insignificant) in physical activity level (PAL) and a similar shift in obesity prevalence among the sub-population.

Conclusion: The PAL among the population, emphasize the adverse effect of sedentarism on the increasing risks of obesity and chronic degenerative diseases. Participation in physical activity programs should be emphasized.

Keywords: PAL; BMI; Adults; Obesity

Introduction

Physical activity plays a critical role in obesity development, because urban women are more sedentary than are their rural counterparts [11]. Sedentary lifestyles are increasing influencing the prevalence of obesity [11,17]. The process modern transport, labour saving devices, TV viewing etc. has brought about a number of consequences affecting physical activity patterns that contribute to obesity [4,8,30].

Over the years, large number of studies evaluating westernized dietary changes in various population in developing countries have been carried out, but little is known about the physical activity levels (PAL) in these populations [1,3,6,14-16]; which has been said to be due to difficulties in measuring total daily energy expenditure (TEE) in free-living situations [4,5,15,25,44].

PAL is needed to quantitatively compare the patterns of physical activity between urban and rural dwellers who shared the same genetic traits and cultural background (often seen in rural-urban migrants and rural residents of the same group), which should be

useful for determining the effect of urbanization [22-24,40,41,43]. The PAL is useful as it allows for comparison of individuals of different body size. PAL values are a universally accepted expression of energy expenditure and help to convey easy understanding of the concept of physical activity pattern [6,27,28,31-33,35]. In this study, the effect of the physical activity level on the prevalence of obesity in both the rural and urban adults (20 - 64yr) in Abeokuta was examined.

Materials and Methods

In all, 240 adults (male = 120, female =120) representative sample within the age 20 years and 64 years were randomly and systematically selected from the enumerated households in both the rural and urban areas of Abeokuta (which is a city in Ogun State, western part of Nigeria in the western region of Sub-Saharan Africa). The inhabitants of the urban area are mostly elites and civil servants, living in modern structured residential buildings and settings. The samples were the participants that were measured and assessed, which represented the population of the localities that were selected for the study.

Validated questionnaires were used to record detailed information. The heights and weights of the subjects were taken to compute the BMI (kg/m^2) adopting the World Health Organization (2000) classification of body weight in adults: normal weight 18.5 - 24.9; Overweight 25.0 - 29.9; obese ≥ 30.0 . The height measurements were obtained using calibrated standing rule and measured to the nearest 0.1 cm. The weight measurements were performed on the same digital scale, which was periodically checked for precision with known weights; and measured to the nearest 0.1kg. The data were taken in the morning, and all the subjects wore light clothing without shoes and socks; with their hands resting on their sides.

Physical activity patterns were based on the mode of transportation, number of hours devoted to sleep, type and nature of occupation, and recreational activities within the previous 24 hours. The duration of each activity was multiplied by its energy cost and summed up; using energy cost of activities as given by FAO/WHO/UNU [5,18,21,26]:

$$\text{TEE} = (t_1 \text{AEE}_1) + (t_2 \text{AEE}_2) + (t_3 \text{AEE}_3) + (\dots t_n \text{AEE}_n)$$

Where t_1, t_2, t_n = time (min)

$\text{AEE}_1, \text{AEE}_2, \dots, \text{AEE}_n$ = Activity Energy Expenditure

$\text{PAL} = \text{TEE}/\text{BMR}$

$$= [(\sum \text{SleepingEE}) + (\sum \text{sedentaryEE}) + (\sum \text{moderateEE}) + (\sum \text{activeEE})]/\text{BMR} [22,34-36,48]$$

Sleeping EE = BMR = $2\text{MJ}/8\text{hr}$ (male), $1.75\text{MJ}/8\text{hr}$ (female) [7,15,24,25,27].

Statistical analysis was performed with t-tests were used for comparison between the urban and rural subjects. Differences were considered to be significant at $p < 0.05$. Results were expressed as means \pm S.D and percentages [37,38,45,47].

Results

In the analysis of patterns of distribution of the anthropometric and physical activity characteristics of the subjects, according to their sex groups and localities/areas, using statistical computing shown in table 1 [37,38,45]. The mean values of body weights and heights were significantly moderate; but are higher among the urban subjects than the rural subjects except in the height of the rural females.

The mean BMI was also higher among the urban subjects than the rural subjects in both sex groups, which may be said to be associated with changes in lifestyles of the urban subjects as observed in this study. Mean BMI was higher above the cut-off among the

rural male by 7.2% ($p = 0.169$), urban male by 11% ($p = 0.169$), urban female by 28.3% ($p = 0.006$), but fell among the rural female by 7.7% ($p = 0.006$).

The pattern of physical activity fell on the sedentary side significantly among the urban subjects than the rural subjects, as seen in the mean total energy expenditure. The TEE is lower significantly among the urban male (2191 ± 410) as compared to the rural male (2639 ± 344) by 17% ($p < 0.005$), and among the urban female (2092 ± 580) as compared to the rural female (2310 ± 530) by 9.4% ($p < 0.005$).

The main focus of this report is to evaluate the PAL value of the sampled population and compared with other findings, such as those described in [6,24,33]. The findings of this study compared very well with those described by [6] for developing countries (i.e. 1.85 ± 0.31 for males and 1.65 ± 0.17 for females, and that 1.8 was the threshold to trigger obesity). There were large, insignificant, positive changes in the PAL values of the urban subjects as compared to the rural subjects. The values as compared to the threshold, showed the rural male (RM) with +3.9% ($p < 0.005$), urban male (UM) with -2.8% ($p < 0.005$), rural female (RF) with +2.8% ($p < 0.005$) and urban female (UF) with -6.7% ($p < 0.005$) (Table 1).

Variables	Rural (n 60)	Urban (n 60)	p-value
Males (n 120)			
Body weight (kg)	75.6 ± 8.11	773 ± 7.63	< 0.001
Height (cm)	172.1 ± 8.5	179.6 ± 3.4	< 0.001
BMI (kg/m^2)	32.16 ± 3.63	33.30 ± 4.69	0.169
Total Energy Expenditure (kcal/day)	2639 ± 344	2191 ± 410	< 0.050
PAL (Physical Activity level)	1.87 ± 0.31	1.75 ± 0.24	< 0.050
% Obese (BMI $30 \text{ kg}/\text{m}^2$)	36.7	51.9	
% walking less than 1 hr/day	42	57	
% not participating in any PA or Sports	30	60	
Females (n 120)			
Body weight (kg)	81.5 ± 9.60	89.8 ± 17.5	< 0.001
Height (cm)	169.3 ± 5.2	163.8 ± 6.5	< 0.001
BMI (kg/m^2)	27.69 ± 6.01	38.49 ± 5.80	0.006
Total Energy Expenditure (kcal/day)	2310 ± 530	2092 ± 580	< 0.050
PAL (Physical Activity Level)	1.85 ± 0.49	1.80 ± 0.83	< 0.050
% Obese (BMI $30 \text{ kg}/\text{m}^2$)	53.3	62.3	
% walking less than 1 hr/day	52	72	
% not participating in any PA or Sports	38	65	

Table 1: Anthropometry, physical activity characteristics and % obesity of the subjects (n = 240) in the rural and urban areas.

Values are given as mean \pm SD and percentage values within the sex group are compared with paired t-tests.

In addition, the relationship between the physical activity level and obesity was most evident in the prevalence, which was 36.7% among the rural male (RM), 51.9% among the urban male (UM), 53.3% among the rural female (RF) and 62.3% among the urban female (UF) (Table 1) and shown in figure 1 and 2 and when compared with those obtained globally in some other studies, they showed some similarities in their values and distributions around their percentages [32,33,40,41].

Cross-sectional studies have shown that increased acquisition of cars, labour saving household devices, TV, viewing etc. can make people to be walking less than 1 hour per day or making people not to be physically active or engage in any sports (Table 1); which have a number of consequences affecting physical activity patterns that are increasingly involved in the prevalence of obesity [1,8,17,30,32,40]. These studies also confirmed that physical activity pattern is most strongly associated with increasing prevalence of obesity in the world [2,10,11,19,30,32,40]. As can be seen from this finding and the figures 1 and 2 presented, that as PAL increases, the percentage prevalence decreases, and vice versa. Whereas some studies did not support that physical activity as reported in the freely living adult population in the long term is associated with the development of obesity, but the studies indicate that obesity may lead to physical inactivity [9,12,14,25,30,44]. There are some studies showing no association [3,13,20].

It has been well known for many decades that physical activity (PA) and exercise provide substantial benefits for the prevention of obesity, as well as overall health, functional capability, quality of life and longevity [23-26,28].

Although much progress has been made in this field, existing longitudinal epidemiological studies have found that PA levels are negatively correlated with weight gain. While some findings also relate the physical activity level (PAL) and body weight to the risk of developing cardiovascular diseases and coronary artery diseases in young adults [27-29]. Increasing PA levels from low to moderate ensures maintenance of body weight, while increasing PA levels from low to high helps individuals lose weight and reduced the risk of contracting CVD and CAD [30-32]. Showing and affirming that PAL would be a good predictor of the effects of incidence and prevalence of most of the obesity related diseases [2,28,29,42,46].

These findings, however, should be interpreted with some caution. Changes (persistence and non-persistence) in PA in this study were defined based on the levels of PA and the prevalence of obesity among the participants. In some cases, participants in the 'sedentary' group are not completely sedentary, and participants in the 'high' active group are not completely highly active. There are limitations in the result presented in the finding which could potentially underestimate the true effects of PA on obesity prevalence and the relative risk estimations [39,42,46]. One of them is the assessment of the correlation that exists between the PA and the percentage of the population that are obese. Future studies with more comprehensive longitudinal data analyses could provide data and evidence supporting the recommendation for the population to improve PAL and maintain this active lifestyle throughout their lifespan for the prevention of obesity, overweight and associated CVD [7,10,28,34].

Figure 1: PAL values in relation to the areas and genders.

Figure 2: % prevalence of obesity in relation to the areas and genders.

Discussion

The BMI distribution and the physical activity level were in agreement with those described in [6,24], showing that the people were transiting towards sedentary lifestyles. Age was an aspect which greatly affects the number of hours spent sitting down at work, in leisure time as well as time spent walking [11,25,27]. The results revealed that most males engage in one sport or the other, or in running around for businesses, while their female counterparts are mostly in sedentary state even while working, which may have contributed to the high prevalence of obesity experienced among the females than the males; thus confirming the difference in the prevalence of obesity among sexes [11,19,20,36,40,41].

Conclusion

The conclusions drawn from several recent reviews, although not fully adequate from the point of view of the problem discussed in this study, attest to the relevance of investigating whether various levels of physical activity are related to increasing prevalence of obesity. These results support the effect of physical activity level on the development and prevalence of obesity in the society. The study indicates that less physical activity leads to obesity and should encourage future studies into establishing the relationship between physical activity and obesity prevalence incorporating all other phenomena in the methodology.

Acknowledgments

This manuscript has been presented as a poster at the 18th International Congress of Nutrition, Durban, South Africa, in 2005. The author is responsible for the opinions expressed which were based on the results of findings of research thesis from the Federal University of Agriculture, Abeokuta, Nigeria. They are not attributed to the sponsors or the editors of this manuscript.

We thank the participants of this study for their contribution and technical assistance. We acknowledged the contributions of the supervisors, reviewers and all other researchers and colleagues towards the success of this article.

The authors appreciate statistical support and expertise from the Statistical Consulting Group. We also thank the anonymous reviewers for feedback to improve the review and approval of the manuscript.

Author Contributions

The authors conceived and designed the experiments, performed the experiments, analyzed the data, interpreted the data, wrote the first draft, and contributed to the final paper. All authors approved the final version. All authors are involved in the preparation of manuscript.

Funding

The author received no specific funding for this project, but acknowledges previous salary support from the Academics Works. The funds played role in the design and conduct of the study; collection, management, analysis and interpretation of the data; and preparation, review or approval of the manuscript.

Conflict of Interest

The authors declare that they have no conflicts of interest.

Bibliography

1. Bar OR., et al. "Physical activity, genetic and nutritional considerations in childhood weight management". *Medicine and Science in Sports and Exercise* 30 (1998): 2-10.
2. Bild DE., et al. "Correlates and predictors of weight loss in young adults: the CARDIA study". *International Journal of Obesity* 20 (1996): 47-55.
3. Crawford DA., et al. "Television viewing, physical activity and obesity". *International Journal of Obesity* 23 (1999): 437-440.
4. Durnin JVGA. "Low energy expenditure in free-living population". *European Journal of Clinical Nutrition* 44 (1990): 95-102.
5. FAO/WHO/UNU. "Energy and Protein requirements". Technical Report Series. Geneva, WHO (1985): 724.
6. Ferro-Luzzi A and Martino L. "Obesity and Physical activity". In: The origins and consequences of obesity. Ciba Foundation Symposium 201 (1996): 207-227.
7. Goldberg GR., et al. "Overweight and basal metabolic rates in men and women". *European Journal of Clinical Nutrition* 42 (1988): 137-144.
8. Goldblatt PB., et al. "Social Factors in obesity". *The Journal of the American Medical Association* 191 (1995): 1039-1044.
9. Haapanen N., et al. "Association between leisure time physical activity and 10-year body mass change among working-aged men and women". *International Journal of Obesity* 21 (1997): 288-296.
10. Klesges RC., et al. "A longitudinal analysis of the impact of dietary intake and physical activity on weight change in adults". *The American Journal of Clinical Nutrition* 55 (1992): 818-822.
11. Mokhtar N., et al. "Diet, culture and obesity in Northern Africa. Symp: Obesity in Developing Country". *American Society for Nutrition - Nutrition Research and Practice* 131 (2001): 887S-892S.
12. Owens JF., et al. "Can physical activity mitigate the effect of aging in middle-aged women?" *Circulation* 85 (1992): 1265-1270.
13. Parker DR., et al. "Dietary factors in relation to weight change among men and women from two southeastern New England communities". *International Journal of Obesity* 21 (1997): 103-109.
14. Petersen I., et al. "Longitudinal study of the long-term relation between physical activity and obesity in adults". *International Journal of Obesity* 28 (2004): 105-112.
15. Spurr GB., et al. "Energy expenditure of urban Colombian women: a comparison of patterns and total daily expenditure by the heart rate and factorial methods". *The American Journal of Clinical Nutrition* 63 (1996): 870-878.

16. Stein TP, et al. "Energy expenditure and socioeconomic status in Guatemala as measured by the doubly labeled water method". *The American Journal of Clinical Nutrition* 47 (1988): 196-200.
17. Taubes B. "As obesity rates rise, expert struggle to explain why?" *Science* 280 (1998): 1367-1368.
18. WHO Expert Committee. "Physical Status: The use and interpretation of anthropometry". Geneva, WHO (2000).
19. Williams PT. "Vigorous exercise and the population distribution of body weight". *International Journal of Obesity* 28 (2004): 120-128.
20. Williamson DF, et al. "Recreational physical activity and ten-year weight change in a US national cohort". *International Journal of Obesity* 17 (1993): 279-186.
21. World Health Organization. Obesity: preventing and managing the global endemic, WHO Technical Report Series 894 WHO: Geneva (2000).
22. Yamauchi T, et al. "Influence of urbanization on physical activity and dietary changes in Huli-speaking population: a comparative study of village dwellers and migrants in urban settlements". *British Journal of Nutrition* 85 (2001): 65-73.
23. Schmitz KH., et al. "Physical activity and body weight: associations over ten years in the CARDIA study. Coronary artery risk development in young adults". *International Journal of Obesity Relat Metab Disord* 24 (2000): 1475-1487.
24. Di Pietro L., et al. "Estimated change in physical activity level (PAL) and prediction of 5-year weight change in men: the Aerobics Center Longitudinal Study". *International Journal of Obesity Relat Metab Disord* 28.12 (2004): 1541-1547.
25. Haapanen N., et al. "Association between leisure time physical activity and 10-year body mass change among working-aged men and women". *International Journal of Obesity Relat Metab Disord* 21 (1997): 288-296.
26. World Health Organization. Global recommendations on physical activity for health. Geneva, Switzerland: *World Health Organization* (2010).
27. Tremblay MS., et al. "Sedentary Behavior Research Network (SBRN) - Terminology Consensus Project process and outcome". *International Journal of Behavioral Nutrition and Physical Activity* 14 (2017): 75.
28. Batty GD and Lee IM. "Physical activity and coronary heart disease". *British Medical Journal* 328 (2004): 1089-1090.
29. Thorp AA., et al. "Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011". *American Journal of Preventive Medicine* 41 (2011): 207-215.
30. Yen CF, et al. "The relationships between body mass index and television viewing, internet use and cellular phone use: the moderating effects of socio-demographic characteristics and exercise". *International Journal of Eating Disorders* 43 (2010): 565-571.
31. Drenowatz C., et al. "The association of physical activity during weekdays and weekend with body composition in young adults". *International Journal of Obesity* (2016): 8236439.
32. Keating C., et al. "Prevalence of class-I, class-II and class-III obesity in Australian adults between 1995 and 2011-12". *Obesity Research and Clinical Practice* 9 (2015): 553-562.
33. Cooper R., et al. "Obesity history and daily patterns of physical activity at age 60-64 years: Findings from the MRC National Survey of Health and Development". *The Journals of Gerontology, Series A* 72 (2017): 1424-1430.
34. Cooper R., et al. "Body mass index from age 15 years onwards and muscle mass, strength, and quality in early old age: findings from the MRC National Survey of Health and Development". *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 69 (2014): 1253-1259.
35. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants". *Lancet* 387 (2016): 1377-1396.
36. Nead KT., et al. "Contribution of common non-synonymous variants in PCSK1 to body mass index variation and risk of obesity: a systematic review and meta-analysis with evidence from up to 331 175 individuals". *Human Molecular Genetics* 24 (2015): 3582-3594.
37. Menard S. "Coefficients of determination for multiple logistic regression analysis". *American Statistical Association (ASA)* 54 (2000): 17-24.
38. R Core Team R. "A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing (2014).
39. Reddon H., et al. "Physical activity and genetic predisposition to obesity in a multiethnic longitudinal study". *Scientific Reports* 6 (2016): 18672.

40. Popkin BM., et al. "Global nutrition transition and the pandemic of obesity in developing countries". *Nutrition Reviews* 70 (2012): 3-21.
41. Swinburn BA., et al. "The global obesity pandemic: shaped by global drivers and local environments". *Lancet* 378 (2011): 804-814.
42. Sallis JF., et al. "Assessing perceived physical environmental variables that may influence physical activity". *Research Quarterly for Exercise and Sport* 68 (1997): 345-351.
43. Powell LM., et al. "Availability of physical activity-related facilities and neighborhood demographic and socioeconomic characteristics: a national study". *American Journal of Public Health* 96 (2006): 1676-1680.
44. Evenson KR., et al. "Physical activity and sedentary behavior patterns using accelerometry from a national sample of United States adults". *International Journal of Behavioral Nutrition and Physical Activity* 12 (2015): 20.
45. Lee PH., et al. "A cluster analysis of patterns of objectively measured physical activity in Hong Kong". *Public Health Nutrition* 16 (2013): 1436-1444.
46. Corder K., et al. "Predictors of change differ for moderate and vigorous intensity physical activity and for weekdays and weekends: a longitudinal analysis". *International Journal of Behavioral Nutrition and Physical Activity* 10 (2013): 69.
47. Stata Corp. "Stata Multiple-Imputation Reference Manual Release 13". College Station, TX: Stata Corp LP (2013).
48. World Health Organization. BMI classification (2018).

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667