



Exercise Testing in Children with Cerebral Palsy

Leunkeu AN^{1*}, Shephard RJ² and Ahmaidi S¹

¹Research Laboratory, "Adaptations Physiologiques à l'Exercice et Readaptation à l'Effort", Faculty of Sciences and Sport, University of Picardie Jules Verne, France

²Faculty of Kinesiology and Physical Education, University of Toronto, Canada

***Corresponding Author:** Leunkeu AN, Research Laboratory, "Adaptations Physiologiques à l'Exercice et Readaptation à l'Effort", Faculty of Sciences and Sport, University of Picardie Jules Verne, France.

Received: April 22, 2021

Published: June 15, 2021

© All rights are reserved by **Leunkeu AN., et al.**

Abstract

Objective: The aim of this study is to examine if cardiorespiratory requirements assess during the 6-minute walking test (6WT) and the cardiopulmonary exercise test (CPET) before and after training are the same in children with cerebral palsy (CP).

Methods: Twenty-four children with CP (16 boys and 8 girls, with the mean age of 14.2 ± 2 years (10 to 16 years) classified at gross Motor Function Classification System level I and II were recruited from a school for special education in France. Participants were divided in two groups of twelve children matched for age, weight, height, sex, type of cerebral palsy and growth motor function classification system (GMFSC). One group ($n = 12$) underwent CPET before and after the 8-week training programme twice per week for 30 minutes (+5min of warm up and 5 minutes of rest), and another group ($n = 12$) assigned the 6WT before and after the 8-week training programme twice per week for 30 minutes.

Results: Cardiorespiratory data did not differ between the CPET and the 6WT before training ($VO_{2\text{Peak}}$: $p = 0.86$; VE_{Peak} : $p = 0.87$ and HR_{Peak} : $p = 0.41$) and after training ($p = 0.41$, $p = 0.26$ et $p = 0.65$) respectively.

Conclusion: These findings demonstrate that cardiorespiratory responses obtained during the 6WT and the CPET before and after training did no differ in children with cerebral palsy.

Keywords: Children with CP; Cardiopulmonary Exercise Test; 6WT; Exercise Training

Introduction

Exercise fitness is commonly used to assess the functional status of children with cerebral palsy (CP) [1,3,20,23]. Some physiotherapists used the six-minute walk test [1,2] or the cardio Pulmonary exercise test (CPET) in interventions [1,2,14,20], or before and after certain surgery or treatments in children with CP [1,2,14]. However, there is paucity studies which think to explore the 6WT and the CPET before and after training. Whereas it could

be informative to provide an excellent tool in clinical rehabilitation. We had already investigated the 6WT and the CPET in children with CP [2,14]. Even if the tests are used in rehabilitation programs, we did not compare the both tests before and after training. However, it could be useful to help and provide some information keys to clinicians. So that, the choice of each test can match with some type of CP children that professional want to evaluate. Meanwhile, no data are yet available concerning the comparison of the 6MWT and the

CPET in children with cerebral palsy. By doing the comparison of the 6MWT and the CPET in children with CP GMFCS levels I and II, the medical staff who work for the rehabilitation and treatments in this population of children can feel free to realize a choice which is appropriated.

Purpose of the Study

The purpose of this study is to provide the comparison of the cardio respiratory parameters of the 6WT and the CPET in children with CP before and after training; and then it overview of it current applications in each type of children with CP.

Methods

Participants

Twenty-four children with CP (age, 14.2 ± 2 years; weight 45.5 ± 7.8 kg; height 154 ± 11cm) familiarized with cycling activity participated in this study. All were volunteers from local specialized school. All subjects underwent a cardiorespulmonary test (CPET) and had a standardized 6WT with gas exchange measurement with a portable gas analyser (Cosmed K4b², Cosmed, Rome, Italy). They were divided in two groups of twelve children matched for age (14 ± 2 years); weight (43 ± 5.7 kg), height (158 ± 14 cm), sex (8 boys and 4 girls), type of CP, and GMFCS. One group underwent CPET before and after the 8-week training programme of the CPET and the other group did 6WT before and after the 8-week training programme of the 6WT with gas exchange measurement with a portable gas analyser (Cosmed K4b², Cosmed, Rome, Italy). All children were spastic classified at level I or II on the gross motor function classification system (GMFCS) as expanded and revised for children > 12 years [17]. Each group represented their own control group. Twenty children were classified as hemiplegic (involvement of one side of the body) and four were classified as diplegic (involvement of the lower part of the body). The participants were independent ambulators and were able to follow simple verbal commands. Participants were excluded if they had had orthopaedic surgery or neurosurgery and/or botulinum toxin injection (s) within 6 months prior to study entry or cardiac or respiratory conditions that could negatively be affected by exercise. The study was approved by a local ethics committee and in accord with the ethical standard of the Helsinki Declaration of 1975 and 1983 [5] and in an investigation that met the ethical standards of this journal as detailed by Harris and Atkinson [10]. Children were excluded if they undergone a surgical procedure influencing mobility during the

past year, were taking medication that could affect muscular function. A medical examination was performed and anthropometric data were collected on each subject. The clinical characteristics of the children with CP are stated in table 1. A physician assessed the spasticity of each child, using the method of Ashworth [4]. All children were independently ambulatory and maturity was assessed by Tanner staging [11,22]. For children, an informed consent was obtained from the parents as well as from the child. Before the testing sessions, a medical examination was performed and their anthropometric data were measured.

Characteristic	Children and adolescents with CP (6WT, n = 12)	Children and adolescents with CP (CPET, n = 12)
Age (years)	14 ± 1.8	14 ± 1.9
Body mass (kg)	45.4 ± 6.3	45.5 ± 5.7
Height (m)	1.58 ± 0.12	1.59 ± 0.13
Sex: boys/girls	8 boys/4 girls	8 boys/4 girls
Hemiplegia	10	10
Diplegia	2	2
GMFCS	Level I: 9, level II: 3	Level I: 9, level II: 3

Table 1: Clinical characteristics of children and adolescents with CP GMFCS levels I and II (Values are mean values ± SD).
 CP = Cerebral Palsy, 6WT = Six Minutes Working Test,
 CPET = Cardiopulmonary Exercise Test.

Cycle-ergometer exercise testing

Exercise testing was performed on a leg cycle ergometer (Ergometer 824 E, Monark Exercise AB, Kroonsväg 1, S-780 50 Vansbro, Sweden) using previously determined protocol [14]. Thorough familiarization and habituation procedures were followed as recommended [14]. Children with CP underwent two cycle ergometer exercise testing sessions. The first test was an habituation test, which was also used to check the protocol and to determine whether the children with CP needed to be attached to the pedals [19]. One week later, children then underwent an incremental protocol exercise test on the cycle-ergometer with an initial power of 30 W and was increased thereafter by 15 W every minute. The pedalling rate was fixed at 50 rpm and the cadence was maintained by using a metronome. The feet were fixed to the pedals by velcro strapping. Individual protocols were constructed such that the total exercise

time ranged between 6 and 12 minutes. All the tests were preceded by a 5 min warm-up period.

Vigorous verbal encouragements were given throughout the tests to achieve maximal effort. During the CPET, cardiorespiratory gas exchange measurements were obtained with the portable gas analyzer (Cosmed K4b², Cosmed, Rome, Italy). The inspiratory air flows and the fraction of expired O₂ (F_E O₂) and CO₂ (F_E CO₂) were calculated by computer every 5 respiratory cycles. One minute averages were then established for minute ventilation (V_E: L.min⁻¹, BTPS), O₂ uptake (VO₂: L.min⁻¹, STPD), CO₂ production VCO₂ (L.min⁻¹, STPD) ventilatory equivalent for O₂ (VE.VO₂⁻¹), and CO₂ (VE.CO₂⁻¹) and breathing frequency (f, cycle.min⁻¹). The test ended when fatigue or dyspnea occurred making the child stop the effort. The VO₂ peak obtained at the end of maximal exercise testing was considered as the maximal VO₂ uptake. Exercise tolerance and cardiorespiratory requirement were assessed through the values obtained at maximal effort: oxygen uptake (VO₂), ventilation (V_E), heart rate (HR). At the end of the test, subjects had 5 minutes of passive recovery.

The 6-minute walking test

Before they began the 6WT, children with CP were fitted with a portable gas analyzer Cosmed K4b², (Cosmed, Rome, Italy) to assess respiratory gas exchanges. Children with CP performed the 6WT between 2 lines set 20 m apart in the corridor, according to usual recommendation [5,16]. Cardiorespiratory requirement was assessed from cardiorespiratory gas exchange measures: oxygen uptake (VO₂ in mL.kg⁻¹.min⁻¹), ventilation (V_E in L.min⁻¹), and heart rate (in beats.min⁻¹). Samples of these variables were taken during rest, 6 minutes during exercise, and 5 minutes during the recovery period. One-minute averages were then established for minute ventilation (VE: L.min⁻¹, BTPS), O₂ uptake (VO₂: L.min⁻¹, STPD), CO₂ production VCO₂ (L.min⁻¹, STPD) ventilatory equivalent for O₂ (VE.VO₂⁻¹), and CO₂ (VE.CO₂⁻¹) and breathing frequency (f, cycle.min⁻¹). All tests were performed between 9:00 AM and 12:00 PM, in the same place, and followed the same protocol to control for circadian variation. Participants were instructed to walk as far as possible and to cover the distance as long as possible in 6 minutes. They also received vigorous verbal encouragements with the standard statements. Every 30 seconds subjects were advised of the remaining time and distance covered. They turned round a marker at each end of the 20 m [8]. Furthermore, they did not use any orthotics or

forearm crutches. The distance covered was measured at the end of the test.

Borg scale

Children quoted the exercise perception during both the 6WT and CPET at 5 minutes after the tests stopped. The Borg Scale with the level range from 6 to 20 was used [7]. Children were instructed to put a cross on a no graduated scale between two adjectives (very very easy to very very difficult) indicating the degree of the difficulty they felt during the tests.

The 6-minute walking test exercise training

The group of the 12 children with CP were tested first and then have a training. Before the exercise training started, a trial of 30 minutes was performed by the training group to check out the speed used as training intensity. The training intensity used corresponded to the mean walking speed recorded on the 2 preceding 6 WT trials. Once the initial training walking speed was determined, the exercise training started 3 days after. Prior to and following the training programme, 6WT with cardiorespiratory gas analysis was conducted: oxygen uptake (VO₂), ventilation (V_E) and heart rate (HR) were measured. The training programme consisted of 30 min of walking three times per week for eight weeks. All training sessions were performed in the morning at 10:00 am on the same place (indoor) and according to the same protocol. Exercise's sessions were performed 3 times per week, each on a separated day. The training was supervised by a physician and a paediatric physical therapist. To enhance the palatability of the training program, verbal encouragement was given to all the children to motivate them to finish the training session. To control for circadian variation, all tests were performed between 9:00 am and 12:00 (noon). To assess the effect of the eight weeks of exercise training, children were tested immediately after the eight-week using the 6 WT.

The cycle-ergometer exercise training

Training sessions were held three times per week, on separated days (Monday, Wednesday and Friday), 40 min (5 min of warm-up + 30 min of test + 5 min of recovery) at 50% the first and the second week, at 55% the third and the fourth week, at 60% the fifth and the sixth week and at 65% the seventh and the eighth week. Cycle ergometer training was chosen because most children with CP who have problems with movement coordination and equinus position of the foot find difficulty with the increasing speed and

slope of treadmill exercise [13]. All training sessions were held at 10 am; immediately previously, the 12 children underwent 30 minutes of stretching of leg, thigh and arm muscles and breathing exercises under the supervision of the physiotherapist. They were given verbal encouragement to complete each training session. Training began with a 5 min warm up; they then undertook 30 minutes of pedalling (50 rpm with an initial power output of 30 w), 50% of the VO_2 observed. The training load increased by 15 w every minute during the training. When they improved their performance, we increased the intensity two weeks by two weeks until they reached eight weeks. To assess the effect of the eight weeks of exercise training, children were tested immediately after the eight weeks.

Statistical analysis

All data are expressed as mean, standard deviation (SD). The normality of the data was tested using Kolmogorov-Smirnov test. The cardio respiratory data (Peak VO_2 , Peak V_E and Peak HR) obtained during 6WT were compared to those assessed during the CPET before and after training period, and the Borg scale using an ANOVA (Statview 5.0, SAS, USA). For the correlation between the 6MWT and the CPET, we used Spearman correlation. A p value less than 0.05 was considered significant.

Results

Participants

A total of twenty-eight participants were assessed for eligibility for this study: four participants did not complete the study. Two participants were awaiting surgery and were obligated to stop the training sessions before the end. Two participants did not wear a face mask because the thought of breathing through the mouth-piece frightened them. Twenty-four participants completed the entire study.

Comparison of the 6WT and the CPET cardio respiratory peak responses before training

The cardiorespiratory responses obtained during the 6WT and the CPET at the start of the 8 weeks in children with CP did not show significant differences for $VO_{2\text{ peak}}$ (6WT: 33.13 ± 7.14 vs. CPET: 32.31 ± 6.28 mL.kg⁻¹.min⁻¹, $p = 0.86$); VE_{peak} (6WT: 50.70 ± 23.50 vs. CPET: 49.03 ± 18.69 L.min⁻¹, $p = 0.87$) and HR_{peak} (6WT: 156.62 ± 22.42 vs. CPET: 148.40 ± 25.1 beats.min⁻¹, $p = 0.41$). Cardio respiratory peak values of the TM6 were 0.90% for the VO_2 ,

1,19% for the VE et 1,05% for the FC high to those that measured in CPET respectively. Moreover there was a correlation for $VO_{2\text{ peak}}$ ($r = 0.80$), VE_{peak} ($r = 1$), and HR_{peak} ($r = 0.80$). The results of the comparison of the cardio respiratory 6WT and the CPET before training are shown in figure 1.




Figure 1: Cardiorespiratory data of $VO_{2\text{ peak}}$ (Figure 3A), V_E peak (Figure 3B) and HR peak (Figure 3C) measured during the 6WT and the CPET before training period. Values are mean values \pm standard deviation (SD). NS: no significant.

Comparison of cardio respiratory data of the 6WT and the CPET after training

Peak VO_2 , peak VE and peak FC of the 6WT after training did not differ from those of the CPET after training ($p = 0,41$, $p = 0,26$ et $p = 0,65$) respectively. Cardio respiratory peak values of the 6MWT were 0.97% for the VO_2 , 1,05% for the VE et 1,04% for the FC high to those that measured in CPET. Moreover, there was a correlation for $VO_{2\text{ peak}}$ ($r = -0.40$), VE_{peak} ($r = 0.92$), and HR_{peak} ($r = 0.50$). The

results of the comparison of the cardio respiratory data 6WT and the CPET after training are shown in figure 2.




Figure 2: Cardiorespiratory data of VO₂ peak (Figure 3A), V_E peak (Figure 3B) and HR peak (Figure 3C) measured during the 6WT and CPET after training period. Values are mean values ± standard deviation (SD). NS: no significant.




Figure 3

Discussion

This study demonstrates that the cardio respiratory responses obtained during the 6WT and the CPET were similar before and after training in children with CP classified at GMFCS level I and II.

The novelty of our investigation was to study comparison of the cardio respiratory requirement during the 6WT and the CPET before and after training. Results obtained demonstrated that the cardio respiratory responses obtained during the 6WT were similar to those obtained during the CPET. Indeed, O₂ uptake, ventilation, heart rate responses and performance did not differ between these two tests. The results from the present study agree with the previous studies conducted in healthy children and children with other pathologic populations [16]. These also indicated the utility of the 6WT and the CPET for assessing functional capacity and cardio respiratory fitness in children with CP classified at GMFCS level I and II. Moreover, Borg scale [7] outlined that the exercise perception was similar in both tests in children with CP. It could be assumed that, under this condition, the quotation of the perception of the effort confirmed the fact that cardio respiratory gas exchanged during the 6WT and the CPET are not different in children with CP GMFCS level I and II.

Given that physiotherapist and clinicians consider making the 6MWT a standard component of assessments for children and adolescents with CP at GMFCS levels I and II [2], the correlation with the CPET can be a type of measurements which potentially aid in treatment planning and in rehabilitation fitness in child with CP [14,21]. We previously found the effect of exercise training in children with CP GMFCS levels I and II during the 6MWT and the CPET [2,14]. We though and feared that CPET could not had been matched, and feet, and appropriate for children with CP, when we found that children with CP usually show quadriceps femoris weakness and thus a poor peak aerobic power when exercise isometric test [12,15]. As it is well known that children and adolescents with CP are very weak and have lower performance, related fitness and cardiorespiratory fitness compared to children who are typically developing [1,3,14]. Then, we focus our study on the comparison of cardiorespiratory data of the 6MW and the CPET instead of the treadmill because speed remained problematic for CP [18]. In the literature there is study assessed a new reference values for cardiorespiratory exercise testing in children [6]. Whereas their study did not mentioned if their methodology can be applied in children with cerebral palsy.

The current study compared the cardiorespiratory data (VO₂, ventilation and heart rate) between the 6MWT and the CPET before and after training in children with CP. The results demonstrate that the cardio respiratory responses obtained when exercise during the both test are similar. To the best of our knowledge, this is the first demonstration of the comparison of two methods of aerobic fitness: the 6MWT and the CPET in children with CP GMFCS level I and II. We found in Borg Scale [7] that children with CP GMFCS level I and II felled the same effort during the 6MWT and the CPET. This result argued with the opinion that the 6MWT and the CPET can be both feasible in children with CP GMFCS levels I and II. We also found a correlation between the 6MWT and the CPET before and after training in those children. However, some studies only shown the comparison without correlation and more over without training. Whereas it is necessary and usual for the fiability and the validity of a test to assess the comparison before and after training. Moreover, in clinical rehabilitation especially in children with cerebral palsy. We have now found that the 6MWT and the CPET in children with CP GMFCS level I and II could similarly assess aerobic fitness in children with CP GMFCS levels I and II. We provided and recommended to medical staff or searcher in children with CP to find this comparison like a good tool of evaluation in fitness, training and the improvement of the healthiness in their practices with their patients.

Conclusion

Viewed in concert the data of the 6MWT matched closely to the findings of the laboratory cycle-ergometer testing before and after training in children with CP GMFCS level I and II. Moreover, the 6MWT and the CPET can be used to evaluate cardiopulmonary function during rehabilitation program in children with CP GMFCS level I and II. These insights in the positive effects of exercise training should ideally lead to implementation in clinical practice.

Acknowledgement

We acknowledge the cooperation and expert technical assistance of the medical staff of the cerebral palsy center during data collection and processing, and the generous cooperation of the children and parents participating in this study. The authors disclose any potential conflict of interest.

Conflict of Interest

None of the author have a financial relationship with any commercial entity that has an interest in the topics covered by this manuscript.

Financial Disclosure

We certify that no party having a direct interest in the results of the research supporting this article has or will confer a benefit on us on any organization with which we are associated AND, if applicable, we certify that all financial and material support for this research (e.g., NIH or NHS grants) and work are clearly identified in the title page of this manuscript.

Bibliography

1. Angeline N Leunkeu., *et al.* "Optimizing Rehabilitation: The potential to Assess cardiorespiratory, Neuromuscular and Biomechanical Adaptations to Exercise of Children with Cerebral Palsy in the face of Intra-Individual Variation". *Journal of Novel Physiotherapy and Physical Rehabilitation* 4.2 (2017): 042-047.
2. Angeline Nsenga Leunkeu and Said Ahmaidi. "Longitudinal assessment of muscle function and oxygen cost during isometric testing in children with cerebral palsy". *Isokinetics and Exercise Science* 20 (2012): 1-7.
3. Angeline Nsenga Leunkeu., *et al.* "Six-Minute Walk Test in Children with Cerebral palsy Gross Motor Function Classification System Levels I and II: Reproducibility, validity and Training effects". *Archives of Physical Medicine and Rehabilitation* 93 (2012): 2333-2339.
4. Ashworth B. "Preliminary trial of carisoprodol in multiple sclerosis". *Practitioner* 192 (1964): 540-542.
5. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. "ATS statement: Guidelines for the six-minute walk test". *American Journal of Respiratory and Critical Care Medicine* 166 (2002): 111-117.
6. Blanchard J., *et al.* "New reference values for cardiopulmonary exercise testing in children". *Medicine and Science in Sports and Exercise* 50.6 (2018): 1125.
7. Borg GA. "Psychophysical bases of perceived exertion". *Medicine and Science in Sports and Exercise* 14 (1982): 377-381.
8. Enright PL. "The six-minute walk test". *Respiratory Care* 48 (2003): 783-785.
9. "Ethical Standards in Sport and Exercise Science Research". *International Journal of Sports Medicine* 30 (2009): 701-702.

10. Harris DJ and Atkinson G. "Update – ethical standards in sport and exercise science research". *International Journal of Sports Medicine* 32.11 (2011): 819-821.
11. Henderson RC, et al. "Skeletal maturation in moderate to severe cerebral palsy". *Developmental Medicine and Child Neurology* 47.4 (2005): 229-236.
12. Leunkeu AN, et al. "Electromyographic (EMG) Analysis of quadriceps femoris muscle fatigue in children with cerebral palsy during a sustained isometric contraction". *Journal of Child Neurology* 25.3 (2010): 287-293.
13. Maltais D, et al. "Repeated treadmill walks affect physiologic responses in children with cerebral palsy". *Medicine and Science in Sports and Exercise* 35 (2003): 1653-1661.
14. Nsenga AL, et al. "Aerobic Training in Children with Cerebral Palsy". *International Journal of Sports Medicine* 34 (2013): 533-537.
15. Nsenga Leunkeu A, et al. "Cardiopulmonary exercise data during quadriceps isometric contraction sustained fatigue in children with cerebral palsy". *Isokinetics and Exercise Science* 17 (2009): 27-33.
16. Paap E, et al. "Physiologic response of the six minute walk test in children with juvenile idiopathic arthritis". *Arthritis and Rheumatology* 53 (2005): 351-356.
17. Palisano R, et al. "Validation of a model of Gross Motor Function for children with Cerebral Palsy". *Physical Therapy* 80 (2002): 974-985.
18. Potter C and Unnithan VB. "Interpretation and implementation of oxygen uptake kinetics studies in children with spastic cerebral palsy". *Developmental Medicine and Child Neurology* 47 (2005): 353-357.
19. Rogers A, et al. "Systematic review of the effectiveness of aerobic exercise interventions for children with cerebral palsy: an AACPD evidence report". *Developmental Medicine and Child Neurology* 50 (2008): 808-814.
20. Takken Tim, et al. "Cardiopulmonary Exercise Testing in Pediatrics". *American Thoracic Society* 14.1 (2017): 123128.
21. Verschuren O, et al. "Exercise programs for children with cerebral palsy: a systematic review of the literature". *American Journal of Physical Medicine and Rehabilitation* 87.5 (2008): 404-417.
22. Worley G, et al. "Secondary sexual characteristics in children with cerebral palsy and moderate to severe motor impairment: a cross-sectional survey". *Pediatrics* 110.5 (2002): 897-902.
23. Zwinkels Maremka, et al. "Body mass index and Fitness in High-Functioning Children and adolescents with Cerebral Palsy; What Happened over a decade?" *Research in Developmental Disabilities* 71 (2017): 70-76.

Volume 4 Issue 7 July 2021

© All rights are reserved by Leunkeu AN, et al.