Acta Scientific NEUROLOGY

Volume 8 Issue 8 August 2025

Review Article

Understanding and Managing Post-Stroke Depression: A Comprehensive Guide to Recovery

Ibrahim Krenawi[1]*, Victor Hugo Post[2], Markus Müller[3] and Mohamed Mustafa[4]

[1]*Consultant Neurologist, Burjeel Royal Hospital, Al Ain, UAE*

[2]*Senior Consultant Psychiatrist & Psychotherapist , LVR-Klinik, Bedburg-Hau, Germany*

[3]*Resident Physician in Psychiatry, Klinikum Hersfeld-Rotenburg, Germany*

[4]*Specialist Psychiatrist & Psychotherapist, Praxis CityNord for Psychiatry und Psychotherapy, Dortmund, Germany*

***Corresponding Author:** Ibrahim Krenawi, Consultant Neurologist, Burjeel Royal Hospital, Al Ain, UAE.

DOI: 10.31080/ASNE.2025.08.0849

Received: July 14, 2025

Published: July 21, 2025

Abstract

The problem of post-stroke depression (PSD) is critical and widespread, as some stroke survivors often delay rehabilitation and experience a decreased quality of life. This review discusses the underlying causes, risk factors, symptoms, diagnostic plans, treatment plans, and lifestyle interventions for PSD. The disorder is a complex interplay of neurological injury, emotional stress, and social isolation following a stroke.

Some of the most typical symptoms and manifestations include persistent depressive affect, fatigue, diminished interest in activities, alteration of appetite, and cognitive problems, which are often confused with the overall impact of stroke recovery. Diagnosis is achieved mainly through clinical interviewing, standardized screening instruments, and medical examination in order to eliminate other disorders. Treatment options include antidepressant drugs, psychotherapy, sociability, exercise, and alternative methods such as mindfulness. Other considerations in the management of PSD include lifestyle changes, which can take the form of physical activity, a healthy diet, and emotional support. Early interaction, constant observation, and the participation of caregivers are key to enhancing outcomes. PSD is manageable with a comprehensive care plan, even though it cannot be prevented. To ensure that physical health and emotional status are in a better condition, this review focuses on the necessity of identifying and treating depression as a means of improving stroke recovery.

Keywords: Post-Stroke; Depression; Guide; Recovery

Introduction

The process of overcoming a stroke is not only a physical one. Although physiotherapy, medicine, etc., usually aim at recovering movement, speech, or coordination etc., another important aspect of recovery being ignored by many, in most cases, is mental health, specifically post-stroke depression (PSD).

Post-stroke depression is prevalent and severe, existing in high numbers of stroke survivors [1]. It involves not only a depressed mood but also constant fatigue, subdued interest in life, anxiety, and emotions of utmost stressfulness. Such emotions can complicate the recovery process, postpone rehabilitation, and even result in an amplification of the chance of additional health problems.

Most individuals believe that depression is just triggered by a traumatic event in their lives, but that is not the case with PSD. Emotional trauma, as well as changes in brain chemistry, both contribute. What adds to the predicament is its tendency to go undiagnosed, since individuals are more concerned with physical ailments.

The good news is that depression that occurs after a stroke is curable. Many survivors are successfully able to move on, partially physically and emotionally, with the appropriate mix of care, which includes medication, therapy, lifestyle modification, and social support [2].

In this paper, we will inform you of all that you ought to know about PSD: what causes it, the symptoms, its diagnosis, as well as the best methods of treatment and prevention. Whether you are a stroke survivor or a caregiver, or even a person with an interest to learn more about this condition, this guide is ready to assist you in taking the next step to full recovery, both in body and mind.

Causes and risk factors of post-stroke depression

Stroke is one of the most common causes of death in the Western world. Numerous epidemiological studies show that post-stroke depression (PSD) is observed as a complication in nearly 50% of cases. Between 25% and 35% of all patients develop major depression within the first year after a stroke. According to ICD-11 and DSM-5, "depressive disorder due to another medical condition" is defined as a depressive mood, loss of interest, or anhedonia, with indications from physical examination, laboratory tests, or technical investigations suggesting that the disorder is a direct pathophysiological consequence of another medical condition. A potential direct pathophysiological connection is discussed for the most common physical conditions associated with depressive disorders, such as diabetes mellitus, cardiovascular diseases, epilepsy, multiple sclerosis, or stroke itself [3]. In recent years, there has been increasing evidence that stroke patients with depression show higher mortality rates [4]. Conversely, a history of depression itself is a risk factor for subsequent stroke, affecting both morbidity and mortality [5,6]. In the multifactorial transmission of this risk, not only psychological factors play a crucial role, but also a range of neurobiological factors, which are shared along with the general risks for cardiovascular diseases or diabetes mellitus [7].

Whether the location of the stroke plays a role is currently not definitively assessable.

The following factors are correlated with the risk of post-stroke depression

- Size of the infarct
- Extent of physical disability
- Age (younger patients are more frequently affected)

The hypotheses regarding the significance of stroke localization in the development of post-stroke depression are contradictory. Influencing factors for the development of post-stroke depression are diverse, with the location and size of cerebral damage, as well as the extent of physical disability, being significant, along with age (younger patients often develop post-stroke depression). While the extent of neurological deficits caused by the stroke appears to be important for the risk of depression, this is less relevant in the acute phase and more so in the long term, considering interactions with other factors, such as certain personality traits or the availability of psychosocial support [7]. Post-stroke depression is significantly associated with poorer recovery of cognitive performance in affected patients, with the relationship between depression and cognitive deficits being discussed bidirectionally in the scientific literature; on one hand, post-stroke depression is seen as a consequence of cognitive deficits, and on the other, cognitive deficits can be a result of post-stroke depression [8].

Post-stroke depression (PSD) lacks a single cause. It normally occurs due to a combination of physical, emotional, and social changes following a stroke [9]. Being aware of these causes can help patients and caregivers identify what is going on and seek the necessary help.

The causes of post-stroke depression (PSD) are diverse and can be both biological and psychological in nature:

- Biological Factors: Damage to the brain from the stroke, changes in neurotransmitter balance, and hormonal influences may play a role.
- Psychological Factors: Processing the traumatic event, coping with losses (physical abilities, independence), difficulties in adjusting to the new living situation, and social factors (isolation, support) can contribute to the development of depression.
- Chemical Imbalance and Brain Alterations
- Emotional Response to Change in Life
- Isolation and Loneliness
- Constant stress and worry

In summary, post-stroke depression has considerable social and medical relevance, as it significantly impairs the rehabilitation of affected patients, delays reintegration into the workforce, and contributes to increased healthcare costs [5].

Symptoms of post-stroke depression

Post-stroke depression often occurs within the first weeks or months following a stroke. It is a serious condition that can significantly impair recovery and quality of life for those affected.

Timeline

- **Early Phase:** Depression can manifest shortly after the stroke, often within the first few weeks.
- **Mid-term:** Many individuals experience depression when struggling with physical limitations and the impact on daily life.
- **Long-term:** In some cases, depression may develop months or even years after the stroke.

The detection of post-stroke depression (PSD) may not be a simple process, particularly because post-stroke survivors are dealing with numerous physical and emotional transformations [10]. However, being aware of the signs can make a difference. Early diagnosis can lead to quicker treatment and an easier recovery process.

These are the most common symptoms to pay attention to

- **Constant Depression or Depressiveness:** A feeling of persistent depressive affect that does not appear to disappear is one of the most evident symptoms of PSD [11]. It becomes more than a bad day; this low mood stays for several weeks and months, and often there is no obvious explanation.
- **Anhedonia (Loss of Interest or Pleasure in Activities):** Things that used to bring pleasure, such as hobbies, social life, or even favorite foods, are suddenly seen as useless. This loss of feeling, referred to as anhedonia, is one of the signs of depression, and it may influence one a lot in terms of motivation for recovery [12].
- **Energy Depletion/Fatigue:** Another common symptom is fatigue even when they are resting [13]. This is not physical, but emotional and mental burnout where one feels that any task is a burden to accomplish.
- **Alteration of Sleep Patterns:** Individuals with PSD can oversleep or suffer from insomnia. Both ends have the potential to interfere with healing and exacerbate emotional health.
- **Guilt, Worthlessness, Hopelessness Feeling:** A sizable number of stroke survivors are burdened or even blame themselves because of the condition. Such negative lines of thinking have the potential to stoke up depression and must not be overlooked [14].
- **Cognitive Impairment:** The cognitive symptoms of mental fogginess, difficulty concentrating, or a feeling of not being integrated are also very common in PSD [14]. This may complicate the process of decision-making and communication.
- **Suicidal or Death Ideation:** In some extreme cases, a person might indicate that he or she wants to die or even mention suicide [15]. These are serious warning signs, and these need to be treated in the hospital at once.
- **Remember:** All these symptoms may not be exhibited by all people, and at times they can be subtle. If you notice that you/or someone you love has an issue after a stroke that could be showing signs of depression, do not hesitate, consult a physician professional immediately. Timely help can transform everything.

Diagnosis Post-Stroke Depression

Even after a stroke, diagnosing post-stroke depression (PSD) may not be easy, since the condition easily mixes with other consequences of the health disaster [16]. For example, it is possible to feel tired, have difficulties in speaking, or feel confused, which may be attributed to the stroke or the depression, or both. This is the reason why the condition of PSD is being diagnosed with a little extra time, attention, and care. Diagnosing post-stroke depression requires a comprehensive clinical evaluation rather than a brief symptom checklist.

The following is how doctors normally go about diagnosing whether a person is depressed after a stroke

Clinical Interview with the Patient and Family: The best initial stage is a talk. A medical practitioner or psychological expert will sit with the stroke patient to ascertain how he or she has been feeling emotionally since the stroke [17]. These are questions they will ask.

- "Are you sad or hopeless?"
- "Do you like the same things that you liked before?"
- "How do you sleep and eat?"
- "Are you ready to live with the idea that life is not worth living?"

Application of Depression Screening Tools: To determine the degree of depression, doctors usually resort to simple questionnaires or checklists in addition to the conversation [18]. The tools are supposed to be quick and simple to fill out, and they provide the doctor with a better idea of the severity of the symptoms.

A few of the screening tools are

- **PHQ-9 (Patient Health Questionnaire-9):** It is a 9-question short form. It enquires how frequently you have experienced things such as depressive effects, tiredness, or even hopelessness in the last two weeks. Every reply is rated, and the score allows the physician to identify whether the individual experiences mild, moderate, or severe levels of depression.
- **Hamilton Depression Rating Scale HAM-D:** Mostly, this one is used in the hospital or rehab facility. A physician will pose a group of questions and will count the responses. It puts emphasis on mood, sleep, appetite, as well as energy.
- **GDS (Geriatric Depression Scale):** This instrument is common among elderly people, particularly stroke patients. It has yes/no questions that are easier and less confusing to people who might not cope well with long conversations [19].
- **Beck Depression Inventory (BDI):** BDI is a widely validated, self-administered questionnaire designed to assess the presence and severity of depressive symptoms. Comprising 21 items, the BDI evaluates a broad range of affective, cognitive, motivational, and somatic symptoms associated with depression, including mood disturbance, pessimism, irritability, guilt, self-criticism, sleep disruption, fatigue, and appetite changes. Each item is rated on a 4-point scale (0-3), reflecting increasing symptom severity, yielding a cumulative score that categorizes depression as minimal, mild, moderate, or severe. The BDI is commonly used in both outpatient and inpatient settings and is particularly valuable for tracking symptom progression and treatment response over time.

Such screening instruments cannot provide a definitive diagnosis, but they aid in the comprehensive assessment and help in monitoring changes over time.

Checking for Other Medical Causes: Before concluding on whether a patient has post-stroke depression, physicians attempt to eliminate other possible causes of symptoms. The reason is that there are situations when the symptoms of vitamin deficiency, thyroid disease, or drug side effects may resemble depression.

For example

- Fatigue and depressive mood can be caused by low thyroid hormone (hypothyroidism).
- Some medicines, particularly those used after the stroke, could influence the mood or energy [20].
- Memory problems and low mood can be triggered by a deficit of vitamin B12 or folate.

Thus, the physician can request blood diagnostics or check which medications you take to be certain that there is no other problem lurking behind the symptoms of depression. This is a very significant step to ensure that the plan of action regarding treatment is correct and sufficient.

Treatment options for post-stroke depression

In the treatment of post-stroke depression, psychological and psychotherapeutic approaches, medication options, and non-invasive brain stimulation methods have been scientifically investigated [21].

Post-stroke depression (PSD) is not a self-limiting syndrome; it needs attention and care. The positive thing is that it is curable, and most individuals can greatly recover in mood, energy, and general recovery with the help of the right blend of therapies [34].

Now it is time to take a closer look at the primary treatment options

- **Medication (Antidepressants):** Medication, particularly antidepressants, is one of the most popular and successful routes of treatment for PSD [35]. They are medications that restore the levels of chemicals (such as serotonin and norepinephrine) in the brain, which affect moods.

The effectiveness of antidepressants in treating post-stroke depression was first demonstrated in a meta-analysis by Chen et al., based on 16 randomized controlled trials (RCTs) involving 1,320 patients compared to placebo [22]. These results have since been replicated multiple times, establishing antidepressants as an integral component in the treatment of post-stroke depression [23-25]. Chen., *et al.* (2007) also conducted a meta-analysis on the efficacy of antidepressants in preventing depression in non-depressed stroke patients (10 RCTs, 703 patients) and concluded that antidepressants can significantly reduce the risk of first-time occurrence of post-stroke depression [26].

Furthermore, results from the so-called FLAME study and a Cochrane meta-analysis indicated positive effects of SSRIs on rehabilitation after stroke, even in non-depressed patients [27,28]. Based on these positive findings, Flaster., *et al.* (2013) and Cholett., *et al*. (2013) authored reviews on the benefits of prophylactic antidepressant treatment in the rehabilitation of stroke patients [29,30]. However, two recently published large randomized, placebo-controlled studies found no positive effects of the SSRIs Citalopram or Fluoxetine on rehabilitation outcomes in non-depressed patients following acute stroke [31,32]. A recent Cochrane meta-analysis by Legg., *et al*. (2019) thus found no robust evidence for the routine use of SSRIs in stroke rehabilitation [33].

Based on the existing data, the prescription of antidepressants, regardless of the presence of depressive symptoms, for improving functional outcomes after strokes remains unclear and is therefore a subject of controversy.

- How it works: The damage to the brain that regulates emotions in a brain stroke is possible. It may cause a chemical imbalance, and it is difficult to be happy and motivated. Antidepressants assist in therapy to gain that balance.

Common types

- SSRIs (Selective Serotonin Reuptake Inhibitors) such as fluoxetine (Prozac), sertraline (Zoloft), and citalopram (Celexa) are the drugs most likely to be prescribed due to the lack of side effects and the fact that they are quite safe among stroke survivors.
- In some cases, SNRI (Serotonin-Norepinephrine Reuptake Inhibitors) such as venlafaxine (Effexor) could be applied.

Important notes

- Antidepressants may need 2 to 6 weeks to work.
- Serious side effects such as nausea, headache, or dizziness are quite uncommon and fade away with time.
- They are the medications that always have to be taken under medical supervision; never quit or change the dosage without a doctor's consultation.

However, the effectiveness of antidepressant treatment for post-stroke depression is scientifically well established. This treatment should be initiated as early and consistently as possible and should include an adequate treatment duration as well as an appropriate antidepressant dosage. Due to their anticholinergic side effects and the risk of cardiac complications, tricyclic antidepressants should be used cautiously. Overall, the favorable side effect profile and established efficacy support the use of SSRIs for treating post-stroke depression [21,25,26], provided there are no specific contraindications. Caution is advised when co-prescribing with anticoagulants or platelet aggregation inhibitors due to a potentially increased risk of bleeding.

Psychological Therapy (Talk Therapy or Counseling): Another effective treatment of depression that follows a stroke is therapy [36]. It assists the survivors to discuss their feelings, fears, and difficulties, and helps them on how to deal with emotional alterations.

The current psychological and psychotherapeutic treatment approaches primarily show moderate efficacy for cognitive behavioral therapy [7,21].

- **Cognitive Behavioral Therapy (CBT):** It is the most popular type of talk therapy. It assists individuals to become aware and transform unconstructive thoughts, and substitute them with more positive, real thinking [37].
- **Behavioral Activation Therapy:** The therapy focuses on boosting activities that enhance mood even in the event of low motivation.
- **Supportive Counseling:** In some cases, it is enough to be able to talk to a person who is understanding, and this will help to stop being lonely and feeling depressed.
- **Social Support and Peer Groups:** The recovery process after a stroke may be isolating. Most of the survivors respond that they do not feel understood by others about what they are experiencing. Here comes social support.

What it involves

- Face-to-face or online Stroke support groups where survivors can exchange stories, challenges, and tips.
- Family and friends support the guest by offering emotional support and motivation.
- Community-based programs provide resources, activities, and companionship.
- Why it's effective: Emotional health can be improved by spending time with like-minded individuals who understand them. One of the most natural antidepressants that we have is social connection.

Physical Activity and Exercise: Even light motion can brighten your mood-yes indeed! Research indicates that when you engage in physical exercise regularly, your depression levels are reduced since feel-good chemicals such as endorphins are released and blood flow into the brain improves.

Safe physical exercise tips

- A little stroll every day (even around the house or the garden)
- Sitting down for exercise or stretch exercises
- Tai Chi
- Gentle yoga
- Specific physical therapy exercises imposed by an expert

Benefits

- Lessens fatigue
- Better sleeping
- Boosts confidence
- Assists in anxiety and stress
- Naturally, it is always worth considering discussing an exercise program following a stroke with a doctor or a physiotherapist.

Healthy Diet and Nutrition: Food does influence mood, believe it or not. Eating well promotes the activity of the brain, energy, and sanity [38].

Good mood-boosting foods include

Regular and good eating positively aids the brain in recovering and operating well. A dietitian can develop a plan when it comes to problems of swallowing or appetite following a stroke [39].

Food Group	Examples	Mood-Boosting Benefits
Leafy Greens and Vegetables	Spinach, kale, broccoli, carrots	Keeps the brain healthy as it is full of vitamins and antioxidants
Omega-3-Rich Fish	Salmon, tuna, mackerel	Anti-inflammatory; promoted mood and brain functions
Whole Grains	Brown rice, oats, quinoa	Helps in stabilizing the blood sugar level and energy level. `
Fruits, Nuts and Seeds	Berries, oranges, almonds, and chia seeds	Fiber-rich and such nutrients as magnesium, vitamin B6
Lean Proteins	Chicken, turkey, eggs, lentils	The neurotransmitters and muscle recovery foster.

Table a

Avoid

Mindfulness, Meditation, and Relaxation Techniques

Foods/Substances to Avoid	Examples	Why Avoid (Potential Negative Effects)
Excess Sugar	Candy, pastries, and sweetened drinks	Leads to energy crashes, mood swings, and aggravates inflammation
Junk Food	Fried snacks, fast food, processed meats	Devoid of nutrients, it is a cause of fatigue, weight gain, and poor brain health.
Too Much Caffeine or Alcohol	Coffee (in excess), energy drinks, beer, wine, spirits	Diverts sleep, causes anxiety, and possibly interacts with drugs

Table b

Depression is aggravated by mental tension and anxiety. Through mindfulness and relaxation, the mind will be calm and will have fewer emotional pressures [40].

Examples include

- Deep breath in and out breathing exercises
- Instructed meditation applications or videos
- Progressive muscle relaxation
- Natural physical activity

You can easily experiment with these methods at home, and within even 10 minutes a day, they can produce a rather significant impact on the emotional state.

Occupational and Recreational Therapy: These therapies assist stroke survivors to gain independence and live their normal lives once again, a major aspect in raising their moods and fighting depression.

The packages they provide

- Assistance with the acquisition of daily skills (such as dressing or cooking)
- Confidence building via creative activities or recreational activities
- Modifying hobbies or workplace activities to keep up with the present levels of abilities

Every tiny task can make you feel productive and good, and it can help greatly with feeling helpless and sad.

Involving Caregivers and Loved Ones: Depression is something more than a weapon in the arsenal of the stroke victim: it comes into play in all the people around him or her. This is the reason why the role of caregivers and family members in the treatment process cannot be overestimated.

How they are useful

- Encourage medication and follow-ups for treatment
- Provide love and patience
- Assist in developing a favorable home atmosphere
- Keep in touch with PSD and recovery advice

Dealing with the entire support system, the survivor does not have to feel lonely and is even more inclined to be cured.

Non-Invasive Brain Stimulation

In recent years, neuromodulatory and non-invasive brain stimulation techniques have also been increasingly studied in the treatment of post-stroke depression. Bucur and Papagno (2018) provide a comprehensive overview of the data on these treatment options in a review [45].

Final thoughts on treatment

Not a single treatment fits the treatment of post-stroke depression. A combination of medication, therapy, lifestyle changes, and support tends to work very well most of the time. The most significant thing is not to lose time and be persistent with the program.

PSD is not a vice; it is a medical condition, and it qualifies to be given the necessary attention, care, and love as is the case with any other health condition.

Lifestyle Changes for Managing Post-Stroke Depression (PSD)

Treating post-stroke depression does not only concern drugs or psychology; the process can be strongly influenced by daily lifestyle [41]. Although the healing process is long, you can control it. It is better to make a few steps toward changing your lifestyle, which will allow you to manage the situation, feel happier, and live a better life.

Setting a simple daily routine is one of the first things that may change things. Life following a stroke is usually uncertain. Structure produces a sense of calm in terms of routine. Waking up at the same time every day, working on meal planning, inserting rest stops, and performing simple tasks, even such a mundane thing as dressing or stretching, may restore sanity and comfort to your routine. Routine also lowers stress, and you have something to anticipate.

One more important aspect of recovery is to remain physically active. You do not have to make exercise very intense. It is also important to include short daily walks, light stretching sessions, and physiotherapy exercises to trigger mood improvement through the release of endorphins, the so-called happy hormones. Movement also alleviates stress, enhances sleep, and lifts energy. Any new activity should always be recommended by the doctor or physical therapist and have to be adapted to your concrete condition.

Together with movement, a healthy, properly balanced diet can produce a large impact on mental health as well. Brain-boosting and emotionally stabilizing healthy foods such as fruits, vegetables, whole grains, nuts, and fish contain a lot of vitamins, omega-3 fatty acids, and fiber. Avoid as much as possible junk food, sugary snacks, and caffeine, which may harm mood and performance. In case you have problems with chewing or swallowing since your stroke, there are soft, nutritious meals that a nutritionist can assist with.

Quality sleep is very important both to the mind and the body. Setting a regular sleep routine, a good sleeping environment, and ensuring a minimum amount of screen time before bed can all enhance sleep quality [42]. But when you can't sleep, or can't stay asleep, talk to your doctor, because bad sleeping only increases depression.

Another effective method of combating depression is by maintaining a link with others. Most stroke patients complain of feeling lonely, particularly when they experience speech or movement disability. However, keeping in touch with friends, family, or support groups via a visit, a call, or online chats may help patients ease the feeling of loneliness and bring emotional support. Simple brief conversations, even once a day, are the way you can feel that you are not alone. It also helps to practice mind and stress management techniques.

Mindfulness can be defined as the act of being conscious of the present, and it can be used to stave off the negative thoughts that routinely accompany depression. Clearing your mind and bringing up your mood can be done by simple breathing exercises, guided meditation, some calming music, or writing in a journal for a few minutes a day.

Goals can be set as small and achievable daily to gain confidence. Heavy chores may be intimidating following a stroke, and a simple approach like brushing your hair, pouring a cup of tea, or speaking to a loved one will help get the feeling of achievement. It only takes baby steps, which slowly lead to success and eventually to recovery.

It can also be useful to reduce the triggers of negativity [43]. Depression can be made worse by looking at bad news, comparing yourself with people who recover faster, and performing stressful activities. Instead, you can substitute them with positive habits, such as reading inspirational stories, going to a park, or watching a favorite movie/hobby.

Talking about hobbies, it is worth noting that pursuing even a small, meaningful activity can restore a feeling of joy and meaning. Painting, gardening, listening to music, and even volunteering, as much as possible, will do something that matters to you, and it will restore a sense of identity as well as a sense of self-worth that will be built after the stroke.

Last but not least, the most critical lifestyle change can be the ability to ask for assistance. Depression can make one feel that he or she has to deal with it on his/her own, but support is the key. Do not be afraid to rely on others, whether it is family, friends, doctors, or support groups. It is not a weakness to share your feelings and request help; it is an action of courage, and it is very significant in the healing process.

Prevention of post-stroke depression

Although it is not always possible to avoid post-stroke depression (PSD) completely, some measures can be taken to minimize it and ensure a stable mood and emotional well-being in the process of recovery. Early intervention is the key, both physically and emotionally, to play an important role in avoiding the issue of long-term mental health issues in the future.

Emotional support in the early stages is an excellent prevention strategy. Survivors of a stroke are usually caught by a feeling of ferocity, depression, or perplexity. These emotions can be negotiated in a healthy manner by actually introducing counseling or support groups early into recovery. Family members and caregivers also contribute a lot by providing listening without judgment, encouragement, and being there emotionally.

The second necessary measure is the periodic screening for depression. The mental condition must be monitored regularly along with the physical development. It is also more effective to treat early manifestations of depression because they are caught early, usually with referrals to mental health workers.

Prevention is also supported through remaining mentally and socially active [44]. Loneliness can be overcome by keeping in touch with friends, family, or community groups. Simulating the mind through puzzles, music, or hobbies helps keep the mind alert and helps slow down emotional deterioration.

There is steadiness due to organizing a daily routine. Emotional stability is maintained through a structured program of good sleep, eating, and light exercise. Simple activities such as expending energy through grooming or doing small chores can give a purpose.

Good nutrition and physical exercise are also important. Exercise contributes to the release of hormones that put you in a good mood, whereas a healthy diet enhances the performance of the brain and energy levels.

Lastly, training to cope with stress using one or the other techniques, such as relaxing, meditating, or taking a walk in nature, as well as learning how to stay abreast of the information regarding PSD, will remove the fear and promote early intervention. By providing education to survivors and caregivers, it is possible to empower them and make them aware of the problems sooner or later.

Conclusion

A not-so-well-known aspect of stroke recovery is post-stroke depression, yet it is very prevalent. It is not only depressive effects, but also influences motivation, relationships, and general healing. What is positive is that a proper combination of medical intervention, emotional support, as well as lifestyle changes can help treat PSD. Support from loved ones, early diagnosis, therapy, and medication can play a significant role in the future of the survivor. Good mental health and quality of life can be achieved impressively with simple and daily activities such as regular exercise, healthy consumption, a regular schedule, and social interaction with people. Although not entirely preventable, early recognition of PSD symptoms significantly improves the chances of timely intervention and better recovery outcomes. The entire process of stroke recovery is not solely physical, and recovery of the mind is imperative as well. Sufferers may just come out of depression by patience, consideration, and appropriate interventions, thus going on in life with a new vigor and hope.

Bibliography

1. Guo J., *et al*. "The advances of post-stroke depression: 2021 update". *Journal of Neurology* 269.3 (2021): 1236-1249.

2. Ore ML and Foli KJ. "Reintegration for post-treatment cancer survivors: A concept analysis". *Journal of Holistic Nursing* 38.3 (219): 300-317.

3. Regen F and Benkert O. „Antidepressiva. In: Benkert O, Hippius H. (Hrsg) Kompendium der Psychiatrischen Pharmakotherapie,. 12.. Auflage, Springer; Berlin (2019): S1-208.

4. Ayerbe L., *et al*. "Explanatory factors for the increased mortality of stroke patients with depression". *Neurology* 83 (2014): 2007-2012.

5. Robinson RG. "Post stroke depression: a review". *The American Journal of Psychiatry* 73 (2016): 221-231.

6. Pan A., *et al*. "Depression and the risk of stroke morbidity and mortality: a meta-analysis and systematic review". *JAMA* 306 (2011): 1241-1249.

7. Kapfhammer HP. "Psychische Störungen bei somatischen Krankheiten. in:. Möller HJ, Laux G, Kapfhammer HP. "(Hrsg) Psychiatrie Psychosomatik Psychotherapie. 5. Auflage Springer-Verlag; Berlin (2017): 2693-2805.

8. Robinson RG., *et al*. "Poststroke depression: a review". *The Canadian Journal of Psychiatry* 55 (2010): 141-149.

9. Zhou J., *et al*. "Post-stroke neuropsychiatric complications: Types, pathogenesis, and therapeutic intervention". *Aging and Disease* 14.6 (2023): 2127.

10. Aldrich M. "Employability skills: Economics dissertation at University of East Anglia [Preprint] (2019).

11. Frank D., *et al*. "Pathophysiology and current drug treatments for post-stroke depression: A Review". *International Journal of Molecular Sciences* 23.23 (2022): 15114.

12. De Fruyt J., *et al*. "Anhedonia in depressive disorder: A narrative review". *Psychopathology* 53.5-6 (2020): 274-281.

13. Komsiyska D. "Symptoms of post-stroke depression". *Trakia Journal of Sciences* 17.3 (2019): 235-242.

14. Suksatan W., *et al*. "Burdens among familial caregivers of stroke survivors: A literature review". *Working with Older People* 26.1 (2021): 37-43.

15. Lee D. "Winslow, the anatomy of suicide". *The History of Suicide in England* 1650-1850 (2021): 167-359.

16. Bilek AJ and Richardson D. "Post-stroke delirium and challenges for the rehabilitation setting: A narrative review". *Journal of Stroke and Cerebrovascular Diseases* 32.8 (2023): 107149.

17. Lanctôt KL., *et al*. "Canadian stroke best practice recommendations: Mood, cognition and fatigue following stroke, 6th edition update 2019". *International Journal of Stroke* 15.6 (2019): 668-688.

18. Abuhadra BD., *et al*. "The prevalence of post-traumatic stress disorder, depression, and anxiety in Libya: A systematic review". *Middle East Current Psychiatry* 30.1 (2023).

19. Huang F., *et al*. "Is geriatric depression scale a valid instrument to screen depression in Chinese community-dwelling elderly?" *BMC Geriatrics* 21.1 (2021).

20. Tay J., *et al*. "Apathy after stroke: Diagnosis, mechanisms, consequences, and treatment". *International Journal of Stroke* 16.5 (2021): 510-518.

21. Lanctot KL., *et al*. "Canadian stroke best practice recommodations: mood, cognition and fatigue following stroke, 6th edition update 2019". *International Journal of Stroke* 15 (2020): 688-688.

22. Chen Y., *et al*. "Treatment effects of antidepressants in patients with post stroke depression: a meta analysis". *Annals of Pharmacotherapy* 40 (2006): 2115-2122.

23. Starkstein SE and Hayhow BD. "Treatment of post-stroke depression". *Current Treatment Options in Neurology* 21.07 (2019): 31.

24. Mortensen JK and Andersen G. "Pharmacological management of post-stroke depression: an update of the evidence and clinical guidance". *Expert Opinion on Pharmacotherapy* (2021): 1-10.

25. Deng L., *et al*. "Efficacy and tolerability of pharmacotherapy for post stroke depression: a network meta analysis". *Oncotarget* 9 (2018): 23718-23728.

26. Chen Y., *et al*. "Antidepressant prophylaxis for post stroke depression: a meta analysis In". *Journal of Clinical Psychopharmacology* 22 (2007): 159-166.

27. Chollet F., *et al*. "Fluoxetine for motor recovery after acute ischaemic stroke (FLAME): a randomised placebo-controlled trial". *Lancet Neurology* 10 (2011): 123-130.

28. Mead GE., *et al*. "Selective serotonin reuptake inhibitors (SSRIs) for stroke recovery". *Cochrane Database of Systematic Reviews* 11 (2018).

29. Flaster M., *et al*. "Poststroke depression: a review emphasizing the role of prophylactic treatment and synergy with treatment for motor recovery". *Topics in Stroke Rehabilitation* 20 (2013): 139-150.

30. Cholett F., *et al*. "Use of antidepresant medications to improve outcomes after stroke". *Current Neurology and Neuroscience Reports* 13 (2013): 318.

31. Kraglund KL., *et al*. "Neuroregeneration and vascular protection by citalopram in acute ischemic stroke (TALOS)". *Stroke* 49 (2018): 2568-2576.

32. FOCUS trial Collaboration. Effects of fluoxetine on functional outcomes after acute stroke (FOCUS): a pragmatic, double-blind randomised, controlled trial". *Lancet* 393 (2019): 265-274.

33. Legg LA., *et al*. "Selective serotonin reuptake inhibitors for stroke recovery". *Cochrane Database of Systematic Reviews* (2019): 1 CD 009286.

34. Wijeratne T., *et al*. "A narrative review on the non-pharmacologic interventions in post-stroke depression". *Psychology Research and Behavior Management* 15 (2022): 1689-1706.

35. Costello H., *et al*. "Antidepressant medications in dementia: Evidence and potential mechanisms of treatment-resistance". *Psychological Medicine* 53.3 (2023): 654-667.

36. Zhang S., *et al*. "Neuropsychiatric issues after stroke: Clinical significance and therapeutic implications". *World Journal of Psychiatry* 10.6 (2020): 125-138.

37. Curran J. "Cognitive behavioural therapy for dummies Rob Willson rhena branch cognitive behavioural therapy for dummies john wiley and sons first 334 £14.99 0470018380 0470018380". *Mental Health Practice* 9.9 (2006): 26-26.

38. Constantin ET and Fonseca S. "The effect of food on mental health". *Revista Internacional de Educação, Saúde e Ambiente* 3.2 (2020): 1-17.

39. Izaham A. "Ethical guidelines for Nutritional Management of stroke". *Nutritional Neurosciences* (2024): 319-333.

40. BARIŞ EREN N. "A stress management tool in nurses: Mindfulness- based stress reduction". *Black Sea Journal of Health Science* 6.3 (2023): 508-510.

41. He C., *et al*. "Healthy Lifestyles and Post-Stroke Depression: A NHANES-based cross-sectional study of stroke survivors (2025).

42. Moosavi-Movahedi F and Yousefi R. "Good sleep as an important pillar for a healthy life". *University of Tehran Science and Humanities Series* (2021): 167-195.

43. Sanson M., *et al*. "Trigger warnings are trivially helpful at reducing negative affect, intrusive thoughts, and avoidance". *Clinical Psychological Science* 7.4 (2019): 778-793.

44. Enos G. "Mental Health-Focused Foundation contributes to suicide prevention plan". *Mental Health Weekly* 32.25 (2022): 1-7.

45. Bucur M Papagno CA. "Systematic review of noninvasive brain stimulation for post stroke depression". *Journal of Affective Disorders* 238 (2018): 69-78.