



Contemporary, Novel and Modern Approaches in the Neurorehabilitation of Cerebral Palsy: A Narrative Review

J Afrin Sithara^{1*}, S Cathrine² and MK Franklin Shaju³

¹MPT in Neurology, RVS College of Physiotherapy, The Tamil Nadu Dr. M.G.R. University, India

²Associate Professor, RVS College of Physiotherapy, India

³Principal, RVS College of Physiotherapy, The TamilNadu DR. M.G.R. Medical University, Chennai, India

***Corresponding Author:** J Afrin Sithara, MPT in Neurology, RVS College of Physiotherapy, The Tamil Nadu Dr. M.G.R. University, India.

Received: May 08, 2026

Published: May 31, 2026

© All rights are reserved by J Afrin Sithara, et al.

DOI: 10.31080/ASNE.2026.09.0917

Abstract

Background: Cerebral palsy (CP) is a non-progressive neurodevelopmental disorder characterized by motor, postural, and functional deficits following damage to the developing brain [1]. It has a profound effect on motor function, independence and quality of life. Neurorehabilitation now emphasizes neuroplasticity-based, task-specific, and technology-assisted approaches to improve functional recovery [2].

Objective: This narrative review aims to evaluate the evidence regarding contemporary approaches such as Constraint-Induced Movement Therapy (CIMT), Neurodevelopmental Therapy (NDT), Modern approaches such as Robotic-Assisted Gait Training (RAGT), Virtual Reality (VR), novel approaches such as Dynamic Intensive Therapy (DIT), Action Observation Therapy (AOT) and Motor Imagery (MI) techniques.

Search Strategy: A narrative review of literature published between 2010 and 2025 was conducted using PubMed, Google Scholar, Ovid, and Embase databases. Randomized controlled trials, systematic reviews, meta-analyses, and clinically relevant studies on neurorehabilitation in cerebral palsy were selected based on type of intervention, patient characteristics and clinical outcomes.

Result: A total of 27 studies were included. Overall, the findings indicate that contemporary, modern and novel neurorehabilitation approaches are effective in individuals with cerebral palsy, with each intervention demonstrating benefits in different ways.

Conclusion: Available evidence suggests that CIMT and RAGT are highly effective in enhancing upper limb function and gait performance, respectively. DIT, VR, AOT, and MI demonstrated promising effects on motor learning, patient engagement and neuroplasticity. However, variability in treatment protocols and limited long-term evidence indicate the need for further high-quality research [3-7].

Keywords: Cerebral palsy, neurorehabilitation, constraint-induced movement therapy, robotic-assisted gait training, motor imagery, action observation therapy, virtual reality

Introduction

Cerebral palsy (CP) is an umbrella term for a group of non-progressive, permanent neurological conditions that mainly affect posture, motor function, and movement, resulting from abnormalities in the developing brain of a fetus or infant. It continues to be a major cause of childhood disability, affecting approximately 2-3 children per 1000 live births [1,8]. The clinical presentation of CP varies depending on the location and severity of the brain lesion and may include spasticity, muscle weakness, coordination difficulties, abnormal reflexes, and limitations in activities of daily living [1,8].

Traditionally, treatment approaches for cerebral palsy have focused on passive interventions such as stretching, positioning, and reflex inhibition [9,10]. While these techniques helped to prevent joint stiffness and contractures, they had limited benefits in improving function. As our understanding of the brain has improved, neuroplasticity has emerged as a key principle in neurorehabilitation [2,4]. Neuroplasticity refers to the ability of the brain to reorganize its structure and function in response to training, experience, and environmental stimulation. Recent evidence suggests that individuals with CP achieve better functional outcomes through intensive, goal-directed, and activity-based rehabilitation approaches that encourage active patient participation [11]. As a result, current rehabilitation approaches emphasize evidence-based, activity-oriented and technology-based interventions. Several innovative rehabilitation approaches based on these principles include the following Constraint-Induced Movement Therapy, Neurodevelopmental Therapy, Robotic-Assisted Gait Training, Dynamic Intensive Therapy, Action Observation Therapy (AOT), Motor Imagery and Virtual Reality rehabilitation. Although these approaches vary in their mechanisms and methods of application, they all have a common goal of improving motor function, improving participation and quality of life in individuals with cerebral palsy [3-5,7,9].

This narrative review discusses these contemporary rehabilitation strategies with emphasis on their underlying principles, clinical application, target population, and evidence regarding their effectiveness.

Methodology

This narrative review was conducted to evaluate contemporary, modern, and emerging neurorehabilitation approaches used in the management of cerebral palsy. Electronic databases

including PubMed, Google Scholar, Ovid, and Embase were searched for relevant literature published between January 2010 and October 2025. The search strategy included keywords such as “cerebral palsy,” “neurorehabilitation,” “constraint-induced movement therapy,” “robot-assisted gait training,” “virtual reality rehabilitation,” “motor imagery,” and “action observation therapy,” combined using Boolean operators (AND, OR) to identify relevant studies.

Randomized controlled trials, systematic reviews, meta-analyses, and clinically relevant studies focusing on neurorehabilitation interventions in cerebral palsy were considered for inclusion. Studies unrelated to cerebral palsy rehabilitation, non-neurological studies, conference abstracts, and articles without accessible full text were excluded.

A total of 74 articles were initially identified through database searching. After removal of duplicates and exclusion of unrelated or non-full-text articles, a total of 27 studies were included in this final review based on their relevance to emerging and evidence-based neurorehabilitation approaches in cerebral palsy. The selected studies were analysed based on intervention type, study design, participant characteristics, and reported clinical outcomes, including motor function, gait, balance, coordination, and functional independence. Particular emphasis was placed on interventions aimed at improving functional recovery, movement performance, and rehabilitation outcomes in children with cerebral palsy.

Intervention

Constraint-induced movement therapy (CIMT)

Constraint-Induced Movement Therapy is a widely recognized rehabilitation approach used to improve upper limb function in children with hemiplegic cerebral palsy. The approach is based on the concept of “learned non-use”, in which reduced use of the affected limb results in further disuse and decreased cortical representation [3,12]. CIMT promotes the use of the affected limb by restricting the sound limb by using mitts, slings, or casts during therapeutic activities. CIMT is based on experience-dependent neuroplasticity, where repetitive task practice strengthens neural connections and enhances cortical reorganization. CIMT protocols commonly focus on functional tasks such as reaching, grasping, object manipulation, eating, and dressing [3,12,13]. There is robust research evidence on the effectiveness of CIMT, including many randomized controlled trials and reviews [13]. Several

studies demonstrated improvements in upper limb function, bimanual coordination, and independence in activities of daily living following CIMT interventions [3,12,13]. However, CIMT may require high-intensity training and prolonged therapy duration, which can affect patient compliance and accessibility [13,27].

Neurodevelopmental therapy (NDT)

Neurodevelopmental Therapy, also known as the Bobath approach, is a well-established rehabilitation approach used in the management of cerebral palsy [9,10]. It is based on the principles of facilitating normal movement patterns while inhibiting abnormal postures and reflex activity through therapeutic handling techniques [11]. The fundamental concept of NDT is that abnormal muscle tone and impaired postural control interfere with normal movement patterns. Through appropriate sensory cues and postural alignment, therapists aim to facilitate normal motor responses and improve functional movement [6]. NDT primarily focuses on postural control, alignment and movement patterns via manual facilitation techniques. NDT has broad clinical application and can be used in various types of cerebral palsy, including spastic, dyskinetic, and ataxic forms, across different age groups, including early intervention programs. NDT is often combined with other rehabilitation interventions as part of a multidisciplinary treatment approach.

Robotic-assisted gait training (RAGT)

Robotic-Assisted Gait Training is an advanced neurorehabilitation approach that provides repetitive and controlled gait training for individuals with cerebral palsy [4]. It involves the use of robotic systems, such as treadmill-mounted exoskeleton devices, to facilitate controlled walking patterns [14]. RAGT enables intensive, repetitive, and task-specific gait training by providing controlled movement patterns with adjustable parameters such as speed, step length, and body-weight support. These features contribute to motor learning and functional recovery by promoting repetitive practice and neuroplastic adaptation [4,15]. Randomized controlled trials and systematic reviews support the use of RAGT for improving gait speed, endurance, walking symmetry, and functional mobility in individuals with cerebral palsy [14-16]. Better outcomes are generally observed in individuals receiving intensive and longer-duration training programs. However, RAGT has certain limitations, including high equipment costs, limited availability in clinical settings, and the requirement

for skilled professionals [17]. Despite these limitations, RAGT continues to emerge as a promising intervention for improving walking ability in individuals with cerebral palsy.

Dynamic intensive therapy (DIT)

Dynamic Intensive Therapy (DIT), also known as intensive therapy or suit-based therapy, is a high-frequency, high-intensity rehabilitation approach that aims at improving motor function in individuals with cerebral palsy. It is based on the principle that intensive and repetitive practice may enhance neuroplasticity in the developing brain [7,18]. DIT emphasizes repetitive functional movements, task-specific training, and active patient participation. Some approaches involve the use of therapeutic suits, such as the Thera Suit or Adeli Suit, to provide external supports, enhance proprioceptive input and improve postural alignment during therapeutic activities [19]. The intervention commonly includes strengthening, balance training, gait practice, and functional task performance [18]. This approach is particularly suitable for children with moderate levels of cerebral palsy who are able to actively participate in intensive training sessions. However, DIT has certain limitations, including the lack of standardized treatment protocols and variability in clinical outcomes across studies [7]. In Addition, the effectiveness of DIT often depends on high patient motivation and significant therapist involvement.

Despite these challenges, DIT remains a promising intervention because of its strong emphasis on intensive practice and functional rehabilitation [19].

Action observation therapy (AOT)

Action Observation Therapy is an emerging neurorehabilitation technique that utilizes the mirror neuron system, which is activated during both observation and performance of goal-directed movements [6,20]. This mechanism plays a vital role in motor learning and cortical reorganization. In AOT, patients observe specific motor tasks such as reaching and grasping movements, followed by active imitation of the observed actions [20]. This process helps stimulate the motor cortex and movement execution. AOT is particularly beneficial for children with impaired motor planning and individuals who have difficulty in initiating voluntary movements. It is often combined with conventional physiotherapy to enhance treatment outcomes. One of the major advantages of AOT is that it is simple, cost-effective, and easy to

implement in clinical and home settings, and it does not require advanced equipment.

Motor imagery (MI)

Motor Imagery is a cognitive process in which an individual mentally rehearses a movement without actually performing it [21]. It activates neural pathways similar to those involved in actual movement, and it may facilitate motor learning and improve cortical connectivity. MI stimulates several motor cortical areas, including primary motor cortex, premotor cortex and supplementary motor area. This neural activation promotes synaptic plasticity and improves motor planning .MI is particularly beneficial for individuals with limited physical ability and those unable to perform active movements because of weakness or spasticity. It is often used as an adjunct to conventional physiotherapy to enhance overall rehabilitation outcomes. Evidence suggests that MI may improve the motor planning, coordination and functional performance, especially when combined with active physical training [21].

Virtual reality (VR) rehabilitation

Virtual Reality is a technology-assisted rehabilitation approach that uses computer-generated simulations to perform functional activities in an interactive environment [5]. It enhances patient motivation and engagement through immersive therapeutic experiences [22]. VR-based rehabilitation provides real-time visual and auditory feedback, along with task-specific and repetitive practice. These features contribute to motor learning, patient participation and neuroplastic adaptation. VR is particularly effective in children with cerebral palsy because of its game-based environment, high levels of engagement and improved treatment compliance [23]. VR rehabilitation is commonly used to improve balance and coordination, upper limb function, and functional mobility. Several studies have demonstrated improvements in motor function, balance and coordination following VR-based interventions in individuals with cerebral palsy. In addition, VR provides a safe and controlled rehabilitation environment and allows individualized treatment programs. However, certain limitations exist, including the need for technological resources and limited accessibility in low- resource clinical settings [24]. Despite these limitations, VR continues to gain importance as an innovative and effective rehabilitation tool in cerebral palsy management [5].

| Author and Year | Study Design | Intervention | Sample Characteristics | Main Findings |
|-------------------------------|-----------------------------|--------------------------------------------------------------------|-------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|
| Bailes AF, <i>et al.</i> 2010 | Intensive therapy study | Suit-wear intensive therapy | Children with cerebral palsy: ambulatory participants with motor impairments. | Improved gross motor function following intensive suit-based therapy. |
| Buccino, <i>et al.</i> 2012 | Pilot study | Action observation therapy | Children with cerebral palsy with upper limb motor impairment. | AOT improved upper limb motor function and functional activity performance in children with cerebral palsy |
| DeLuca, <i>et al.</i> 2012 | Clinical intervention study | Constraint-Induced Movement Therapy (CIMT) | Young children with cerebral palsy and upper limb impairment. | Children tolerated therapy dosage well and showed functional improvement. |
| Dewar, <i>et al.</i> 2015 | Systematic review | Exercise interventions for postural control | Children with cerebral palsy with postural impairment. | Moderate evidence supports exercise interventions for improving postural control. |
| Jung, <i>et al.</i> 2018 | Case series | Virtual Reality training using Xbox Kinect | Children with spastic cerebral palsy with balance impairment. | Improvement noted in balance and physical functioning after VR training. |
| Sah, <i>et al.</i> 2019 | Randomized clinical trial | Task-oriented activities based on Neurodevelopmental Therapy (NDT) | Children with cerebral palsy with trunk control and balance impairment. | Improved trunk control, balance, and gross motor function. |

| | | | | |
|---------------------------------------|-------------------------------------|--------------------------------------------|---------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------|
| Hoare., <i>et al.</i> 2019 | Randomized controlled trial | Constraint-Induced Movement Therapy (CIMT) | Children with unilateral cerebral palsy with upper limb impairment. | CIMT showed better upper limb outcomes compared to low-dose therapy. |
| Karadag-Saygi., <i>et al.</i> 2019 | Systematic review | Suit therapies | Children with cerebral palsy with motor and postural impairment. | Intensive task-specific suit therapy showed promising results. |
| Souto., <i>et al.</i> 2020 | Cross sectional study | Motor imagery | Children with cerebral palsy and typically developing children. | Children with CP demonstrated motor imagery ability, but performance was slower and less accurate than typically developing children. |
| Vitrikas., <i>et al.</i> 2020 | Review article | Overview of cerebral palsy management. | Children with cerebral palsy. | Discussed clinical presentation, diagnosis, and multidisciplinary management approaches in CP. |
| Pool., <i>et al.</i> 2021 | Clinical intervention study | Robotic-Assisted Gait Training (RAGT) | Children with cerebral palsy with gait impairment. | Participants completed intervention safely with functional gait improvement. |
| Ramey., <i>et al.</i> 2021 | Randomized trial | Constraint-Induced Movement Therapy (CIMT) | Children with unilateral cerebral palsy | High-dose CIMT produced greater improvement than moderate-dose therapy |
| Montoro-Cárdenas., <i>et al.</i> 2021 | Systematic review and meta-analysis | Nintendo Wii Balance Board therapy | Children with cerebral palsy with balance impairment | Positive effects observed on dynamic balance |
| Belizón-Bravo., <i>et al.</i> 2021 | Systematic review | Dynamic suit orthoses | Children with cerebral palsy with postural and gait impairment | Improved gait, balance, and functional outcomes |
| Moll., <i>et al.</i> 2022 | Randomized controlled trial | Robot-Assisted Gait Training | Pediatric patients with cerebral palsy with gait impairment | Improvement observed in walking speed and gait performance |
| Cortés-Pérez., <i>et al.</i> 2022 | Systematic review and meta-analysis | Robot-Assisted Gait Therapy | Children with cerebral palsy with gait impairment. | RAGT demonstrated better post-intervention outcomes than conventional therapy. |
| Ungureanu., <i>et al.</i> 2022 | Pilot study | Bobath and Vojta methods | Children with cerebral palsy with motor impairment. | Improvement in symmetry and postural facilitation was observed. |
| Volpini., <i>et al.</i> 2022 | Systematic review and meta-analysis | Assisted robotic gait training | Individuals with cerebral palsy with gait impairment. | Walking distance improved significantly following RAGT. |
| Te Velde., <i>et al.</i> 2022 | Meta-analysis | Neurodevelopmental Therapy (NDT) | Children with cerebral palsy with motor impairment. | Moderate effect favoring activity-based rehabilitation approaches. |

| | | | | |
|-----------------------------------|---------------------------------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------|----------------------------------------------------------------------------------------|
| Khanna, <i>et al.</i> 2023 | Literature review | Neurodevelopmental treatment | Children with cerebral palsy with motor impairment. | Further large-scale studies are recommended to confirm effectiveness. |
| Klobucká, <i>et al.</i> 2023 | Cost-effectiveness analysis | Robotic-Assisted Gait Training (RAGT) | Children with cerebral palsy with motor impairment | RAGT was effective but associated with high treatment cost. |
| Julien, <i>et al.</i> 2024 | Clinical intervention study | Robotic-Assisted Gait Training (RAGT) | Children with cerebral palsy with gait impairment. | Significant improvement observed in gait performance. |
| Ziab, <i>et al.</i> 2024 | Double-blinded randomized controlled trial | Overground gait training compared with balance-specific and conventional training | Children with cerebral palsy with gait and balance impairment | Improvement observed in GMFM scores and goal attainment scales. |
| Takes, <i>et al.</i> 2025 | Randomized comparative study | Robot-assisted gait training | Children with cerebral palsy with gait impairment. | Combined conventional therapy and RAGT showed functional gait improvement. |
| Martin-Moreno, <i>et al.</i> 2025 | Systematic review and meta-analysis of follow-ups | Constraint-induced movement therapy and bimanual training | Children with cerebral palsy with upper limb impairment. | Improved upper limb function: further long-term studies needed. |
| Cortes-Perez, <i>et al.</i> 2025 | Systematic review and meta-analysis | Dual task training interventions | Individuals with cerebral palsy with balance and walking | It improved postural balance and walking speed. |
| Perez-Carcamo, <i>et al.</i> 2025 | Systematic review and meta-analysis | Non-immersive virtual reality rehabilitation | Children with cerebral palsy with gross motor and balance impairment. | VR rehabilitation improved gross motor function, balance, and functional independence. |

Table 1: Summary of contemporary, novel, and modern neurorehabilitation approaches in cerebral palsy.

Discussion

This narrative review outlines current trends in neurorehabilitation for cerebral palsy, highlighting a shift from conventional passive treatment to more active, intensive, and neuroplastic-based approaches [2,4]. Neuroplasticity refers to the ability of the brain to reorganize neural pathways in response to repeated practice, motor learning, and environmental stimulation [2]. Contemporary rehabilitation approaches in cerebral palsy are increasingly designed to utilize neuroplastic mechanisms through repetitive, task-specific, and intensive training, thereby improving motor control and functional independence. This review highlights the variable effectiveness of different types of rehabilitation approaches based on their mechanisms and implementation. Although all interventions demonstrated beneficial effects,

their effectiveness varied depending on treatment goals and patient characteristics. Constraint-Induced Movement Therapy showed stronger evidence for improving upper limb function [12], whereas Robotic-Assisted Gait Training was more effective for improving gait symmetry, walking endurance, and mobility outcomes [16]. Neurodevelopmental Therapy remains widely used in clinical practice, however, recent evidence suggests that task-oriented and intensive approaches may produce greater functional improvements in some children with cerebral palsy [10]. Virtual Reality rehabilitation also plays a role by enhancing engagement and compliance through interactive and motivating environments [24]. The provision of real-time feedback improves motor learning and function. However, interventions such as robotic-assisted therapy and virtual reality rehabilitation may have

limited accessibility in low-resource clinical settings because of high equipment costs, the need for specialized training, and limited technological availability. In contrast, approaches such as CIMT and task-oriented therapy are comparatively more feasible and cost-effective for broader clinical implementation. The selection of rehabilitation strategies should be individualized according to the severity of impairment, functional goals, age, cognitive status, and availability of rehabilitation resources.

Novel interventions, such as Dynamic Intensive Therapy, Action Observation Therapy, and Motor Imagery have shown encouraging outcomes by promoting motor learning, cortical activation and functional outcomes [19-21]. These approaches demonstrate the importance of integrating cognitive and physical components of rehabilitation for better results.

Limitations

Although there is increasing evidence supporting contemporary neurorehabilitation strategies to improve motor function in cerebral palsy, several limitations remain. Few studies assessed new therapeutic interventions such as Motor Imagery and Action Observation Therapy, limiting the evidence. Lack of treatment protocols for each intervention leads to inconsistencies in clinical effects [7]. Many studies have small sample sizes, which could influence the reliability and generalizability of the results [19]. Differences in the duration, intensity and characteristics of the patient populations make it challenging to develop standardized clinical recommendations. Additionally, variability in intervention intensity, duration, outcome measures, and participant characteristics across studies limits direct comparisons between rehabilitation approaches. Another limitation was the use of different outcome measures across the included studies. Various tools were used to assess gait, balance, motor function, and functional independence, this heterogeneity in assessment tools made comparison of intervention effectiveness across studies more challenging.

Conclusion

The present review highlights the recent shift in the rehabilitation of cerebral palsy toward neuroplasticity-driven, task-specific, and intensive therapies [2]. The most robust evidence for enhancing upper limb function and walking ability comes from Constraint-Induced Movement Therapy and Robotic-Assisted Gait Training,

respectively. Novel approaches such as Action Observation Therapy, Motor Imagery, and Virtual Reality have potential to improve motor learning, enhance cortical activation, and increase engagement, but more research is needed to determine their long-term efficacy [20,21,24,25]. Future studies should focus on long-term follow-up, standardized treatment protocols, and larger multicenter trials to better understand the sustainability of therapeutic outcomes across cerebral palsy rehabilitation approaches [26,27].

The results suggest that repetition, intensity and active engagement are critical elements to successful rehabilitation interventions. In summary, a multimodal, personalized approach is essential in the rehabilitation of individuals with cerebral palsy. Additional studies are needed to establish protocols, and assess long-term outcomes. Although these approaches differ in their mechanisms and methods of application, these interventions collectively aim to enhance motor function, promote participation, and improve the quality of life in individuals with cerebral palsy [4].

Bibliography

1. Wimalasundera N and Stevenson VL. "Cerebral palsy". *Practice Neurology* 16.3(2016): 184-194.v
2. Dewar R., et al. "Exercise interventions improve postural control in children with cerebral palsy: a systematic review". *Developmental Medicine & Child Neurology* 57.6 (2015): 504-520.
3. DeLuca SC., et al. "Constraint-induced movement therapy (CIMT) for young children with cerebral palsy: effects of therapeutic dosage". *Journal of Pediatric Rehabilitation Medicine* 5.2 (2012): 133-142.
4. Pool D., et al. "Locomotor and robotic assistive gait training for children with cerebral palsy". *Developmental Medicine & Child Neurology* 63.3 (2021): 328-335.
5. Jung SH., et al. "Does virtual reality training using the Xbox Kinect have a positive effect on physical functioning in children with spastic cerebral palsy? A case series". *Journal of Pediatric Rehabilitation Medicine* 11.2 (2018): 95-101.
6. Sah AK., et al. "Effects of Task-oriented Activities Based on Neurodevelopmental Therapy Principles on Trunk Control, Balance, and Gross Motor Function in Children with Spastic Diplegic Cerebral Palsy: A Single-blinded Randomized Clinical Trial". *Journal of Pediatric Neuroscience* 14.3 (2019): 120-126.

7. Karadağ-Saygı E and Giray E. "The clinical aspects and effectiveness of suit therapies for cerebral palsy: A systematic review". *Turkish Journal of Physical Medicine and Rehabilitation* 65.1 (2019): 93-110.
8. Vitrikas K., et al. "Cerebral Palsy: An Overview". *American Family Physician* 101.4 (2020): 213-220.
9. Te Velde A., et al. "Neurodevelopmental Therapy for Cerebral Palsy: A Meta-analysis". *Pediatrics* 149.6 (2022): e2021055061.
10. Khanna S., et al. "Neurodevelopmental Treatment in Children With Cerebral Palsy: A Review of the Literature". *Cureus* 15.12 (2023): e50389.
11. Ungureanu A., et al. "Balance Rehabilitation Approach by Bobath and Vojta Methods in Cerebral Palsy: A Pilot Study". *Children* 9.10 (2022): 1481.
12. Ramey SL., et al. "Constraint-Induced Movement Therapy for Cerebral Palsy: A Randomized Trial". *Pediatrics* 148.5 (2021): e2020033878.
13. Hoare BJ., et al. "Constraint-induced movement therapy in children with unilateral cerebral palsy". *Cochrane Database System Review* 4.4(2019): CD004149.
14. Volpini M., et al. "Clinical effects of assisted robotic gait training in walking distance, speed, and functionality are maintained over the long term in individuals with cerebral palsy: a systematic review and meta-analysis". *Disability and Rehabilitation* 44.19(2022): 5418-5428.
15. Cortés-Pérez I., et al. "Efficacy of Robot-Assisted Gait Therapy Compared to Conventional Therapy or Treadmill Training in Children with Cerebral Palsy: A Systematic Review with Meta-Analysis". *Sensors* 22.24(2022): 9910.
16. Moll F., et al. "Use of Robot-Assisted Gait Training in Pediatric Patients with Cerebral Palsy in an Inpatient Setting-A Randomized Controlled Trial". *Sensors* 22.24 (2022): 9946.
17. Klobucká S., et al. "Cost-effectiveness analysis of robot-assisted gait training in patients with bilateral spastic cerebral palsy". *Cost-Effective Resource Allocation* 21.1 (2023): 60.
18. Bailes AF., et al. "Changes in two children with cerebral palsy after intensive suit therapy: a case report". *Pediatric Physical Therapy* 22.1 (2010): 76-85.
19. Belizón-Bravo N., et al. "Effects of Dynamic Suit Orthoses on the Spatio-Temporal Gait Parameters in Children with Cerebral Palsy: A Systematic Review". *Children* 8.11 (2021): 1016.
20. Buccino G., et al. "Improving upper limb motor functions through action observation treatment: a pilot study in children with cerebral palsy". *Developmental Medicine & Child Neurology* 54.9 (2012): 822-828.
21. Souto DO., et al. "Motor imagery in children with unilateral cerebral palsy: a case-control study". *Developmental Medicine & Child Neurology* 62.12 (2020): 1396-1405.
22. Montoro-Cárdenas D., et al. "Nintendo Wii Balance Board therapy for postural control in children with cerebral palsy: a systematic review and meta-analysis". *Developmental Medicine & Child Neurology* 63.11 (2021): 1262-1275.
23. Ziab H., et al. "Effectiveness of virtual reality training compared to balance-specific training and conventional training on balance and gross motor functions of children with cerebral palsy: A double blinded randomized controlled trial". *Journal of Pediatric Rehabilitation Medicine* 17.3 (2024): 353-368.
24. Perez-Carcamo J., et al. "Effectiveness of Non-Immersive Virtual Reality on Gross Motor Function, Balance, and Functional Independence in Children with Cerebral Palsy: A Systematic Review with Meta-Analysis". *Journal of Clinical Medicine* 14.21 (2025): 7582.
25. Cortés-Pérez I., et al. "Dual-Task Training Interventions for Cerebral Palsy: A Systematic Review and Meta-Analysis of Effects on Postural Balance and Walking Speed". *Medicina (Mex)* 61.8 (2025): 1415.
26. Tekeş SEÖ., et al. "Robot-assisted gait training in children with cerebral palsy: a randomized comparative study". *Developmental Neurorehabilitation* 28.4-5 (2025): 163-72.
27. Martin-Moreno G., et al. "Constraint-Induced Movement Therapy Versus Bimanual Training to Improve Upper Limb Function in Cerebral Palsy: A Systematic Review and Meta-Analysis of Follow-Ups". *Children* 12(6 (2025): 804.