



The Use of Enzymes in Poultry Diets trends and Approaches, A Minimum Review

Youssef A Attia^{1,2*}

¹Arid Land Agriculture Department, Faculty of Meteorology, Environment and Arid Land Agriculture, King Abdulaziz University, Saudi Arabia

²Animal and Poultry Production Department, Faculty of Agriculture, Damanhour University, Damanhour, Egypt

***Corresponding Author:** Youssef A Attia, Arid Land Agriculture Department, Faculty of Meteorology, Environment and Arid Land Agriculture, King Abdulaziz University, Saudi Arabia and Animal and Poultry Production Department, Faculty of Agriculture, Damanhour University, Damanhour, Egypt.

Received: May 10, 2019; **Published:** June 11, 2019

Abstract

Approximately 50 years ago, enzymes industry was established to meet different demands from detergents to feeds and foods. Enzymes play an important role in improving the utilization of feed, especially feed materials containing antinutritional factors and high proportion of fiber and undigested materials, especially in young and chickens. Recent research has shown confirmed results in this area, especially with the current trend of using enzymes to improve the utilization of low-nutrient feed materials, particularly when using agro-by products in poultry feed [1-4]. The results suggest that the addition of commercial enzymes for fodder containing agricultural residues such as the by-products of rice (broken rice, rice hulls, rice bran and rice polishing) the age of wheat or the products of distillation of dry grains have improved the benefits of feed, growth recipes and egg production [5-7]. However, in general, results showed that enzymes effectiveness are affected by strain of chickens as enzymes induced pronounced effect on younger ages and on broilers than old chickens and layers, type and composition of diets, dose and type of enzymes and nature of feedstuffs [8]. The addition of enzymes to layer chicken's diets containing maize and soybean gain did not affect egg production and its properties and the activity of pancreatic enzymes [9-15].

Keywords: Enzymes; Poultry; Diet

Recent trend in enzymes utilization

The fluctuation in the prices of feed materials always leads to the search for suitable alternatives to use instead of the traditional feed materials, especially with the recent rise in prices due to increased demand and competition between humans and animals on these sources, giving absolute primacy to meet food needs [7,9,10]. The aim of using non-traditional feed is to reduce production costs. However, the rate of replacement depends on its availability easily and easily in markets or the surrounding environment with attention and caution from the presence of anti-feeding and chemical composition of feed materials. Many non-nutritional substances and sugars found in plant components of feed materials negatively affect the health of the intestines and poultry performance [4,7,16-18]. Non-starch polysaccharides (NSP) are important components of plant cell wall for feed which can't be digested by poultry because of their structural nature and their resistance to hydrolysis in the digestive system [3,19]. In some feeds, the cell wall is difficult to digest, especially fiber,

in addition to reducing the feed energy, which is the main cause of the low nutritional values. The use of these feedstocks is then restricted by NSP contents [7].

This occurs mainly when the fibers are soluble, which has a great ability to absorb water and form a gel in the intestine. This increased viscosity of the food bubble leads to slow passage of food along the digestive system, as well as hinder the efficiency of digestive enzymes and reduce the action and absorption rate of food [20]. Moreover, increased viscosity of food bolus interferes with microorganisms (promotes the growth of harmful organisms because they are a suitable food for them). Generally, enzymes that are able to digest xylans and β -glucan, and many noncancerous sugars make it possible to be used in foods and feeds [20].

The main enzymes used in animal feed, especially stomach monocytes, are protease, amylase, lipase, phytase, as well as enzymes for many noncancerous sugars and cellulose. The newly published results recommended using enzymes in animal feed to

improve the utilization of feed materials, particularly those rich in non-starch polysaccharides or containing phytic acid [3,8-10,20]. Enzymes cocktail (multienzymes) are usually a combination of several various enzymes that are active on a different variety of feed materials. Enzymes that have established efficient in animal feed include xylanase, β -glucanase, arabinoxylanase, cellulose, and phytase [7,8,21-23].

Mode of action

The objectives of using enzymes as food supplements include (a) increasing the amount of enzymes available to the animal [19]; (b) mitigating the negative influences of anti-nutritional factors, such as β glucan and arabinoxylans [3]; and (c) increasing availability of nutrients and energy and enhancing their absorption from feed components [8,21]. And (d) alter the microbial environment of the gut by raising beneficial organisms at the expense of harmful organisms [6,24-26].

The external enzymes xylanases or exoxylanase improve the availability of complex nutrients. For example, endoxylanase or exoxylanase enzymes work on analyzing xylan units, which are the backbone units of arabinoxylan, into smaller molecules, which produce many valuable effects and make xylose units more disposable [6,27]. They also increase the prevalence of nutrients and their interaction with internal enzymes [3,20]. Endoxylanase enzymes release nutrient constituents of the plant cell wall and are not susceptible to digestion by monogastrics [28,29]. This group of enzymes could reduce the spread of harmful microorganisms in the small intestine as a result of increased rate of passage swallowing [30] and thus improve the utilization of food by reducing competition between beneficial and harmful intestinal microflora and tolerance to heat stress [31] and reduce blood plasma cholesterol [5].

Conclusion

Enzymes is a useful tool to improve feed utilization, animal performance, immunity through improve gut health and economic efficiency and the mode of action may include reduce gut viscosity of viscous grain, digestibility of nutrients, utilization of nutrients and overcome the anti-nutritional substances.

Bibliography

1. Attia YA., et al "The effect of supplementation of enzyme on laying and reproductive performance in Japanese Quail hens fed nigella seed meal". *The Journal of Poultry Science* 45.2 (2008): 110-115.
2. Lee SA., et al. "Age and dietary xylanase supplementation affects ileal sugar residues and short chain fatty acid concentration in the ileum and caecum of broiler chickens". *Journal of Animal Feed Science and Technology* 234 (2017): 29-42.
3. Aftab U and MR Bedford. "The use of NSP enzymes in poultry nutrition: myths and realities". *World's Poultry Science Journal* 74.2 (2018): 277-286.
4. Alsaffar AA., et al "Productive and reproductive performance and egg quality of laying hens fed diets containing different levels of date pits with enzyme supplementations". *Tropical Animal health and Production* 45.1 (2013): 327-334.
5. Al-Harhi MA. "The Effect of Olive Cake, with or Without Enzymes Supplementation, on Growth Performance, Carcass Characteristics, Lymphoid Organs and Lipid Metabolism of Broiler Chickens". *Revista Brasileira de Ciência Avícola* 19 (2017): 83-90.
6. Al-Harhi MA., et al. "The effects of autoclaving or/and multi-enzymes complex supplementation on performance, egg quality and profitability of laying hens fed whole Prosopis juliflora pods meal diet". *European Poultry Science* 82 (2018): 22.
7. Attia YA., et al. "Effect of zinc bacitracin and phytase on growth performance, nutrient digestibility, carcass and meat traits of broilers". *Journal of Animal Physiology and Animal Nutrition* 100.3 (2016): 485-491.
8. Attia Y A., et al. "Utilization of vegetable diets containing different levels of rice bran with or without commercial enzymes in Norfa laying hen diets". *The Journal of Agricultural Science* 26 (2001): 3557-3577.
9. Attia YA., et al. "Value of rice bran, its maximal utilization and upgrading by phytase and other enzymes and diet- formulation based on available amino acid for broiler chicks". *Archiv Für Geflügelkunde* 67.4 (2003): 157-166.
10. Attia YA and SA Abd El-Rahman. "Impact of multienzymes or Yea Sacc supplementation on growth performance and some carcass parameters of broiler chicks fed triticale containing-diets". *Archiv Für Geflügelkunde* 65.4 (2001): 168-177.
11. Jaroni D., et al. "The effect of dietary wheat middlings and enzyme supplementation. 1. Late egg production efficiency, egg yields, and egg composition in two strains of Leghorn hens". *Poultry Science* 78.6 (1999): 841-847.

12. Flores-Cervantes, S., *et al.* "Use of multienzymatic mix on sorghum-soybean mealbased ration on hen performance". *Italian Journal of Animal Science* 10.2 (2011): e25.
13. Zarei M., *et al.* "Productive performance of laying hens fed wheat-based diets included olive pulp with or without a commercial enzyme product". *African Journal of Biotechnology* 10 (2011): 4303-4312.
14. Wen C., *et al.* "Effect of enzymes preparation on egg production, nutrient retention, digestive enzymes activities and pancreatic enzyme messenger RNA expression of late-phase laying hens". *Animal feed Science and Technology* 172 (2012): 180-186.
15. Sindhu AA., *et al.* "Agro-Industrial by-product as a potential source of livestock feed". *International Journal of Agriculture and Biology* 4.2 (2002): 307-310.
16. Neto RM., *et al.* "Productive performance, intestinal morphology and carcass yield of broilers fed conventional and alternative diets containing commercial enzymatic complex". *International Journal of Poultry Science* 11(2012): 505-516.
17. Jia W., *et al.* "Effects of diet type and enzyme addition on growth performance and gut health of broiler chickens during subclinical *Clostridium perfringens* challenge". *Poultry Science* 88.1 (2009): 132-140.
18. Ramos LSN., *et al.* "Use of exogenous enzymes in diets of broilers chickens". *Revista científica de produção animal* 9 (2007): 84-94.
19. Cozannet P., *et al.* "Next-generation non-starch polysaccharide-degrading, multi-carbohydrase complex rich in xylanase and arabinofuranosidase to enhance broiler feed digestibility". *Poultry Science* 96.8 (2017): 2743-2750.
20. Choct M. "Enzymes for the feed industry: past, present and future". *World's Poultry Science Journal* 62 (2006): 5-16.
21. Odetallah NH., *et al.* "Keratinase in starter diets improves growth of broiler chicks". *Poultry Science* 82.4 (2003): 664-670.
22. Santos AA., *et al.* "Dietary pentosanase supplementation of diets containing different qualities of wheat on growth performance and metabolizable energy of turkey poults". *International Journal of Poultry Science* 3.1 (2004): 465-468.
23. Brufau J., *et al.* "The use of enzymes to improve cereal diets for animal feeding". *Journal of the Science of Food and Agriculture* 86 (2006): 1705-1713.
24. Engberg RM., *et al.* "Influence of whole wheat and xylanase on broiler performance and microbial composition and activity in the digestive tract". *Poultry Science* 83.6 (2004): 925-938.
25. Hogberg A and Lindberg JE. "Influence of cereal non-starch polysaccharides and enzyme supplementation on digestion site and gut environment in weaned piglets". *Animal Feed Science and Technology* 116.1-2 (2004): 113-128.
26. Wang ZR., *et al.* "Effects of enzyme supplementation on performance, nutrient digestibility, gastrointestinal morphology, and volatile fatty acid profiles in the hindgut of broilers fed wheat-based diets". *Poultry Science* 84.6 (2005): 875- 881.
27. Odetallah NH. "The use of dietary enzymes to alleviate enteric disorders of turkeys". Ph.D. Thesis, North Carolina State University (2000): 197.
28. Bedford MR. "Exogenous enzymes in monogastric nutrition: Their current value and future benefits". *Animal Feed Science and Technology* 86.1-3 (2000): 1-13.
29. Lee SY., *et al.* "Effects of multiple enzyme (Rovabio Max) containing carbohydrases and phytase on growth performance and intestinal viscosity in broiler chicks fed corn-wheat-soybean meal based diets". *Asian-Australasian Journal of Animal Science* 23.9 (2010): 1198-1204.
30. Choct M., *et al.* "A comparison of three xylanases on the nutritive value of two wheat's for broiler chickens". *British Journal of Nutrition* 92.1 (2004): 53-61.
31. Attia YA. "Responses of broiler chicks raised under constant relatively high ambient temperature to enzymes, amino acid supplementations, or diet density". *Archiv Für Geflügelkunde* 70.2 (2006): 80-91.

Volume 3 Issue 7 July 2019

© All rights are reserved by Youssef A Attia.