

Mineralization is also greatest when soil moisture is near field capacity and declines with soil drying [9]. Linn and Doran [13] showed that in moist soils microbial activity is optimum when soil water-filled pore space (WFPS) is near 60%, resulting in maximum organic matter decomposition. Anaerobic conditions tend to occur at a WFPS greater than 60%, thereby slowing the decomposition process. Knoepp and Swank [14] found that there was a significant interaction between moisture content and soil temperature in N mineralization. Wetting and drying cycles have an influence on microbial activity, there by affecting decomposition of SOM [15-17]. Drying followed by rapid rewetting cycles generally causes an increase in organic substrate available for microbial attack [15]. The labile substrates that become available (i.e., C and N compounds in the form of organic material) are rapidly mineralized by the remaining soil microbes, which causes a pulse in mineralization of N and C [18]. Wetting and drying cycles have also been shown to cause soil aggregates to break apart, exposing physically protected organic matter to further degradation [19,20].

Keeping in view the importance of this studies, and limited work done on this aspect in the study area, the current experiment was hypothesis to find out that both urea and FYM amended soil whether used alone or in combination had no effects on the soil organic matter mineralization and other physico-chemical properties over a period of five weeks.

Material and Methods

The study was conducted to asses the release of mineral N and other soil physico-chemical properties from farmyard manure (FYM) and urea alone and in combinations amended soil during laboratory incubation experiment. The details of laboratory incubation experimental treatments are;

Treatments	Urea (%N)	FYM (%N)	Pool N (kg ha ⁻¹)
T ₁ Control	0	0	0
T ₂ FYM	0	100	120
T ₃ Urea	100	0	120
T ₄ FYM +UREA	50	50	120

Table 1

These four treatments were based on the chemical composition of the FYM. All these four treatments were incubated for 42 days, at two different temperature i.e. 15°C and 25°C, and the experiment was arranged in a CR design with three repeats.

Sample preparation

Soil simple at 0-30 cm depth was collected from NDF The University of Agriculture, Peshawar. Known amount of soil sample (70g) was taken in an incubation petri dish and amended with required amount of fertilizers as per above treatments and 60%

moisture content was added according to weight of samples and maintained moisture content by adding after two days again weighting the samples.

Laboratory analysis

Samples were taken out at 0, 7, 14, 21, 28,35,and 42 days of incubation and analyzed for soil physico-chemical properties including soil mineral N (NH₄-N and NO₃-N), total N, soil organic C, soil EC, pH and salinity.

Soil mineral N

Mineral N was determined by the methods as described by Mulvaney [21]. In this methods, 20 g of soil samples was shaken with 100 ml of 1M KC1 for 1hr and filtered. An aliquot of 20 ml was distilled with 0.2gMgO to recover NH₄-N and NO₃-N. The distillate was collected in 5ml boric acid mixed indicator solution and titrated against 0.005 M HCL. Nitrogen was calculated as 1 ml of 0.005 HCL equals 70µg N. NO3_N was determined by difference.

Soil total nitrogen

Total nitrogen in soil sample was determined by Kjeldahl procedure [22]. In this method, 0.2g of soil sample was digested with 3ml of concentrated H₂SO₄ in the presence of digestion mixture containing K₂SO₄, CuSO₄ and Se on block digested for about 4-5hours. The digestion mixture started at 50°C and then gradually raised to 100, 150, 200, 250, 300, and finally to 350°C which was maintained at least for 1hour. After cooling the digest was transfer to 100ml flask, out of which 20ml was distilled in the presence of 5ml of 40% NaOH solution into a 5ml boric acid mixed indicator solution. The distillate was titrated against standard 0.005N HCL. Nitrogen was calculated as 1ml of 0.005M HCL Equal 70µg N.

Soil organic matter

Organic matter in sample was determined by the walkaly-Black procedure as described in Nelson and Sommers (1996). In this method 1.0g of soil sample was treated with 10ml of 1 N K₂Cr₂O₇ and 20ml of concentrated H₂SO₄. After cooling, 200ml of distilled water were added filtered and titrated against 0.5N FeSO₄ solution using ortrophenolphthalein as indicator with the appearance of maroon colour as end point. A blank titration was also run. Organic matter was calculated from the amount of ferrous sulphat used in titration which determined the amount of unused dichromate and in turn calculated the amount of chromate used in the oxidation of organic C in soil. Organic C was converted to soil organic matter.

Electrical conductivity

Ten gram of soil was taken from soil sample from Petri dish of each treatment in a 100mL bottle and shaken for 60 minutes with 50ml of water (1:5). After shaking the suspension was left over night to allow the soil to settle down. A pipette type conductivity

cell was filled with the supernatant and electrical conductivity was measured in dSm^{-1} (Richard, 1954) using the EC meter.

Soil PH.

Soil pH was determined in 1:5 soil water suspensions [23] using the soil pH meter.

Statistical analysis

The data was analyzed by procedure described in Steel and Torrie [24] using computer software Statistic 8.1 relevant to repeated measurement analysis, basically trialed in CR design. The treatments mean were separated by on the basis of standard errors of mean.

Results and Discussion

Soil mineral Nitrogen ($mg\ kg^{-1}$)

Soil mineral nitrogen is an imporatnt parameter contributing to soil fertility (available nutrient) and is directly or indirectly affected by addition of organic manurs to soil. Soil nitrogen as affected by FYM and urea alone and in combination at various temprature (15°C and 25°C) are presented in Figure 1. It revealed that at 15°C soil mineral nitrogen increased as incubation intervals increased from 0 to 42 days for all all treatments. Similar trend was found at 25°C. Lower soil mineral nitrogen at 15°C and 25°C was found in soil samples collected from experimental units treated with FYM at first interval while higher soil mineral nitrogen was recorded in soil treated with FYM and combination with urea after 5th in both 15°C and 25°C. Fertilizer application in combination with FYM resulted in a greater increase in mineral nitrogen pool than with FYM application alone. Crop availability of nitrogen from FYM is generally lower than the availability of nitrogen from inorganic fertilizer, possibly due to slow release of organically bonded nitrogen immobilization of nitrogen and volatilization of NH_3 from surface applied manure. These results are supported by the finding of Jokela [25], who reported that fertilizer application with FYM greater increase in mineral nitrogen.

Soil total nitrogen (%)

Soil total nitrogen is an imporatnt parameter contributing to soil fertility, and shows the capability of soil to affect the mineralization. The total N content is directly or indirectly affected by addition of organic manurs and fertilizer to soil. Soil nitrogen as affected by FYM and urea alone and in combination at various temprature (15°C and 25°C) are presented in Figure 2. It revealed that at 150C soil nitrogen increased as incubation intervals increased from 0 to 42 days for the all treatments. Similar trend was found at 25°C. Low soil nitrogen at 15°C and 25°C was found in soil samples collected from experimental units at zero interval while higher soil N was recorded in soil treated with FYM alone, compare to control. Our hypothesis that FYM decomposed was rapidly due to low C:N ratio resulted lower aggregate stability and soil nitrogen accumulation. In support of our observations, high N and low C/N ratio OR inputs have been shown to decompose rapidly [26,27].




Figure 1: Graphical representation of soil mineral N amended with manure at different temperature over a period of time.




Figure 2: Graphical representation of soil total N amended with manure at different temperature over a period of time.

Soil organic carbon (%)

Soil organic carbon is an imporatnt parameter contributing to soil fertility which are directly or indirectly affected by addition of organic manurs and fertilizer to soil. Soil organic carbon as affected by FYM and urea alone and in combination at various temprature (15°C and 25°C) are presented in Figure 3. It revealed that at 15°C soil organic C increased as incubation intervals increased from 0 to last sampling for all treatments. Similar trend was found at 25°C. Low soil C at 15°C and 25°C was found in soil samples collected from experimental units treated at first interval while higher soil C was recorded in soil treated with FYM alone compared to low content in control for the same interval. The hypothesis that FYM decompose rapidly due to low C:N ratio resulted lower aggregate stability and Soil organic carbon accumulation. In support of our observations, high N and low C/N ratio OR inputs have been shown to decompose rapidly [26,27].

Soil pH

Soil pH as an important parameter it ranges from 1 to 14. At the range of pH 6.6 To 7.5 most of the nutrient are available and it affect the availability of nutrients at high and low pH. PH of the amended soil as affected by FYM and urea alone and in combination are presented in Figure 4. It revealed that in first intervals the pH is same in all treatment but the pH of manure amended (FYM) soil

decreased and in urea alone increased at 15°C the similar trend was also found in 25°C.

The reduction in soil pH with FYM could be associated with release of H⁺ as result of various organic acids production during decomposition process or ionization of various functional groups associated with organic sources. Our result of reduction in soil pH with application of FYM was in line with the results reported by Sharif, *et al.* [28] who found that addition of organic manure or compost caused reduction in soil pH. Addition of urea to the soil decline CO₂ evolution and increase the soil pH our results was supported by Sarah J., *et al.* [29] Addition of urea showed a significant decline in CO₂ evolution with increasing soil acidity.




Figure 3: Graphical representation of soil Organic carbon (%) of soil amended with manure at different temperature over a period of time.




Figure 4: Graphical representation of pH in soil amended with manure at different temperature over a period of time.

SOIL EC (μscm^{-1})

Figure 5 showed the data of soil EC. It revealed that initially the soil EC was decreased in period of of time at 4th intervals and in last intervals the EC increased and high increased ocured in a soil amended with FYM in both temperature (15°Cand 25°C) which is essential for plant growth. The electrical conductivity (EC) of soil-water mixtures indicates the amount of salts present in the soil. All soils contain some salts, which are essential for plant growth. However, excess salts hinder plant growth by affecting the soil-water balance. The high EC values determined in this experiment

may be explained by the decomposition of the larger organic fragments and their intermediates or end product of FYM during incubation periods. EC values of the soils treated with the FYM were lower than 4dS m⁻¹, the traditional value above which soils present a salinization hazard [23]. The result is in agreement with those of Erdem Yilmaz. And Zeki Alagöz [30] who reported that Soil EC affected with the incorporation of FYM [31-39].




Figure 5: Graphical representation of EC in soil amended with manure at different temperature over a period of time.

Conclusion and Recommendation

- Soil mineral, total N was higher in FYM + urea amended soil, and increased linearly with the incubation period,
- Organic C was higher in FYM either used sole or combined with N and was increased with the incubation time.
- Higher increase rate for Total N was observed in FYM + urea treatment.
- There was a slight change in pH for different treatments; however soil pH was higher for urea treatment than others.

Based on the results, it is recommended that mixed application of FYM and urea release higher mineral N and hence would be better if applied to soil for improved productivity.

Bibliography

1. Kaboneka S, *et al.* "Carbon decomposition of nitrogen form from control released fertilizer". *Communications in Soil Science and Plant Analysis* 28 (1997).
2. Chang C., *et al.* "Barely performance under heavy application of cattle feedlot manure". *Agronomy Journal* 85 (1993): 1013-1018.
3. Ismaily AL., *et al.* "Effect of osmotic and matric potentials on nitrogen mineralization in un amended and manure amended soils". *Soil Science* 173.3 (2008): 203-213.
4. Katterer T, *et al.* "Temperature dependence of organic matter decomposition: A critical review using literature data analyzed with different models". *Biology and Fertility of Soils* 27 (1998): 258-262.

5. Kruse J., *et al.* "Effects of drying and rewetting on carbon and nitrogen mineralization in soils and incorporated residues". *Nutrient Cycling in Agro-ecosystems* 69 (2004): 247-256.
6. Torbert HA and C Wood. "Effects of soil compaction and water-filled pore space on soil microbial activity and N losses". *Communications in Soil Science and Plant Analysis* 23 (1992): 1321-1331.
7. Schjonning P., *et al.* "Turnover of organic matter in differently textured soils, I: Physical characteristics of structurally disturbed and intact soils". *Geoderma* 80 (1999): 177-198.
8. Gordillo R.M., *et al.* "Mineralizable nitrogen in broiler litter, II: Effect of selected soil characteristics". *Journal of Environmental Quality* 26 (1997): 679-1689.
9. Cassman KG and DN Munns. "Nitrogen mineralization as affected by soil moisture, temperature, and depth". *Soil Science Society of America Journal* 44 (1980): 1233-1237.
10. Eghball B. "Nitrogen mineralization from field-applied beef cattle feedlot manure and compost". *Soil Science Society of America Journal* 64 (2000): 2024-2030.
11. Vigil MF and DE Kissel. "Rate of nitrogen mineralized from incorporated crop residues as influenced by temperature". *Soil Science Society of America Journal* 59 (1995): 1636-1644.
12. Campbell CA., *et al.* "Mineralization rate constants and their use for estimating mineralization in some Canadian prairie soils". *Canadian Journal of Soil Science* 64 (1984): 333-343.
13. Linn DM and JW Doran. "Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and no tilled soils". *Soil Science Society of America Journal* 48 (1984): 1267-1272.
14. Knoepp JD and WT Swank. "Using soil temperature and moisture to predict forest soil nitrogen mineralization". *Biology and Fertility of Soils* 36 (2002): 177-182.
15. Soulides DA and FE Allison. "Effect of drying and freezing soils on carbon dioxide production, available mineral nutrients, aggregation, and bacterial population". *Soil Science* 91 (1961): 291-298.
16. Bloem J., *et al.* "Microbial numbers and activity in dried and rewetted arable soil under integrated conventional management". *Soil Biology and Biochemistry* 24 (1992): 655-665
17. Magid J., *et al.* "Drying and rewetting of a loamy sand soil did not increase the turnover of native organic matter but retarded the decomposition of added 14-C labeled plant material". *Soil Biology and Biochemistry* 31 (1999): 595-602.
18. Keift TL., *et al.* "Microbial biomass response to a rapid increase in water potential when dry soil is wetted". *Soil Biology and Biochemistry* 19 (1987): 119-126. 262-286.
19. Adu JK and JM Oades. "Physical factors influencing decomposition of organic materials in soil aggregates". *Soil Biology and Biochemistry* 10 (1978): 109-115.
20. Lundquist EJ., *et al.* "Wet dry cycles affect DOC in two California agricultural soils". *Soil Biology and Biochemistry* 31 (1999): 1031-1038.
21. Mulvaney RL. Nitrogen inorganic form. In: D.L. Sparks (Ed). Method of soil analysis part.3-chemical method, SSSA book series No.5, Inc. ASA, inc. Madison, Wisconsin, USA (1996): 1123-11184.
22. Bremner JM. "Nitrogen Total". In: D.L. Sparks(Ed). Method of soil analysis part 3. *American Society of Agronomy* 37 (1996): 1085-1122.
23. Richards LA. "Diagnosis and Improvement of Saline and Alkali Soils". USA Department of Agriculture Handbook 60. USDA Government Printing Office, Washington, DC (1954).
24. Steel G and JH Torrie. Principles and procedures of statistics: A Biometrical Approach, 2nd edition New York: Mc-Graw-Hill (1980): 633.
25. Jokela WE. "Nitrogen fertilizer and dairy manure effects on corn yield and soil nitrate". *Soil Science Society of America Journal* 56 (1992): 148-154.
26. Melillo J., *et al.* "Carbon and nitrogen dynamics along the decay continuum: plant litter to soil organic matter". *Plant and Soil* 115 (1989): 189-198.
27. Hoorens B., *et al.* "Does initial litter chemistry explain litter mixture effects on decomposition?" *Oecologia* 137 (2003): 578-586.
28. Sharif M., *et al.* "Effect of organic and inorganic fertilizers on the yield and yield components of maize". *Pakistan Journal of Agriculture, Agricultural Engineering and Veterinary Sciences* 20.1 (2004): 11-15.
29. Sarah JK., *et al.* "PH regulation of carbon and nitrogen dynamics in two agricultural soils". *Soil Bio and Biochemistry* 38 (2006): 898-911.
30. Yilmaz E and Z Alagöz. "Effects of short-term amendments of farmyard manure on some soil properties in the Mediterranean region – Turkey". *Journal of Food, Agriculture and Environment* 8.2 (2010): 859-862.

31. Eneji AE., *et al.* "Nitrogen transformation in fores Japanies soil following manures+urea amended". *Communications in Soil Science and Plant Analysis* 33 (2002): 55-66.
32. Gao JQ., *et al.* "Effects of temperature and water saturation on CO2 production and nitrogen mineralization in alpine wetland Soils". *Pedosphere* 19 (2009): 71-77.
33. Guntiñas ME., *et al.* "Effects of moisture and temperature on net soil nitrogen mineralization". *European Journal of Soil Biology* 48 (2012): 73-80.
34. Jha PB., *et al.* "Effect of fertilizer and organic matter inputs on nitrifier populations and N mineralization rates in a dry tropical region". *India Journal of Applied Soil Ecology* 4 (1996): 231-241.
35. Javier EF and RE Tabien. "Nitrogen dynamivs in soil amended with different organic fertilizer". *Journal of Crop Science* 28 (2003): 49-60.
36. Sierra J. "Temperature and soil moisture dependence of N mineralization in intact soil cores". *Soil Bio and Biochemistry Research* 29 (1997): 1557-1563.
37. Statistix. Statistix package, version 8.1. Analytical Software, United state of America (2010).
38. Tian Y., *et al.* "Responses of soil nitrogen mineralization to temperature and moisture in alpine ecosystems on the Tibetan Plateau". *Environmental Sciences* 2 (2010): 218-224.
39. Watts DB., *et al.* "Mineralization of nitrogen in soils amended with dairy manure as affected by wetting/drying cycles". *Communications in soil science and plant analysis* 38 (2007): 2103-2116.

Volume 3 Issue 11 November 2019

© All rights are reserved by Muhammad Bakhtiar., *et al.*