

Chemical, Microbiological and Sensory Profiles of Mixed Fruit Wine from Banana (*Musa acuminata*), Watermelon (*Citrullus vulgaris* L.), Pineapple (*Ananas comosus* L.) and Cucumber (*Cucumis sativus*)

Adeyemi Philips Adebawale*, Gabriel Olaniran Adegoke and Folasade Olabimpe Adeboyejo

Department of Food Technology, University of Ibadan, Ibadan, Nigeria

*Corresponding Author: Adeyemi Philips Adebawale, Department of Food Technology, University of Ibadan, Ibadan, Oyo State, Nigeria.

DOI: 10.31080/ASNH.2020.04.0637

Received: January 31, 2020

Published: February 07, 2020

© All rights are reserved by Adeyemi Philips Adebawale, et al.

Abstract

Blend of several fruits (banana, pineapple, watermelon and cucumber) were used for producing wine. When it is realised that banana on its own may not really be suitable for wine production, however, addition of some fruits and *Saccharomyces cerevisiae* can be a source of obtaining acceptable wine. The aim of this work was to produce and improve the quality of banana wine by using blend that will be acceptable. Watermelon (*Citrullus vulgaris* L.), pineapple (*Ananas comosus* L.) and cucumber (*Cucumis sativus*) musts (In ratio 60:40 and 100% as the control) were added respectively to banana must. Physico-chemical, antioxidants, colour, microbial, organic compounds and sensorial profiles of wine samples were done using standard methods. The pH, volatile acidity and phenolics contents of the samples of wine ranged from 3.52 to 3.62, 0.30 to 0.60%, 602.76 mg GAE/l to 858.51 mg GAE/l respectively. Hue angles of the samples of wine were within 0-90°; from 57.45° to 69.40°, and were significantly different (p<0.05) from each other. Colour of the wines was generally orange-red and microbiological examination of the samples showed no contamination with spoilage bacteria like acetobacter, pediococcus and gluconobacter species. Using Fourier Transform Infrared Spectroscopy (FTIR), the samples of wine were found to contain probable compounds like water, alcohol and esters. Principal Component Analysis (PCA) was useful in assessing the relationship and correlations of some of the parameters examined. In this study, blends rich in pineapple and watermelon were more acceptable to taste panelists than 100% banana wine. Our findings with this research work has shown that good quality mixed fruit wine can be produced using combination of banana and other fruits in right proportions.

Keywords: Fruits; Banana; Watermelon; Pineapple; Cucumber; Blend

Introduction

Apart from water and milk no other drink has universal acceptance and esteem throughout ages as wine [1]. The natural chemical balance of grape juice aids its fermentation into wine without the addition of sugars, acids, or other nutrients thus making grape to be a preferred fruit for wine production. Grapes are however not readily available and are also costly in some tropical regions [2]. Banana and other fruits can be used in the production of wine but due to the nature of banana fruit juice, it tends to affect its suitability for wine and this has led to use of enzymes (pectinase and α -amylase) and recombinant yeast strains [3,4]. Producing wine from blends of fruits can provide good substrate for wine production [5]. When it is realised that banana juice contains high concentration of sugars and many acids, watermelon has beta carotene and lycopene, pineapple with its flavour and good source of sugar, ascorbic acid, cucumber with naturally-purified moisture content and high levels of flavonoids (apigenin, luteolin, quercetin, and kaempferol) and cucurbitacins — banana wine with all these nutrients from different fruits will improve its quality. Our present

study was therefore carried out to ascertain the quality profiles and acceptability of wines produced from mixtures of banana and other fruits [6-10].

Materials and Methods

The study took place at the Department of Food Technology, University of Ibadan, Ibadan and National Horticultural Research Institute (NIHORT), Jericho, Ibadan, Nigeria between the months of May 2018 and December 2018.

Collection of raw materials

Ripe, mature and unbruised samples of banana (*Musa acuminata*), pineapple (*Ananas comosus* L.), watermelon (*Citrullus vulgaris* L.) and cucumber (*Cucumis sativus*) were purchased (June, 2018) in Ibadan, Nigeria and were identified as authentic samples at the National Horticultural Research Institute (NIHORT), Jericho, Ibadan, Nigeria before processing into wine. Commercial grade sugar (Dangote Sugar, Nigeria) and yeast were obtained locally. All chemicals and reagents used were of analytical grade.

Sample preparation

Preparation of must and formulation

The fruits were washed with potable water and peeled, and juice extraction was carried out by homogenising the pulp in a blender (Scanfrost, SFKAB409, China) with distilled water (pulp; water, 1:1). To prevent browning and to inhibit unwanted microflora in the juice, 100mg/l potassium metabisulphite was added. It was boiled at 80°C for 5mins and held overnight at room temperature for clarification. The musts obtained contained low sugar and it was chaptalised to 20° brix with the aid table sugar and held at pH 4. The formulations used were modified from those used described elsewhere [11] and samples were coded as follows: sample 945 (banana 60% + pineapple 30% + watermelon 5% + cucumber 5%), sample 238 (banana 60% + pineapple 5% + watermelon 5% + cucumber 30%), sample 172 (banana 60% + pineapple 5% + watermelon 30% + cucumber 5%), sample 561 (banana 60% + pineapple 13.3% + watermelon 13.3% + cucumber 13.3%) and sample 476 (banana 100%).

Starter culture, fermentation and clarification

Saccharomyces cerevisiae; active dry yeast, was activated in distilled water at 30°C and added to the must at a concentration of 3%. After which fermentation was carried out at ambient temperature (27°C) for 6days with monitoring using the hydrometer until the reading reached 1.000-1.020. Secondary fermentation proceeded until all the sugars in the must were utilised completely. Thereafter, the 'young wines' were allowed to age for 4 months. Yeast and other materials that settled at the bottom of the container used for fermentation, were siphoned off or/racked, then 150 ml of prepared gelatin (10%) was added into each of the wine container followed by stirring to dissolve properly. 100 ml of the mixture was collected into a sterile bottle which was covered tightly and used to monitor the process of clarification. Filtration followed after the wines clarification: using muslin cloth, sieve and syphon tubes sterilised by 70% alcohol. The wines was syphoned into each sieve containing four layers of muslin cloth followed by pasteurisation and addition of potassium sorbate [1].

Determination of pH, Total Titrable Acidity and Volatile Acidity pH was determined using a pH meter (HANNA Instruments, H12210-01, Benchtop, Rhode Island, USA).

Total Titrable Acidity (TTA) was determined by titrating 9 ml of wine sample, against 0.1N sodium hydroxide using phenolphthalein as an indicator and pink colour obtained marked the end point of titration with the percentage titratable acidity expressed as tartaric acid (milliequivalent, 0.075).

Volatile acidity was expressed as percentage acetic acid and was measured by steam distillation; 100 ml of wine sample were boiled for 15 mins under reflux to expel carbon dioxide. The flask was cooled and the liquid steam-distilled to reduce the volume to 50 ml. This was titrated with 0.1N NaOH using 2 drops of phenolphthalein, 1 ml of NaOH with the milliequivalent 0.006 of acetic acid [12].

Determination of total solids, soluble solids, alcohol content and temperature

Total solids and, specific gravity of each wine sample were carried out. Hand Held Digital Abbe refractometer (Wincom Company Limited, Pocket Type, Changsha, China) was used to determine the soluble solids of the wine blends samples according to Daramola and Asunni. The specific gravities of the wines were determined using the hydrometer method and the results were the reading stem which was used for alcohol content [5,13].

Temperature of each wine sample was measured using mercury-in-glass thermometer [13].

$$\%ABV = \frac{\text{Initial Sg} - \text{Final Sg} \times 100}{7.37} \quad (1)$$

Antioxidative potentials analyses

Determination of free radical scavenging

Free radical scavenging ability of the wine extract against DPPH (1, 1- diphenyl-2-picrylhydrazyl) was determined. 1 ml of the extract to was mixed with 1 ml of 0.4 mM methanolic solution of DPPH (SigmaAldrich) [14]. The mixture was left in the dark for 30 min and thereafter absorbance of the mixture measured at 516 nm. The percentage inhibition was calculated as a percentage of DPPH discolouration using the equation:

$$\% \text{ Inhibition} = \frac{[(ADPPH - AS)]}{ADPPH} \times 100 \quad (2)$$

Where, AS is the absorbance of the solution when the sample extract has been added at a particular level and ADPPH is the absorbance of the DPPH solution.

Determination of reducing antioxidant property

Reducing property of wine extract was determined using method described elsewhere [15]. 0.25 ml of the wine extract was mixed with 0.25 ml of 200 mM of sodium phosphate buffer pH 6.6 and 0.25 ml of 1% potassium ferric cyanide. The mixture was incubated at 50°C for 20 min, after which 0.25 ml of 10% trichloroacetic acid (TCA) was also added and centrifuged at 2000 rpm for 10 min, 1 ml of the supernatant was mixed with 1 ml of distilled water and 0.1% of FeCl₃ and the absorbance was measured at 700 nm in a spectrophotometer (GENESYS 10SVis, Thermo Scientific, city, USA). Ferric reducing antioxidant power (FRAP) values were obtained by comparing the absorption change in the test mixture with those obtained from increasing concentrations of Fe²⁺, and expressed as mmol of Fe²⁺ equivalents per litre of sample.

Determination of scavenging ability

2, 2'-azino-bis (3-ethylbenthiazoline-6-sulphonic acid) (ABTS) scavenging ability of what the wine extract was determined according to the method described by Re., *et al.* [16]. ABTS in full here was generated by reacting ABTS stock solution (7mM) with 2.45 mM K₂S₂O₈ (1/1, v/v) in the dark for 16 hours until the reaction was completed and the absorbance was stable. The (ABTS) solution was diluted with ethanol to an absorbance of 0.700 ± 0.05 at 734 nm for measurements. Thereafter, 0.2 ml of the appropriate dilution of the

wine extract was then added to 2.0 ml of ABTS solution and mixed for 45 sec; and the absorbance was read at 732nm after 15 mins. The trolox equivalent antioxidant capacity was subsequently calculated (264.32g). The results were expressed as mmol of Trolox per litre of the sample.

Determination of total phenolic content

The Total Phenolic Content of the extract was determined using the method described by Singleton [17]. 0.2 ml of the extract was mixed with 2.5 ml of 10% Folin ciocalteau's reagent (Merck, Darnstadt, Germany) and 2 ml of 7.5% sodium carbonate. The reaction mixture was subsequently incubated at 45°C for 40 mins, and the absorbance was measured at 700nm using a spectrophotometer—Gallic acid was used as the standard and data obtained were expressed as milligram of Gallic Acid Equivalents (GAE) per litre of extract.

Determination of Total Flavonoid Content

Total flavonoid content of the extract was determined using a colourimeter assay developed by Bao [18]. 0.2 ml of the extract was added to 0.3 ml of 5% NaNO₃ at zero time. After 5 mins, 0.6 ml of 10% AlCl₃ was added and after 6min, 2 ml of 1M NaOH was added to the mixture followed by the addition of 2.1 ml of distilled water. Absorbance was read at 510nm against the reagent blank and flavonoid content was expressed as mg catechin equivalent.

Determination of tannin

Tannin determination was done according to the method described by AOAC with some modifications: 1 ml of the wine extract was filtered into 100 ml volumetric flask, followed by the addition of 20 ml of distilled water, 2.5 ml of Folin-Denis reagent and 10 ml of 17% aqueous. Na₂CO₃ was also added and thoroughly mixed together. The mixture was then made up to 100ml with distilled water, mixed and allowed to stand for 20 mins. The bluish-green colour developed at the end of the reaction mixture of different concentrations ranging from 10-50ppm. The absorbance of the tannic acid standard solutions as well as sample was measured after colour development at 760nm using spectrophotometer AJI-CO₃ UV-VIS [12].

Colour determination

Colour (L*a*b) parameters were determined using the colourimeter (Colour Tec PCMTM Colour Tec Associates, Inc., Clinton, NJ, USA). L, a, and b parameters were determined by placing the sensor of the colourimeter on the sample (19). From the data obtained, the following parameters were also derived:

$$\text{Hue angle} = \tan^{-1} (b/a) \quad (3)$$

$$\text{Whiteness Index (WI)} = 100 - [(100 - L^*)^2 + (a^*)^2 + (b^*)^2]^{0.5} \quad (4)$$

$$\text{Yellowness Index (YI)} = 142.86 b^* / L^* \quad (5)$$

$$\text{Chroma} = [(a^*)^2 + (b^*)^2]^{0.5} \quad (6)$$

$$\text{Colour difference } \Delta E^* = [\Delta a^* \Delta b^* \Delta c^*]^{0.5} \quad (7)$$

Sensory evaluation

Sensory evaluation was carried out by a panel of 20 semi-trained panelists from Food Technology Department, University of Ibadan and National Horticulture Research Institute, Ibadan, Nigeria. Each member compared wines for colour, flavour, taste, clarity, and overall acceptability on 9-point hedonic scale where 9 denotes excellent and 1 very poor.

The sensory evaluation procedure was modified as described by Ogodo, Ugboogus, and Ezeonu and the exercise was carried out on the second day of wine production [11].

Microbiological analysis

Microflora of different blends was determined using Potato Dextrose Agar, Nutrient Agar, MacConkey Agar, de Man, Rogosa Sharpe and Carr Medium as described elsewhere [20,21].

Fourier transform infrared spectroscopy (FTIR) analysis

The wines were subjected to IR spectroscopy (Shimadzu, Japan). The infrared analysis of the wine was carried using Nujol as the sample holder; operated in a wavelength range of 4000-500 cm⁻¹ using FT-IR 8300 series [12].

Statistical analyses

Data obtained from samples analysed in triplicates were subjected to analysis of variance (ANOVA) and Duncan's multiple range test using the Statistical Package for the Social Sciences (SPSS) version 20.0 software (SPSS Inc., Chicago, IL, USA); **P<0.05 was considered statistically significant; reported results are expressed as the Mean ± SD. Principal Component Analysis by XLSTAT version 2018.7 (Addinsoft, U.S.A).

Results

The physico-chemical properties of wines examined in this study are shown in Table 1. pH values in the mixed wines which were in the range of 3.10~3.60 were acidic throughout the period of fermentation. Values for total titrable acidity (TTA) were between 0.60~0.82%, with samples 945 and 172 being more acidic with no significant difference (p>0.05). However, sample 945 had the highest volatile acidity of 0.66%.

The percentage alcohol in the wines at the end of fermentation were 11% (v/v) for samples 476 and 561 while samples 172, 238 and 945 had 12% (v/v) respectively. Total soluble solids (TSS) values ranged from 3.2~4.9 after the fermentation stopped. Temperature of the final mixed wines ranged from 20.7°C to 23.1°C. The antioxidative assays (Tables 2,3) were as follows (Table 2 and Table 3). DPPH assay in the samples was in the decreasing order sample 945 (80.70% Ihb) > sample 172 (78.26% Ihb) > sample 238 (64.99% Ihb) > sample 561 (62.05% Ihb) > sample 476 (55.67% Ihb) correlating with other methods used to determine antioxidative potential; Ferric Reducing Property (FRAP) and Trolox Equivalent Antioxidant Capacity (TEAC). Total Phenols in the wine samples increased in order of 602.76 mg GAE/l~ 858.51 mg GAE/l)

Wine	pH	TTA%	Volatile Acidity %	ABV (%)	Total Soluble Solids (°B)	Total Solids (%)	Fermentation Temperature (°C)
238	3.57 ^b ± 0.01	0.64 ^b ± 0.02	0.46 ^b ± 0.01	12	3.2	2.40 ^c ± 0.02	21.3 ± 1.2
561	3.60 ^a ± 0.01	0.60 ^b ± 0.02	0.35 ^d ± 0.02	11	4.7	2.95 ^b ± 0.01	20.4 ± 1.3
172	3.53 ^c ± 0.02	0.78 ^a ± 0.03	0.64 ^a ± 0.01	12	3.9	2.42 ^c ± 0.01	23.1 ± 1.5
945	3.52 ^c ± 0.01	0.82 ^a ± 0.03	0.66 ^a ± 0.02	12	3.7	2.35 ^d ± 0.01	22.3 ± 1.7
476	3.62 ^a ± 0.02	0.62 ^b ± 0.01	0.42 ^c ± 0.01	11	4.9	3.09 ^a ± 0.01	20.7 ± 0.8

Table 1: Physico-Chemical properties of the wines.

Values are mean ± SD of 3 replicates. Different letters (a-e) within same row are significantly different (P<0.05) according to Duncan's multiple range test. 238: (60+5+5+30)%, 561: (60+13.3+13.3+13.3)%, 172: (60+5+30+5)%, 945: (60+30+5+5)%, 476: (100%).

which follows the same pattern with tannins. Sample 172 had the highest amount of tannin (128 mg/l) while sample 561 (90.42 mg/l) had the least amount of tannins. It was also evident from the colour analysis, a* redness to greenness level of the sample, highest a* value was observed in sample 172. The results of hue angles (E.q 2.3) for the blends fell within 0-90°; from 57.45° to 69.40°.

Acceptability of the wines as were as follows: sample 945 > sample 172 > sample 238 > sample 561 > sample 476 (Table 4). There were no significant differences in the microbial loads in the of samples 172, 238 and 945 but these samples were significantly different to sample 561 and no growth was observed in sample 476 (Table 5).

Wine	DPPH (%Ihb)	TEAC ¹ (mMolTrolox/l)	FRAP ² (mMolFe ²⁺ /l)	Phenols ³ (mgGAE/l)	Flavonoids ⁴ (mg/l)	Tannin (mg/l)
238	64.99 ^b ± 1.91	49.62 ^c ± 1.39	44.80 ^c ± 1.00	730.35 ^c ± 18.9	309.44 ^c ± 25.0	109.55 ^c ± 2.76
561	62.05 ^c ± 0.93	42.84 ^d ± 1.72	20.70 ^d ± 2.10	602.76 ^e ± 2.3	476.11 ^b ± 13.89	90.42 ^b ± 0.34
172	78.26 ^a ± 0.91	62.50 ^b ± 1.17	50.61 ^b ± 1.10	858.51 ^a ± 29.31	509.45 ^a ± 8.33	128.78 ^a ± 4.39
945	80.70 ^a ± 1.36	86.39 ^a ± 1.27	69.11 ^a ± 1.11	786.10 ^b ± 2.87	312.23 ^c ± 5.56	117.91 ^c ± 0.37
476	55.67 ^d ± 1.45	38.62 ^e ± 1.94	19.40 ^e ± 1.00	634.37 ^d ± 0.58	245.56 ^d ± 5.55	95.16 ^d ± 0.09

Table 2: Antioxidants properties of the wines.

Values are mean ± SD of 3 replicates. Different letters (a-e) within same row are significantly different (P<0.05) according to Duncan's multiple range test. ¹FRAP: Ferric reducing antioxidant power. ²TEAC: Trolox equivalent antioxidant activity ³GAE: gallic acid equivalent. ⁴Cat: Catechin equivalent. 238: (60+5+5+30)%, 561: (60+13.3+13.3+13.3)%, 172: (60+5+30+5)%, 945: (60+30+5+5)%, 476: (100+0+0+0).

Wine	L*	a*	b*	h*	C*	WI	YI	ΔE*
238	50.57 ^c ± 0.7	-3.50 ^e ± 0.02	9.58 ^c ± 0.06	69.93 ^a ± 0.05	10.19 ^d ± 0.01	49.52 ^c ± 0.03	27.06 ^c ± 0.10	23.13 ^c ± 0.08
561	48.63 ^d ± 0.4	-5.19 ^d ± 0.06	8.13 ^e ± 0.26	57.45 ^e ± 0.27	9.65 ^e ± 0.02	47.73 ^d ± 0.08	23.88 ^d ± 0.30	21.96 ^d ± 0.19
172	59.56 ^a ± 0.0	5.44 ^c ± 0.01	9.20 ^d ± 0.05	59.40 ^d ± 0.08	10.69 ^c ± 0.01	58.17 ^a ± 0.01	22.07 ^e ± 0.02	31.97 ^a ± 0.02
945	56.01 ^b ± 0.1	-6.27 ^a ± 0.11	11.15 ^b ± 0.18	60.64 ^c ± 0.09	12.79 ^b ± 0.02	54.19 ^b ± 0.06	28.44 ^b ± 0.13	28.74 ^b ± 0.14
476	47.12 ^e ± 0.1	-5.71 ^b ± 0.01	13.23 ^a ± 0.01	66.65 ^b ± 0.01	14.4 ^a ± 0.02	45.19 ^e ± 0.07	40.11 ^a ± 0.01	20.45 ^e ± 0.06

Table 3: Colour determination of the wines.

Values are mean ± SD of 3 replicates. Different letters (a-e) within same row are significantly different (P<0.05) according to Duncan's multiple range test. L*: Lightness; a*: red (+)/green (-); b*: yellow (+)/blue (); C*: Chroma; ΔE*: Total colour difference; h*: Hue angle; WI: Whiteness Index; YI: Yellowness Index. 238: (60+5+5+30)%, 561: (60+13.3+13.3+13.3)%, 172: (60+5+30+5)%, 945: (60+30+5+5)%, 476: (100+0+0+0)%.

Wine	Taste	Aroma	Colour	Clarity	Overall A.
238	6.96 ^a ± 1.50	6.00 ^b ± 1.60	7.04 ^{ab} ± 1.68	6.62 ^{ab} ± 1.33	6.8 ^{ab} ± 1.02
561	6.81 ^a ± 0.90	6.38 ^{ab} ± 1.65	6.27 ^{bc} ± 1.66	6.46 ^{ab} ± 1.24	6.46 ^{bc} ± 1.30
172	6.30 ^a ± 1.67	7.00 ^a ± 1.50	7.27 ^a ± 1.43	7.00 ^a ± 1.01	7.03 ^a ± 1.31
945	6.40 ^a ± 1.77	7.19 ^a ± 1.09	7.70 ^a ± 0.96	6.96 ^a ± 1.92	7.46 ^a ± 0.99
476	6.90 ^a ± 1.30	5.5 ^b ± 1.80	5.62 ^c ± 1.50	5.84 ^b ± 1.95	6.00 ^c ± 1.64

Table 4: Sensory evaluation of the wines.

Values are mean ± SD of 3 replicates. Different letters (a-e) within same row are significantly different (P<0.05) according to Duncan's multiple range test. 238: (60+5+5+30)%, 561: (60+13.3+13.3+13.3)%, 172: (60+5+30+5)%, 945: (60+30+5+5)%, 476: (100%).




Wine	NA (cfu/ml)	PDA (cfu/ml)	MCA (cfu/ml)	CAR (cfu/ml)	MRS (cfu/ml)
238	NG	NG	NG	2.1×10 ^{1a} ± 10	NG
561	NG	2.4×10 ^{3b} ± 200	NG	1.1×10 ^{1b} ± 10	NG
172	NG	NG	NG	2.1×10 ^{1a} ± 10	2.13×10 ^{1a} ± 30
945	NG	NG	NG	2.0×10 ^{1a} ± 10	1.10×10 ^{1b} ± 20
476	NG	7×10 ^{3a} ± 200	NG	NG	NG

Table 5: Microbial analysis of the wines.

Values are mean ± SD of 3 replicates. Different letters (significantly different (P<0.05) according to Duncan's multiple range test. NA: Nutrient Agar; PDA: Potato Dextrose Agar; MCA: MarConkey Agar; CAR: Car medium; MRS: de Man, Rogosa Sharpe; NG: No Growth. 238: (60+5+5+30)%, 561: (60+13.3+13.3+13.3)%, 172: (60+5+30+5)%, 945: (60+30+5+5)%, 476: (100+0+0+0)%.

Fourier Transform Infrared results of the wines; provided the functional groups of the organic compounds present in the samples. The organic compounds were: hydroxyl and carboxylic acids (O-H) - 3300-3600 cm⁻¹, esters (C-O) - 1000-1300 cm⁻¹ (2 bands) cm⁻¹, ether (C-O) - 1050-1200 cm⁻¹, and 1070-1150 cm⁻¹, amines (NH₂ and N-H, C-N, NH₂, and N-H) - 600-900 cm⁻¹, 1000-1250 cm⁻¹, 1550-1650 cm⁻¹, and 3300-3500 cm⁻¹ (Figures 1(a)-(e)).

Using principal component analysis for the wines and their quality - was employed. Samples 172 and 945 were on one side of the plot while samples 238, 561 and 476 were on the other side (Figures 2 and 3). Principal Component Analysis permits visual interpretation of the data represented by two-dimensional scatter plots [22,23].



Discussion

In this study, changes in pH of wines (Table 1) which were similar to the ones reported for banana wine elsewhere could be due to the production of acids within the period of fermentation probably arising from microbial succession [24,25]. When it is realised that increase in acidity of the samples of wine examined in this study could be due to accumulation of organic acids during fermentation low pH of the wine samples protected them against microbial spoilage and also produced at the same time more rapid natural clarification with greater effectiveness of stabilisation treatments and a longer shelf life [26]. In this work, the concentrations of ethanol contributed to the whole characteristic quality and flavour of the wine produced [27]. Total solids of the wines examined in this study were low; low values of total solids can however be attributed to the efficiency of yeasts used for fermentation.

Temperature is one of the most important parameters in oenology because it influences the fermentation kinetics and the chemical quality of wine [26]. During the period of fermentation there were fluctuations in the temperature, however, wine fermentation should be between 13-35°C [28]. The observed changes in the temperature of the wines could be due to microbial succession arising from microbial metabolic activities [29]. Bacteria have both positive and negative influences on wine production. From the morphological examination and biochemical tests done, the samples are not colonised with spoilage bacteria like pediococcus, acetobacter and gluconobacter: invariably, this shows that the wine has not been spoiled; the differences in microbial loads could be due to differences in the fruit wine blends.

Wines are healthy beverages that have been used as natural remedy for man's illness from early days [29]. In evaluating the antioxidative of any food product; it is essential to use at least three radical, as the solvent and substrate do impact on the values obtained [30]. Sun., *et al.* have also proved the influence of pineapple in scavenging free radical through DPPH assay [31]. Besides, watermelon in the work of Melo., *et al.* showed remarkable free radical scavenging and antioxidant activity owing to potent antioxidant lycopene [32]. According to Gawel, tannins are polymeric flavonoid compounds containing subunits of flavan-3-ol, responsible for colour stability, astringency and active antioxidants—results for flavonoids shows positive correlation while sample 476 (245.56 mg Cat/l) had the least amount of flavonoids [33]. And also the intensity of the sample a* could be indication of concentration level of lycopene, while sample 476 had the highest level of yellowness index (E.q 2.5) due to the level of high concentration of the banana which was 100%, which implies orange-red colouration. The most whitish sample was 945 and the least was sample 476, while samples 172 and 945 had the highest L* value of 59.56 and 56.01 respectively, this could be due to the low pH as reported by Ribereau-Gayon [26].

With respect to the sensory analysis of the samples of wine examined in this present study, the mean scores for overall acceptability showed no significant difference between samples 172, 238

and 945 meaning that the samples were equally accepted while samples 476 and 561 were significantly different ($p > 0.05$).

From the FTIR spectra (Figures 1(a)–(e)), the hydroxyl group in the wine sample was confirmed to be phenolics—Phenolics have one phenol group (at least one hydroxylated benzene ring), and the other more complex phenol group that probable to be present is flavonoids likewise tannins: which are complex esters of glucose and gallic acid.

From the analysis carried out with PCA for the physico-chemical properties, colour determination and sensory characteristics—the closeness of each of the parameters show their relationship; Also, these might be important in defining the acceptability of samples 172 and 945—alcohol, colour, volatile acidity, total titrable acidity clarity, aroma, whiteness index, colour difference and lightness index have appeared to be positively correlated owing from results. Thus, the different blends of the fruit in each fruit wine led to the differences and most importantly the acceptability of samples 172 and 945. Better still, the samples also have shown to be better in terms of antioxidant properties.

Conclusion

Mixed fruit wine was produced from blending of banana, pineapple, watermelon and cucumber musts. From the results of this present study; fruit wine from appropriate levels of banana 60% + pineapple 30% + watermelon 5% + cucumber 5% and banana 60% + pineapple 5% + watermelon 30% + cucumber 5% were found to be acceptable to panelists particularly with respect to aroma, colour and clarity.

Acknowledgements

Authors are grateful to staff of the Department of Food Technology, University of Ibadan, Ibadan and National Horticulture Research Institute (NIHORT), Ibadan, Nigeria for their assistance during this research work.

Bibliography

1. Swami SB., *et al.* "Fruit Wine Production: A Review". *Journal of Food Research and Technology* 2.3 (2014): 93-100.
2. Aloba AP and Offonry SU. "Characteristics of coloured wine produced from Roselle (*Hibiscus sabolaritts*) calyx extract". *Journal of the Institute of Brewing* 115.2 (2009): 91-94.
3. Chilaka CA., *et al.* "Evaluation of the efficiency of yeast isolates from palm wine in diverse fruit wine production". *African Journal of Food Science* 4.12 (2010): 764-774.
4. Jackson T and Badrie N. "Quality Changes on Storage of Caribbean Banana (*Musa acuminata*) Wines: Effects of Pectolase Concentration and Incubation Period". *Journal of Wine Research* 13.1 (2012): 43-56.
5. Awe S. "Production and microbiology of pawpaw (*Carica papaya* L) wine". *Current Research Journal of Biological Sciences* 3.5 (2011): 443-447.

6. Mohapatra D., et al. "Post-harvest Processing of Banana: Opportunities and Challenges Food". *Bioprocess Technology* 4 (2011): 327-339.
7. Collins JK., et al. Sensory evaluation of low sugar watermelon by consumers". *Horticulture Science* 40 (2015): 883.
8. Baruwa OI. "Profitability and constraints of pineapple production in Osun State, Nigeria". *Journal of Horticultural Research* 21.2 (2013): 59-64.
9. Mukherjee PK. "Phytochemical and therapeutic potential of cucumber". *Fitoterapia* 84 (2013): 227-236.
10. Idise OE and Odum EI. "Studies of wine produced from banana". *International Journal of Biotechnology and Molecular Biology Research* 2 (2013): 209-214.
11. Ogodo AC., et al. "Production of mixed fruit (pawpaw, banana and watermelon) wine using *Saccharomyces cerevisiae* isolated from palm wine". *Springer Plus* 4 (2015): 683.
12. AOAC. Official Method of Analysis. 18th ed. Association of Official Analytical Chemists, Washington, DC, USA (2005).
13. Daramola B and Asunni OA. "Preparation, physicochemical and sensory assessment of pawpaw-red ginger food drink". *American-European Journal of Scientific Research* 22 (2007): 101-105.
14. Gyamfi MA., et al. "Free radical scavenging action of medicinal herbs from Ghana: *thonningia sanguine* on experimentally induced liver injuries". *General Pharmacology* 32 (1999): 661-667.
15. Publiido R., et al. "Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay". *Journal of Agricultural and Food Chemistry* 48 (2000): 3396-3402.
16. Re R., et al. "Antioxidant cavity applying an improve ABTS Radication decolourisation assay". *Free Radical Biology and Medicine* 26 (1999): 1231-1237.
17. Singleton VL., et al. "Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Cioalteau Reagents". *Methods in Enzymology* 299 (1999): 152-178.
18. Meda A., et al. "Determination of the total phenolic, flavonoid and proline contents in Burkina Fasan honey, as well as their radical scavenging activity". *Food Chemistry* 91 (2005): 571-577.
19. Hunt RWG and Pointer MR. "Measuring colour". John Wiley and Sons Ltd., Publication (2011).
20. Cowan I and Steel ST. "Cowan and Steel's Manual for the identification of medical bacteria". 3rd Edn. Cambridge University Press, London (2004).
21. Yamada Y., et al. "Identification of acetic acid bacteria isolated from Indonesian sources, especially of isolates classified in the genus *Gluconobacter*". *The Journal of General and Applied Microbiology* 45 (1999): 23-28.
22. Destefanis G. "The use of principal component analysis (PCA) to characterise beef". *Meat science* 56.3 (2000): 255-259.
23. Šnirc M., et al. "Application of principal component analysis method for characterization chemical, technological, and textural parameters of farmed and pastured red deer". *International Journal of Food Properties* 20.4 (2017): 754-761.
24. Robinson J. *The Oxford Companion to Wine* (3rd Edn.), Oxford University Press, USA. (2006): 840.
25. Akubor PI., et al. "Production and evaluation of banana wine". *Plant Foods for Human Nutrition* 58 (2013): 1-6.
26. Ribe´reau-Gayon P. "Handbook of enology: The microbiology of wine and vinifications". ed., John Wiley and Sons Ltd., Chichester (2006).
27. Reddy LV and Reddy OVS. "Production, optimization and characterization of wine made from mango (*Mangifera indica*)". *Natural Product Radianc* 8.4 (2009): 426-435.
28. Molina A.M., et al. "Influence of wine fermentation on the synthesis of yeast-derived volatile aroma compounds". *Applied Microbiology and Biotechnology* 77.3 (2007): 675-687.
29. Okafor N. "Modern industrial microbiology and biotechnology". Science publishers, Enfield, (2007): 262-278.
30. Badejo AA., et al. "Changes in Nutrient Composition, Antioxidant Properties, and Enzymes Activities of Snake Tomato (*Trichosanthes cucumerina*) during Ripening". *Preventive Nutrition and Food Science* 21.2 (2016): 9096.
31. Sun J., et al. "Antioxidants and antiproliferative activities of common fruits". *Journal of Agricultural and Food Chemistry* 50 (2002): 7449-7454.
32. Melo EA., et al. "Capacidade antioxidante de frutas". *Journal of Pharmaceutical Sciences* 44 (2008): 194-201.
33. Gawel R. "Red wine astringency: A review". *Australian Journal of Grape and Wine Research* 4 (1998): 74-95.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: <https://www.actascientific.com/>

Submit Article: <https://www.actascientific.com/submission.php>

Email us: editor@actascientific.com

Contact us: +91 9182824667