



## Comparative Characteristics of the Mineral Composition of the Meat of the Black Sea Grass Shrimp *Palaemon Adspersus*, Rathke, 1837 in Different Fishing Periods

Lebskaya Tatyana Konstantinovna<sup>1\*</sup>, Bal-Prilipko Larisa Vatslavovna<sup>2</sup>,  
Lebsky Sergey Olegovich<sup>3</sup> and Slobodyanyuk Natalia Mikhailovna<sup>4</sup>

<sup>1</sup>Doctor of Technical Sciences, Professor of the Department of Technology of Meat, Fish and Seafood, Ukraine

<sup>2</sup>Doctor of Technical Sciences, Professor, Dean of the Faculty of Food Technologies and Quality Management of Agricultural Products, Ukraine

<sup>3</sup>postgraduate student of the Department of Technology of Meat, Fish and Seafood, Ukraine

<sup>4</sup>Candidate of Agricultural Sciences, Head of the Department of Technology of Meat, Fish and Seafood, Ukraine

**\*Corresponding Author:** Lebskaya Tatyana Konstantinovna, Doctor of Technical Sciences, Professor of the Department of Technology of Meat, Fish and Seafood, Ukraine.

**DOI:** 10.31080/ASNH.2022.06.1029

**Keywords:** Nutritional Value; Shrimp Meat; Essential; Toxic; Poorly Studied Mineral Elements; Physiological Needs; Safety

### Annotation

The nutritional value of the mineral composition of the meat of one of the species of crustaceans of the Black Sea-*Palaemon adspersus* Rathke, 1837 in the spring and autumn periods of fishing-has been studied. Essential mineral elements. Calcium (Ca) Potassium (K), Sodium (Na) Magnesium (Mg), Copper (Cu), Zinc (Zn), Iron (Fe), Manganese (Mn), Nickel (Ni), Lithium (Li), Little Studied (Silver (Ag), Aluminum (Al), Barium (Ba), Boron (B), Bismuth (Bi), As Well as Toxic (Arsenic (As), Cadmium (Cd), Swine c (Pb), Mercury (Hg).. The level of most mineral elements in the meat of this crustacean is lower than the physiological needs for an adult, with the exception of Cu, Cr, Ni and Al. Their amount significantly exceeds the recommended values for humans, however substantially below the toxicity threshold. The content of heavy metal in the meat of *Palaemon adspersus* Rathke, 1837 during catch periods does not exceed the permissible norms.

### Introduction

The nutritional value of raw materials is determined by many constituent factors, among which the qualitative and quantitative composition of mineral elements [1-5], as well as their safety [6] play a significant role in ensuring the physiological needs of a person [7,8] Mineral components of nutrition are characterized by

various physiological functions: play an important role in plastic processes, the formation and construction of body tissues, are necessary to maintain acid-base balance in the body, create a certain concentration of hydrogen ions in tissues and cells, interstitial and intercellular fluids, participate in enzymatic processes as activators and cofactors of enzymes [7,8].

Hydrobionts are a source of biologically valuable proteins, lipids, carbohydrates, mineral components, vitamins with health benefits for all age categories of people and their consumption is promoted as a healthy food choice [9-11].

Based on this, the World Health Organization and the Food and Agriculture Organization of the United Nations recommend regular consumption of seafood in amounts of one to two servings per week [12].

### Study of mineral components in the meat of hydrobionts, including

crustaceans, a significant number of works are devoted [13-21]. In these studies, it is shown that their meat contains all the essential, poorly studied and toxic elements. Qualitative and quantitative composition reveals significant fluctuations that are associated

**Received:** February 14, 2022

**Published:** March 22, 2022

© All rights are reserved by **Lebskaya Tatyana Konstantinovna., et al.**

with their type, size, sex, degree of development of the reproductive system, season and catch area, nutritional characteristics. Thus, on the example of the Antarctic krill *Euphausia superba*, it was found that sexually mature females and males differ between themselves and immature individuals in the content of manganese, zinc, copper, nickel, strontium and cadmium [16,20]. *Pandalus borealis*, which lives in the Pacific region [17-19], reveals a number of differences in both the qualitative and quantitative content of the mineral components of meat compared with the meat of related species-*Euphausia superba* [16,20] and *Maja brachydactyla* [21].

The Black Sea *Palaemon adspersus*, Rathke, 1837 is a commercial species of crustaceans living in the Black Sea [22]. In many countries of the world, since ancient times, this type of crustacean has been used for food in boiled and boiled frozen form. Many issues relating to the features of biology, in particular, various stages of the life cycle, migration, nutrition are covered in the works [23-26].

The chemical composition and biological value of this species of crustaceans were studied only in the pre-spawning period for amino acid and fatty acid compositions [27,28]. Information on the mineral component of the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 is limited to indicating their total content [27]. This type of crustacean is in great demand in the Ukrainian market. Therefore, the study of the nutritional value of the mineral composition of meat and the assessment of its contribution to the provision of physiological needs, as well as the determination of safety, represent an urgent task.

The purpose of the work was to assess the nutritional value of the mineral composition of the meat of the Black Sea grass shrimp *Palaemon adspersus* Rathke, 1837 during the fishing periods (spring and autumn), as well as to assess the contribution of the mineral elements of its meat to the satisfaction of human physiological needs.

## Material and Methods

The collection of material was carried out during the spring and autumn fishing of the Black Sea grass shrimp *Palaemon adspersus* Rathke, 1837 in the area of the northwestern coast of the Black Sea of the Odessa region. Fishing was carried out by venters with a mesh size of 2.8 to 8 mm. The total number of analyzed individuals was 650 specimens. After catching shrimp were cut, in meat was isolated, minced meat was cooked and in an average sample in an amount of 300-350g the total ash content was determined by weight after mineralization of the hanging in a muffle oven at a

temperature of 500-600 ° C. By inductively coupled plasma mass spectrometry (ICP-MS) in the samples obtained, the content of the following elements was determined: silver (Ag), aluminum (Al), boron (B), barium (Ba), bismuth (Bi), calcium (Ca), cadmium (Cd), cobalt (Co), chromium (Cr), copper (Cu), iron (Fe), potassium (K), lithium (Li), magnesium (Mg), manganese (Mn), molybdenum (Mo), sodium (Na), nickel (Ni), lead (Pb), zinc (Zn). The analysis was carried out on the atomic emission spectrometer IRIS Interpid II XSP (Termo Fisher Scientific, USA) equipped with a gas-filled cell of the DRC system for the removal of interference and a seven-port dosing valve FAST, as well as an autosator ESISCDX4 (Elemental Scientific Inc., Omaha, NE 68122, USA). librated by external calibration to multi-element standards. Standards containing a full range of detectable elements (0.5, 5, 10 and 50 µg/L) were prepared prior to operation from the Universal Data Acquisition Standards Kit (#N9306225, Perkin Elmer Inc.) by dilution in distilled deionized water acidified with 1% HNO<sub>3</sub>. To account for the incomplete correspondence of sample matrices and calibration solutions for acidity and viscosity, internal standardization «online» for the isotope yttrium-89 was applied in the analysis. An internal standard containing 10 µg/l Y was prepared from the reference standard of yttrium (#N9300167, Perkin Elmer Inc.) on a matrix containing 8% 1-Butanol ((#1.00988, Merck KGaA), 0.8% TritonX-100 (Sigma #T9284 Sigma-Aldrich, Co.), 0.02% TMAH (#20932, Alfa-Aesar, Ward Hill, MA 01835, USA) and 0.02% EDT Acid (Sigma#431788 Sigma-Aldrich, Co). The detection limit was 0.1 mg/kg.

Analysis of the content of mineral elements was carried out in comparison with the literature data published on crustaceans upholstering in various seas [16-21], as well as on marine mollusks [29], fish of the Black Sea [30-32] and other seas [33].

The reliability of the results was analyzed using Microsoft Excel 365 and SPSS 16.0 at P ≤ 0.05.

## Results and Discussion

Analysis of the results of research shows that the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 in the spring and autumn periods of fishing in terms of the total content of mineral elements does not show significant differences: 1.85 ± 0.21 and 1.92 ± 0.34%, % and on average is 1.88 ± 0.27. According to the literature, the total mineral content of other crustaceans, e.g., *Euphausia superba* ranges from 1.9 to 3.2% [16], *Pandalus borealis* from 1.76 [17] to 4.0% [19] and *Pandalus goniurus* to 3.9 ± 0.1% [19].

To assess the nutritional value of the mineral component of shrimp meat, the contribution of 100g of meat per day to meet the physiological needs of an adult, as well as its toxicity in comparison with the literature data was determined (Table 1).

**Essential mineral elements**

Analysis of the research results indicates that the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 contains all essential mineral elements that are determined by mass spectrometry with

Item Name	Black Sea <i>Palaemon adspersus</i> Rathke, 1837, fishing periods		<i>Pandalus borealis</i> [17]	<i>Euphausia superba</i> [16]	Daily level for an adult	
	spring	autumn			The 20th [7]	Toxicity [5]
Essential mineral elements, mg/100g						
*Are	1,89 ± 0,20	1,19 ± 0,10	125,10	124,00	1200	-
K	0,33 ± 0,01	0,34 ± 0,03	281,50	253,00	2500	6000
On	0,33 ± 0,07	0,27 ± 0,01	118,90	313,00	1300	-
*Mg	90,10 ± 7,78	66,10 ± 0,06	125,10	430,00	400	-
*With	2,95 ± 0,51	1,96 ± 0,24	0,35	2,60	1,00	200
*Zn	2,51 ± 0,28	1,93 ± 0,14	2,12	-	12,00	600
Fe	0,92 ± 0,90	0,86 ± 0,06	4,69	4,10	15	200
*Mn	0,62 ± 0,03	0,32 ± 0,01	0,06	-	2,00	40
Cr	0,07 ± 0,003	0,06 ± 0,008	-	0,36	0,005	5
*Nor	0,04 ± 0,001	0,02 ± 0,002	0,03	2,00	0,01	20
*Li	0,12 ± 0,05	0,04 ± 0,003	-	4,50	-	146
With	≤ 0,01	≤ 0,01	-	0,06	-	0,9

**Table 1:** Nutritional value of the mineral composition of the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 during fishing periods.

Note: \*- statistically significant differences at P ≤ 0.05.

inductively coupled plasma. In the spring of the *Palaemon adspersus* Rathke fishery, 1837, the content in its meat of such elements as Ca, Mg, Cu, Zn, Mn, Ni and Li shows higher concentrations in comparison with the autumn period (differences are significant at P ≤ 0.05). The recommended values of the physiological needs of the daily intake of mineral elements are average indicators and reflect the estimated required amount to maintain a normal healthy state of a person [6,7]. Taking into account these recommendations, it follows that the use of 100g of Black Sea shrimp meat is not able to meet human needs for most of the essential mineral elements. Thus, the content of Ca in the spring period of fishing in the meat of *Palaemon adspersus* Rathke, 1837 was 1.89 ± 0.20, in the autumn-1.19 ± 0.10 against the recommended level per day-1200 mg, which is its deficit of more than 99% (Table 1). Ca has been identified in the meat of many hydrobionts, including fish, invertebrates and algae [12-20]. Thus, in the meat of *Pandalus borealis* [17] and *Euphausia superba* [16], the content of Ca was 125.10 and 124.00 mg/100g, respectively. In the meat of the Black Sea shark *Squalus acanthis*, Linnaeus, 1758, 13.84 ± 0.69 mg/100g were detected [32]. Ca is the determining factor for the normal formation of the skeleton and the achievement of peak, genetically predetermined mass and

density [34-36]. The amount of this element both in the meat of the Black Sea *Palaemon adspersus* Rathke, 1837, and in the meat of other crustaceans [16-20] and fish [30-31] is significantly lower than the recommended daily level of its consumption [7].

Essential mineral elements such as K, Na are also found in *palaemon adspersus rathke*, 1837 in low concentrations (0.33 ± 0.01; 0.34 ± 0.03; and 0.33 ± 0.07; 0.27 ± 0.01, respectively) against a higher number in the meat of other crustaceans-*Pandalus borealis* (K-281.50; Na-118,90 mg/100g) [17], *Euphausia superba* (K-253,00; Na-313,00 mg/100g [16]. K is an important dietary mineral and electrolyte that is necessary for the regulation of the body's electrical signals (maintaining cellular polarity, neuronal signaling, transmission of cardiac impulses and muscle contraction), in the transport of nutrients and metabolites, as well as in the activation of enzymes [5,6]. However, its amount in the meat of crustaceans, as well as in the meat of some Black Sea fish (*Squalus acanthis*, Linnaeus, 1758-166.60 ± 8.33 [32] is significantly lower than the recommended daily intake level of 2500 mg/day [7]. Na is important for the effective functioning of the kidneys, nervous and digestive system, for vascular tone, for normal muscle contraction; Together

with K, it takes part in maintaining the normal water-salt balance in cells, regulating the volume of fluid in the body, is an integral part of the ferment, which is responsible for the biosynthesis of energy and the transport of valuable amino acids and glucose to the cells of the body [6,35,36]. The deficiency of these elements in the meat of the Black Sea shrimp is more than 99%.

The concentration of Mg in the meat of *Palaemon adspersus* Rathke, 1837 in the spring fishing period is higher compared to the autumn- $90.10 \pm 7.78$  and  $66.10 \pm 0.06$  ( $P \leq 0.05$ ) and significantly less than the level of this element in other crustaceans (*Pandalus borealis* - $125.10$  mg/100g [16]; *Euphausia superba*- $430$  mg/100g [16]). The amount of Mg in *Euphausia superba* meat exceeds the recommended daily requirement level for this element (400 vs. 430 mg/day) [7,8]. Low concentrations of Mg have been identified in the meat of the Black Sea shark *Squalus acanthis*, Linnaeus, 1758- $15.69 \pm 0.78$  mg/100g [32]. In the meat of such species of commercial fish as *Merluccius bilinearys* from the Gulf of Oman, the mg level was 137. 5-150, 4 mg/g [33]. This element is essential, which is necessary for cell adhesion and migration, energy metabolism, DNA transcription, stability RNA takes part in the synthesis and degradation of numerous neurotransmitters, in particular, catecholamines [38,39].

The level of Cu in the meat *palaemon adspersus* rathke, 1837 during fishing periods exceeds the recommended amount of daily consumption:  $2.95 \pm 0.5$ ;  $1.96 \pm 0.24$  mg/100g against 1.00-1.50 mg [7] (Table 1). *Euphausia superba* meat also showed high concentrations of this element- $2.60$  mg/100g [16], which is consistent with the results of our studies. At the same time, *Pandalus borealis* has an amount of Cu ranging from 0.19 [17] to 0.35 mg/100g [19]. In the meat of *Rapana venosa* in various regions of the Black Sea, from  $1.39 \pm 0.004$  to  $16.72 \pm 2.33$  mg/100g were detected [29], in the tissues of *Mytilus galloprovincialis* up to 3% of the total content of mineral elements [40].

Cu is an important micronutrient necessary for electron transfer processes, is a central component of many enzymes, is involved in energy metabolism and in the cross-linking of collagen and elastin [41-48]. Studies of the balance of consumption of this element have shown that its absorption from the diet is about 50% for all age groups and groups of all stages of human life [5]. The literature on the recommended level of Consumption of Cu is contradictory. The

authors A.V. Skalny [5,49], V.I. Tsipriyan [7], A.V. Pogozheva [8] have instructions to limit the level of daily consumption of Cu. However, at present, sufficiently reliable, sensitive and specific biomarkers of Cu have not been identified., therefore, it is not possible to limit the daily level of consumption of this element [45]. The content of Cu in the meat of *Palaemon adspersus* Rathke, 1837 during fishing periods does not emit the level of toxicity [5] and is safe in the content of this element.

The presence of the trace element Zn in the meat *palaemon adspersus* rathke, 1837 in different periods of fishing does not show significant differences ( $2.51 \pm 0.28$  and  $1.93 \pm 0.14$  mg/100g, in spring and autumn, respectively) and is consistent with data on its content in meat *Pandalus borealis* ( $2.12$  mg/100g) living in the Sea of Okhotsk [17] (Table 1). According to other researchers of the mineral component, *Pandalus borealis* meat from the same region contains Zn up to 0.9 mg/100g [19]. In the Black Sea *Rapana venosa*, higher amounts of this element were detected-up to  $9.12 \pm 2.15$  mg/100g. [29]. The meat of the Black Sea shark *Squalus acanthis*, Linnaeus, 1758 is characterized by the presence of Zn in concentrations: from  $2.94 \pm 0.03$  mg/100g [30], in 14 species of fish from the Gulf of Oman-from 4,512 to 6,421 mg/g [33]. The importance of zinc for human health and life is due to the fact that this element is a cofactor of more than 300 enzymes and more than 100 variants of transcription factors [49-50]. However, this element exhibits toxicity in certain concentrations [51].

The concentration of Fe in the meat of *Palaemon adspersus* Rathke, 1837 during fishing periods does not show statistically significant differences at  $P \leq 0.05$  and averages  $0.88 \pm 0.04$  mg/100g against the recommended amount of 15 mg per day [7] (Table 1). [7], *Euphausia superba* has 4.10 mg/100g [16]. *Rapana venosa* meat has a content of 0.128 to 0.332 mg/100g [29]. Fe is an important element for ensuring the vital activity of all living organisms, the physiological functions of which are due to the entry into the composition of enzymes to ensure the transport of oxygen, electrons for the synthesis of ATP and DNA [7,8,50].

The concentration of Mn in the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 in the spring fishing period is twice as high as in the autumn:  $0.62 \pm 0.03$  against 0.32 mg/100g (Table 1). The level of Mn in the meat of the Ochotomorian *Pandalus borealis* was 0.063 mg/100g [17], in the meat of the Black Sea *Rapana venosa*-

from  $0.34 \pm 0.012$  to  $0.76 \pm 0.032$  mg/100g [29]. This element is involved in the normal development of bone tissue, helps to strengthen the immune system, the proper course of digestive processes, as well as fat and insulin metabolism and brain function [8]. The need for this element for different sources ranges from 2.00 [5] to 4.00 mg/100g [7].

The content of Cr in the meat of *Palaemon adspersus* Rathke, 1837 exceeds 14 and 12 times in the spring and autumn fishing periods, respectively, compared with the recommended values of the daily consumption of this element:  $0.07 \pm 0.003$ ;  $0.06 \pm 0.008$  against 0.005 mg/day (Table 1). High concentrations of Cr were observed in *Euphausia superba*-0.36 mg/100g [16], as well as in *Rapana venosa* meat-from  $0.03 \pm 0.002$  to  $0.42 \pm 0.016$  mg/100g [29], as well as in many species of commercial fish-at the level of 0.066-0.099 mg /g [31-33]. The daily requirement for this element is 0.005 mg/100g [7,8] with a level of toxicity of 5 mg [5]. Cr is considered an important essential trace element, because it contributes to the structural integrity of nucleic acid molecules; participates in the regulation of the heart muscle and the functioning of blood vessels; promotes the removal of toxins, salts of heavy metals, radionuclides from the body [53,54]. However, to date, the mechanisms of these Cr functions in metabolism have not been conclusively substantiated. The bioavailability of chromium from inorganic compounds in the gastrointestinal tract is low, only 0.5-1%, and increases to 20-25% with the intake of chromium in the form of complex compounds (picolinate, asparaginate) [52]. Taking into account these data, the identified content of Cr in the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 does not pose a danger to human health.

The level of Ni in the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 in the spring fishing period is twice as much as in the autumn:  $0.04 \pm 0.001$  and  $0.02 \pm 0.002$  mg/100g, respectively, and exceeds the recommended values-0.01 mg [7], but not toxicity [5,55]. According to the literature data, Ni, coming from food, is absorbed mainly in the small intestine and its bioavailability is 1-10% [5,56].

It is known that one of the main functions of Ni is an indirect participation in the process of hematopoiesis through the mechanism of penetration of iron into the blood [57]. It has also been shown that Ni is a cofactor (activator) of bioligand, capable of

binding iron and transferring it from the insoluble form of  $Fe^{3+}$  to easily digestible  $Fe^{2+}$ , is also included in the composition of blood cells-erythrocytes, participates in the activation of certain enzymes and in redox processes organism [58,59]. Its role in ensuring the necessary conformation and shape of molecules in the structural organization and functioning of DNA, RNA, proteins is shown [56]. A toxic dose of nickel with a single dose is 50 mg, and with a daily dose-20 mg/day; data on the lethal dose are not available [60]. The negative effect of Ni can be manifested in a decrease in immunity, the manifestation of allergic [59], stress reactions and depression of the cardiovascular system [61]. In the human gastrointestinal tract, from 1 to 10% of the received Ni is absorbed. According to our data, when Ni enters the body with an average of 0.33 mg/100g of shrimp and the absorption of 10% in the human body, 0.03 mg will be absorbed (Table 1), which is significantly lower than the toxicity threshold for this element [5].

The content of Li in the meat of the spring shrimp catch exceeds its amount in the autumn by 3 times:  $0.12 \pm 0.05$  against  $0.04 \pm 0.003$  mg/100g (Table 1). In other crustaceans, this element is present in higher concentrations-4.50 mg/100g [16] or is not detected [17-20]. In the meat of many species of commercial fish, this element has not been determined [28-32].

Li is important for the proper functioning of a number of enzymes, hormones, vitamins, growth factors, the immune system, and the nervous system [62]. This mineral also has a wide range of neuroprotective effects [63-65]. The recommended level of daily intake of Li has not yet been established, but there is data on the level of toxicity-146 mg per day [5].

The content of Co in the meat of *Palaemon adspersus* Rathke, 1837 does not show differences depending on the catch season and is  $\leq 0.01$  mg/100g (Table 1). In the meat of other crustaceans, for example, *Pandalus borealis*, this element was not detected [17], while in *Euphausia superba* its content is established at 0.90 mg/100g [16]. This element is a vital and indispensable component of many enzymes and coenzymes [66-68]. In the middle, the intake of Co ions into the human body with food is 0.012 mg/day and the expression of the physiological level of Co consumption can cause toxic effects: activation of inflammatory processes, mutagenesis, carcinogenesis, tissue necrosis [69].



**Toxic and little-studied elements**

One of the groups related to toxic elements are heavy metals: Pb, Cd, As and Hg, which in certain concentrations can have harmful effects on the human body, capable of accumulate in the tissues, causing a number of diseases [70,71]. These elements enter the ocean through the atmosphere and with the burial of multi-faceted waste in the oceans [72,73].

The assessment of the content of heavy metals in the meat of the Black Sea *Palaemon adspersus* Rathke, 1837 during fishing periods and their compliance with permissible levels [70] is presented in table 2.

Pb is found in the meat of *Palaemon adspersus* Rathke 1837 in amounts  $\leq 0.01$  mg/100g (Table 2). These data are consistent with

Name of the element	Heavy metals, mg/kg					Acceptable levels, not more than [70]
	<i>Palaemon adspersus</i> Rathke 1837, fishing period		<i>Pandalus borealis</i> [17]	<i>Pandalus goniurus</i> [19]	<i>Euphausia superba</i> [16]	
	spring	autumn				
Pb	< 0.01	< 0.01	0,15	0,01	0,10	2,0
Cd	0,17 ± 0,01	< 0.01	0,50	0,02	0,01	0,5
As	0,24 ± 0,02	0,24 ± 0,02	2,03	2,57	0,50	0,5
Hg	< 0.01	< 0.01	0,17	0,11	0,01	0,1

**Table 2:** Assessment of the compliance of the heavy metals Pb, Cd, As and Hg (mg/kg) in *Palaemon adspersus* Rathke 1837 meat with their permissible levels [70].

the Pb content of *Pandalus goniurus* meat [19], while in the meat of other crustaceans (*Euphausia superba* and *Pandalus borealis* [16,17] this element is present 15 and 10 times, respectively. High concentrations of lead (from 0.03 to 0.10 mg/100g) are found in the meat of many commercial Black Sea Fish [30-32] and crustaceans [74]. However, the level of this element in the meat of hydrobionts is significantly lower than its permissible concentrations: 0.003-0.005 mg/100g [70]. The toxic effect of Pb is manifested in a negative effect on human mental abilities [7,8]. It is excreted from the body by 90% [5].

The content of Cd in the meat of *Palaemon adspersus* Rathke 1837 in the spring fishing period is higher compared to the autumn period- $0.017 \pm 0.001$  against  $< 0.001$  mg/100g (the differences are significant at  $P \leq 0.05$ ). In *Pandalus borealis* meat, 0.0063 mg/100g were detected [17], in *Euphausia superba*-from  $0.4 \pm 0.04$  to  $1.3 \pm 0.1$  mg/100g [16]. Significant accumulations of this element in the human body lead to increased pressure, renal failure and chronic poisoning [5,75,76]. This element is found in the meat of many crustaceans [13-21] and fish [30-33], however, in most cases, as for the Black Sea *Palaemon adspersus* Rathke 1837 its level is lower than toxicity [5].

The concentration of As in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 of the spring and autumn catch periods does not show any differences and is  $0.002 \pm 0.0001$ , which is more than two times less than the recommended daily intake level (0.005 mg/day) [70]. As is found in the meat of all hydrobionts, but the level of its content differs in all species. In meat, *Euphausia superba*-0.06 [16], *Pandalus borealis*-0.002 and *Pandalus goniurus*-0.002 mg/100g, respectively [19]. As refers to toxic elements, the intake of which in high concentrations can lead to acute or chronic intoxication, and in some cases to the development of malignant neoplasms [4,5,65].

The level of Hg in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 does not show differences depending on the catch period and is  $\leq 0.01$  mg/kg, i.e., significantly lower than the permissible level [7,8,70]. In other crustaceans, this element was detected in higher concentrations: *Pandalus goniurus*-0.11; *Pandalus borealis*-0.17 mg/kg [19]. Low Hg content (0.012-0.049 mg/g) is determined in the meat of many commercial fish species [31-33]. Hg is one of the elements constantly present in

the environment and living organisms [71-74]. It is a highly toxic cumulative poison that affects the hematopoietic, enzymatic, nervous systems and kidneys [75-76]. FAO and the World Health Organization have established a weekly safe intake of total Hg-5 µg per kilogram of human body weight [70], toxic effect occurs at 0.4 mg/day [5]. Thus, the level of Hg in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 is not dangerous for human health.

**Toxic and malo-studied mineral elements**

Toxic and poorly studied mineral elements include Al, Ag, Ba, Bi and B [5], data on the content of which in the meat *palaemon adspersus* rathke 1837 during fishing periods are presented in table 3.

The level of Al reveals statistically significantly large values in the spring period of the Black Sea *Fishery Palaemon adspersus* Rathke 1837 (0.41 ± 0.06 against 0.26 ± 0.03, mg/100g in autumn) (Table 3).

Item Name	Catch periods <i>Palaemon adspersus</i> Rathke 1837		Permissible level, mg [5]
	Spring	Autumn	
To the	0,41 ± 0,06	0,26 ± 0,03	-
At the	0,03 ± 0,003	0,02 ± 0,001	0,05
Three	0,63 ± 0,01	0,36 ± 0,04	-
Would	≤ 0,001	≤ 0,001	-
B	≤ 0,001	≤ 0,001	-

**Table 3:** The content of toxic and poorly studied mineral elements, mg/ 100g in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 during fishing periods.

Al refers to toxic elements, the effect of which can be manifested in the hydronephrotic transformation of the kidneys, the expansion of the urinary ducts, difficulty urinating and/or the formation of cameos, as well as in the genotoxic effect on the chromosomes of bacterial cells, cells of warm-blooded animals and neurotoxic action [78-80].

Aluminum compounds have been shown to dissolve with the release of free Al<sub>3+</sub> ions when released into the acidic environment of the stomach [81,82]. The absorption of various aluminum compounds in the intestine is at a level of 0.01 to 0.5% of the amount consumed [80]. The Joint FAO/WHO Expert Committee on Food Additives (JECFA) has established a new level of safe weekly consumption of aluminum PTWI, which is 0-2.0 mg/kg by weight. Thus, the aluminum content in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 from the standpoint of the established level of safety is within acceptable values.

The level of Ag in the meat of *Palaemon adspersus* Rathke 1837 is higher in the spring fishing period and is 0.03 ± 0.003 mg/100g against 0.02 ± 0.001 (Table 3). In other crustaceans [16-20], mollusks [29] and fish [30,32] this element was not detected. Ag is

important for the human body [7,8], but at a dose of 10g, a lethal outcome is likely [5]. According to the content of this trace element, the meat of *Palaemon adspersus* Rathke 1837 is safe during fishing periods.

The concentration of Ba, as well as Ag and Al in the spring period of fishing is higher compared to the autumn: 0.63 ± 0.01 against 0.36 ± 0.03 (at P ≤ 0.05) (Table 3). In the meat of other crustaceans [16-20], Black Sea rapana [29] and fish [30-32] this trace element was not determined. The physiological and biochemical significance of barium for the body is not well understood [5]. Ba is a synergist of acetylcholine and in low concentrations exhibits the properties of the essential element of vertebrates, without the participation of which the correct functioning of nervous activity is impossible [83,84]. However, high concentrations of Ba are able to block potassium channels not only of neurons, but of all cellular life forms in general [84]. A day in the human body fasts 0.3-0.9 mg of Ba and in general contains 20-22 mg of this heavy metal, which enters with food and water [5]. The toxic dose for humans is 200 mg of Ba per day, the lethal dose varies between 0.8-3.7g. [6]. The content of Va in the meat of the Black Sea during fishing periods does not pose a danger to humans.

The concentration of boron B in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 does not depend on the catch season and is  $< 0.001\text{g}/100\text{g}$  (Table 3). In crustaceans [15-19] and other hydrobionts, this element was not detected [28-32]. It affects the regulation of parathyroid hormone activity and indirectly takes part in the metabolism of calcium and magnesium (prevents their loss through the kidneys), fluoride, phosphorus and vitamin D [48]. As a result, it plays a regulatory role in the formation of bone tissue, prevents the development of osteoporosis, improves the absorption of calcium by bone tissue, is necessary for the normal formation of the skeleton in children, and in older age groups it prevents diseases of the spine and joints (osteochondrosis, arthrosis, arthritis) [52,53]. It is established that healthy adults can consume 1-13 mg of boron with food [5].

Vi was detected in the meat of the Black Sea grass shrimp in the amount of  $\leq 0.001\text{ mg}/100\text{g}$  during fishing periods (Table 3). The current level of knowledge does not allow us to speak definitively about any physiological role of bismuth in the human body [5,6]. It belongs to the category of heavy metals; it is a moderately toxic element.

Thus, studies of the mineral composition of the meat of the Black Sea *Palaemon adspersus* Rathke 1837 showed the presence of all essential, low-toxic, toxic and poorly studied mineral elements. In the spring period of the fishery, statistically significant higher concentrations of elements such as Ca, Mg, Cu, Zn, Mn, Ni were established, Li, Cd, Al, Ba (Table 1-3). This is probably due to the fact that the North-Western part of the Black Sea (NWFM), where fishing and shrimp sampling was carried out, is the most hydrologically dynamic area of the sea [85]. The Dniester, the Dnieper and the Southern Bug, the total flow of which is almost 80% of the total flow into the Black Sea and lead to high chemical pollution of water and bottom sediments in the area of the Odessa coast of the Black Sea in the spring period of the year [86]. It has also been shown that the effluents of these rivers, especially the Danube, are the most important sources of heavy metals. Nevertheless, the accumulation of toxic elements in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 is at an acceptable level and meets the safety requirements [70].

## Conclusion

The mineral composition of the meat of the Black Sea grass shrimp in the spring and autumn periods of fishing in the Black Sea has been studied. The level of essential mineral elements (Ca, K, Na, Mg, Zn, Fe, Mn) is significantly lower than the physiological needs of humans. Statistically significant higher concentrations of Mg, Cu, Mn, Ni, Cd, Al, Ba, in the spring catch period compared to the autumn catch were revealed. However, the level of toxicity of such elements as As, Cd, Pb, Hg, in the meat of the Black Sea *Palaemon adspersus* Rathke 1837 in different fishing periods is significantly lower than the permissible values. Thus, in terms of safety, the meat of the Black Sea *Palaemon adspersus* Rathke 1837 meets the safety requirements and can be used for food purposes.

There are also poorly studied elements, such as Ag, Ba, Bi, the physiological role of which is in the research stage.

The results of the research indicate the need to monitor the mineral component of shrimp meat in the fishing periods of the year, as well as to study the effect of heat treatment regimes on the mineral status of the meat of the Black Sea grass shrimp.

## Bibliography

1. OSHA. Occupational safety and health standards. Limits for air contaminants". Washington, DC: Occupational Safety and Health Administration. 29 CFR 1910.1000, Table Z-1 (2003).
2. Bost M., *et al.* "Literature search and review related to specific preparatory work in the establishment of Dietary Reference Values for Copper (Lot 3)". Project developed on the procurement project CT/EFSA/NUTRI/2011/01. EFSA Supporting publication EN-302 (2012): 63.
3. Dietary Reference Values for nutrients Summary report European Food Safety Authority (EFSA) Update: 4 September TECHNICAL REPORT Approved (2019).
4. Afshin A., *et al.* "Health effects of dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease study 2017". *Lancet* (2019): 1958-1972.
5. Skalny AV. «Chemical elements in human physiology and ecology». M. Izd. Onyx 21st Century House: The World (2004): 216.



6. Oberlis D., et al. "The biological role of macro- and microelements in humans and animals». S-Pb: Nauka (2008): 543.
7. Tsypryan VI., et al. "Food hygiene with the basics of nutriology». book 1.- All-Ukrainian specialty «Medicine», Kyiv (2007): 527 cc.
8. Pogozheva AV., et al. "Healthy nutrition. The role of dietary supplements». M. GEOTAR - Media (2020): 480.
9. Fleming L., et al. The oceans and human health. In: Shugart HH, editor. "Oxford Research Encyclopedia of Environmental Science". Oxford: Oxford University Press (2015).
10. Charrondiere UR., et al. "FAO/INFOODS Food Composition Database for Biodiversity". *Food Chemistry* 140.3 (2013): 408-412.
11. Mayer AMS., et al. "Marine Pharmacology in 2016-2017: Marine Compounds with Antibacterial, Antidiabetic, Antifungal, Anti-Inflammatory, Antiprotozoal, Antituberculosis and Antiviral Activities; Affecting the Immune and Nervous Systems, and Other Miscellaneous Mechanisms of Action". *Marine Drugs* 19.2 (2021): 49.
12. World Health Organization. "Ambition and Action in Nutrition 2016-2025". World Health Organization (2017).
13. Amiard JC., et al. "Bio accessibility of Essential and Non-Essential Metals in Commercial Shellfish from Western Europe and Asia". *Food Chemistry and Toxicology* 46 (2008): 2010-2022.
14. Holthuis LB. "FAO species catalogue. Shrimps and prawns of the world". FAO Fisheries Synopsis 125.1 (1980): 271.
15. Méndez L., et al. "Mineral Concentrations in Muscle and Hepatopancreas of Newly Caught Wild and Hatchery-Exhausted Spawners of Pacific White Shrimp, *Penaeus vannamei*". *Journal of Applied Aquaculture* 8.4 (1999): 17-26.
16. Antarctic Krill: Handbook / Ed. by VM.Bykova. - M. Izd-vo VNIRO (2001): 207.
17. Paulov YuV., et al. "Technochemical characteristics and technological features of promising species of Okhotsk Sea shrimp». *Izvestiya TINRO* 140 (2005): 291-301.
18. Zyuzgina AA., et al. "Comparative characteristics of the total chemical composition and activity of the proteases of the northern *Pandalus borealis*, the angle-tailed *Pandalus goniurus* and the Antarctic *Euphausia superba* shrimp». *Izvestiya TINRO* 147 (2006): 354-360.
19. Yarochkin AP., et al. "Biotechnology of processing small shrimps for use in food products». *Izvestiya TINRO, Technology of Processing Hydrobionts* 200.2 (2020): 460-485.
20. Handbook on the chemical composition and technological properties of algae, invertebrates and marine mammals/Ed. V. A. Bykova. M. VNIRO (1999): 262.
21. Marques A., et al. "Chemical composition of Atlantic spider crab *Maja brachydactyla*: Human health implications". *Journal of Food Composition and Analysis* 23.3 (2010): 230-237.
22. Boltachev AA., et al. "Black Sea herbal shrimp *Palaemon adspersus* (Decapoda, Palaemonidae). Biology, Fishery, Problems». *Journal of Voprosy Ribolovstvo* 18.3 (2017): 313-327.
23. Makarov YuN. "Fauna of Ukraine. Ten-legged crustaceans". Kyiv: Nauk. Dumka (2004): 430.
24. Lisitskaya LA. «Morphometric characteristics of shrimp *Palaemon adspersus* and *Palaemon elegans* (Palaemonidae) from the Black Sea waters of the south-western Crimea (M. Kaya-Bash and Balaklava Bay)». *Uch. Zap. TNU. Ser. Biology, Chemistry* 25.64 (2012): 109-114.
25. Wiktor K. "The food composition of *Palaemon adspersus* (Rathke) from Puck Bay Zesz. Nauk. BINOZ UG". 6 (1979): 145-154.
26. Lapinska E., et al. "Seasonal variations in the occurrence of the prawns *Crangon crangon* (L., 1758), *Palaemon adspersus* (Rathke, 1837) and *Palaemon elegans* (Rathke, 1837) in the littoral zone of the Gulf of Gdańsk". *Oceanological and Hydrobiological Studies* 34.2 (2005): 95-110.
27. Bal-Prilipko LV., et al. "Nutritional and biological value of the Black Sea herbal shrimp *Palaemon adspersus*». *G. Food Industry of the Agro-Industrial Complex* 5 (2018): 28-31.

28. Lebskaya TK., et al. "Lipid profile of the Black Sea herbal shrimp *Palaemon adspersus* Rathke, 1837". G. Nutrition Issues. *Vopr Pitan* 89.1 (2020): 96-100.
29. Apache MV., et al. "Safety of black sea mollusks in terms of the content of heavy metals". *Commodity Newsletter* 9 (2016): 92-100.
30. Omelchenko SO., et al. "The content of heavy metals in the tissues of some Chernomo fish and their effect on the level of oxidative modification of proteins». Scientific notes of the Tauride National University named after V.I. Vernadsky". *Series «Biology, Chemistry»*. 20.59 (2007): 59-64.
31. Boldyrev DA., et al. "Content and distribution of chemical elements in the tissues of the Black Sea merlangus *Merlangius Merlangus Euxinus*». *Ribne hospodarstvo Ukrainy* 6 (2011): 44-48.
32. Sydorenko O., et al. "Characteristics of the safety of meat of the Black Sea shark katran in the content of heavy metals». Internar. Science pact. Journ. «Goods and Markets». 2.20 (2015): 124-132.
33. Al-Amri I., et al. "Determination and comparison of non-essential and essential elements in different species of fish available in Oman markets by using inductively coupled plasma-optical emission spectrometry". *Acta Scientific Nutritional Health* 5.1 (2021): 60-73.
34. Galchenko AV., et al. "Calcium status among vegetarians and vegans". Russian scientific-practical conference with international participation". *Fundamentals of Technological Development of Agriculture. Orenburg* (2019): 209-212.
35. Areco V., et al. "Dietary and pharmacological compounds altering intestinal calcium absorption in humans and animals". *Nutrition Research Reviews* 28.2 (2015): 83-99.
36. Martinchik AN., et al. "Calcium in the diet of preschool and school-age children: the main food sources and factors affecting consumption». *Question Food* 87.2 (2018): 24-33.
37. Oria M., et al. "Dietary Reference Intakes for Sodium and Potassium" Appendix J, Dietary Reference Intakes Summary Tables. National Academies of Sciences, Engineering, and Medicine; Health and Medicine Division; Food and Nutrition Board; Committee to Review the Dietary Reference Intakes for Sodium and Potassium". Washington (DC): National Academies Press (US) (2019).
38. Bobrova EV. «Magnesium metabolism and antihypertensive efficacy of beta-blockers». *Ukrainian Journal of Cardiology* 1 (2003): 75-79.
39. Revko OP., et al. "Physiological role and importance of magnesium in the therapy of internal diseases». *Bulletin of the Club of Pancreatologists* 2 (2010): 60-64.
40. Pospelova VN., et al. "Copper content in organs and tissues of *Mytilus galloprovincialis* Lamarck, 1819 and the flow of its sedimentation into bottom sediments in the farms of the Black Sea aquaculture". *Marine Biological Journal* 3.4 (2015): 64-75.
41. Araya M., et al. "CCS mRNA transcripts and serum CCS protein as copper marker in adults suffering inflammatory processes". *Biomaterials* 27 (2014): 645-652.
42. EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2015h. Scientific Opinion on Dietary Reference Values for copper". *EFSA Journal* 13.10 (2015): 4253-4351.
43. Keskin Y., et al. "Cadmium, lead, mercury and copper in fish from the Marmara Sea, Turkey". *Bulletin of Environmental Contamination and Toxicology* 78 (2007): 258-261.
44. de Romana DL., et al. "Risks and benefits of copper in light of new insights of copper homeostasis". *Journal of Trace Elements in Medicine and Biology* 25 (2011): 3-13.
45. Riley MR., et al. "Effects of metals Cu, Fe, Ni, V, and Zn on rat lung epithelial cells". *Toxicology* 190.3 (2003): 171-184.
46. Hänsch R., et al. "Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl)". *Current Opinion in Plant Biology* 12.3 (2009): 259-266.

47. Skalny AQ and IA Rudakov. «Bioelements in Medicine». M.: Mir (2004): 272.
48. Riley MR., et al. "Effects of metals Cu, Fe, Ni, V, and Zn on rat lung epithelial cells". *Toxicology* 190.3 (2003): 171-184.
49. Mazur AA., et al. "Evaluation of the toxic effects of zinc ions and zinc oxide nanoparticles on the early development of the sea urchin *Scaphechinus mirabilis* (Agassiz, 1864) (Echinodermata: Echinoidea)". *Sea Biology* 46.1 (2020): 53-59.
50. Babenko GA. «Human trace elements: pathogenesis, prevention, treatment». *Trace Elements in Medicine* 2.1 (2001): 2-5.
51. Reutina SV. «The role of chromium in the body». Bulletin of the Peoples' Friendship University of Russia". *Series: Ecology and Life Safety* 4 (2009): 50-55.
52. Panchenko LF. «Clinical biochemistry of microelements». M.: GOU VUNMC MZ RF (2004).
53. Ragsdale SW. "Nickel-based enzyme systems". *Journal of Biological Chemistry* 284 (2009): 18571-18575.
54. Galchenko AV and AA. Sherstneva. «Conditionally essential ultramicroelements in the diet of vegetarians and vegans: nickel, lithium, vanadium, germanium». *A. Trace Elements in Medicine* 22.2 (2021): 3-16.
55. Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for Nickel. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service (2005).
56. Pandey R and SP Srivastava. "Spermatotoxic effects of nickel in mice". *Bulletin of Environmental Contamination and Toxicology* 64.2 (2000): 161-167.
57. "Agency for Toxic Substances and Disease Registry (ATSDR)". Toxicological profile for Nickel. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service (2005): 397.
58. Andrew A and A Barchowsky. "Nickel-induced plasminogen activator inhibitor-1 expression inhibits the fibrinolytic activity of human airway epithelial cells". *Toxicology and Applied Pharmacology* 168.1 (2000): 50-57.
59. Cavallo D., et al. "Evaluation of oxidative damage and inhibition of DNA repair in an in vitro study of nickel exposure". *Toxicology in Vitro* 17.5-6 (2003): 603-307.
60. Dasika UK., et al. "Nickel allergy to the percutaneous patent foramen ovale occuder and subsequent systemic nickel allergy". *The Journal of Thoracic and Cardiovascular Surgery* 126.6 (2003): 2112.
61. Chen CY., et al. "Nickel-induced oxidative stress and effect of antioxidants in human lymphocytes". *Archives of Toxicology* 77.3 (2003): 123-130.
62. Demling JH., et al. "On the physiological function of lithium from a psychiatric view point". *Medical Hypotheses* 57.4 (2001): 506-509.
63. Rohayem J., et al. "Predictors of prophylactic response to lithium". *Encephale* 34.4 (2008): 394-399.
64. Kjølholt J., et al. "The elements in the second Rank-Lithium". Copenhagen, Denmark: Miljoministeriet (2003): 108.
65. Leysens L., et al. "Cobalt toxicity in humans - a review of the potential sources and systemic health effects". *Toxicology* 387 (2017): 43-56.
66. M Yilmaz., et al. "Toxic effects of cobalt parahydroxybenzoate on tissue histopathology and serum proteins in Capoeta capoeta capoeta". *Fresenius Environmental Bulletin* 17.9 (2008): 1322-1327.
67. Kumanto M., et al. "Cobalt (II) Chloride Modifies the Phenotype of Macrophage Activation". *Basic and Clinical Pharmacology and Toxicology* 121 (2017): 98-105.
68. Dolomatov SI., et al. "Modern aspects of regulatory, pathophysiological and toxic effects caused by cobalt ions during oral intake into the human body". *Analysis of Health Risk* 3 (2019): 161-174.
69. CODEX GENERAL STANDARD FOR CONTAMINANTS AND TOXINS IN FOOD AND FEED (CODEX STAN 193-1995).

70. Saghiri MA., *et al.* "Functional role of inorganic trace elements in angiogenesis part III: (Ti, Li, Ce, As, Hg, Va, Nb and Pb)". *Critical Reviews in Oncology/Hematology* 98 (2016): 290-301.
71. Patin SA. «The Impact of Pollution on the Biological Resources and Productivity of the World Ocean». M.: Pisheskaya obshchestvo (1979): 250 s.
72. Sayenko GN. «Metals and Halogens in Marine Organisms». M.: Nauka (1993): 252.
73. Olgunoglu MP., *et al.* "Heavy metal concentrations (Cd, Pb, Cu, Zn, Fe) in Giant Red Shrimp (*Aristaeomorpha foliacea* Risso 1827) from the Mediterranean Sea". *Polish Journal of Environmental Studies* 24. 2 (2015): 631.
74. Malov AM and ML Alexandrova. «Medical and ecological aspects of mercury contamination in the conditions of a megalopolis". *Ecology* 10.4 (2009): 102-112.
75. Clarkson TW., *et al.* "The toxicology of mercury and its chemical compounds". *Critical Reviews in Toxicology* 36 (2006): 609-620.
76. Thomas., *et al.* "Immuno-allergological properties of aluminum oxide (Al<sub>2</sub>O<sub>3</sub>) ceramics and nickel sulfate in humans". *Biomaterials* 24.6 (2003): 959-966.
77. Bagrntseva OV., *et al.* "Aluminium: food-related health risk assessment of the consumers". *Health Risk Analysis* 1.13 (2003): 59-68.
78. Mujika J., *et al.* "Aluminium in Biological Environments: A Computational Approach". *Computational and Structural Biotechnology Journal* 9.15 (2014): 1-13.
79. Aguilar F., *et al.* "Safety of aluminium from dietary intake. Scientific Opinion of the Panel on Food Additives, Flavourings, Processing Aids and Food Contact Materials (AFC)". *The EFSA Journal* 754 (2008): 1-34.
80. Aluminium (from all sources, including food additives)// Evaluation of certain food additives and contaminants: sixty-seventh report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report 940 (2007): 33-45.
81. Ranau R., *et al.* "Aluminum Levels of Fish Fillet Baked and Grilled in Aluminum Foil". *Food Chemistry* 73 (2001): 1-6.
82. Chashchin V., *et al.* "Medical and ecological aspects of the relationship of disorders of human functional systems with the content of trace elements barium and strontium in the body. Literature Review». *Human Ecology* 4 (2019): 39-47.
83. Ribera AB and N Spitzer. "Both barium and calcium activate neuronal potassium currents (calcium-dependent potassium current/transient potassium current/tetraethylammonium ion/whole-cell voltage clamp/ subtraction protocol)". *Proceedings of the National Academy of Sciences of the United States of America Neurobiology* 84 (1987): 6577-6581.
84. Dyatlov SE., *et al.* "Toxicological characteristics of drainage and casting waters of the Odessa coast". of the Sciences. Zap. Ternopil National University". pedagogical university named after Volodymyr Hnatiuk 3.4 (2015): 203-207.
85. Dyatlov SYe. "Heavy metals in water and bottom sediments of Odessa region of the Black Sea". *Journal of Shipping and Ocean Engineering* 5 (2015): 51-58.
86. Ilyin YuP. «Hydrological structure of waters in various wind situations according to polygon and satellite observations. Natural conditions of the danube river and snake islands: the current state of the ecosystem». Ed. Ivanov V.A., Goshovskogo S.V. - Sevastopol: MGI NASU (1999): 134-146.

#### Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: [www.actascientific.com/](http://www.actascientific.com/)

Submit Article: [www.actascientific.com/submission.php](http://www.actascientific.com/submission.php)

Email us: [editor@actascientific.com](mailto:editor@actascientific.com)

Contact us: +91 9182824667