

## Occupational Exposure to Mycotoxin-Producing Fungi in Respirable and Surface Dust of Farms and Feed Factories Located in the State of São Paulo, Brazil: A Pilot Study

Larissa Tuanny Franco<sup>1</sup>, Tatiana Alves Reis<sup>2</sup> and Carlos Augusto Fernandes de Oliveira<sup>1\*</sup>

<sup>1</sup>Departamento de Engenharia de Alimentos, Faculdade de Zootecnia e Engenharia de Alimentos, Universidade de São Paulo, Pirassununga, SP, Brazil

<sup>2</sup>Departamento de Microbiologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, SP, Brazil

**\*Corresponding Author:** Carlos Augusto Fernandes de Oliveira, Departamento de Engenharia de Alimentos, Faculdade de Zootecnia e Engenharia de Alimentos, Universidade de São Paulo, Pirassununga, SP, Brazil.

**DOI:** [10.31080/ASNH.2022.06.1094](https://doi.org/10.31080/ASNH.2022.06.1094)

**Received:** July 04, 2022

**Published:** July 11, 2022

© All rights are reserved by **Carlos Augusto Fernandes de Oliveira., et al.**

### Abstract

In the present study, a preliminary investigation was conducted on the occurrence of mycotoxin-producing fungi in respirable air and surface dust from farms (N = 4) and feed factories (N = 3) located in the state of São Paulo, Brazil. The main genera detected were *Aspergillus* sp., *Fusarium* sp., *Penicillium* sp., and *Cladosporium* sp. *Aspergillus* sp. was identified in more than 50% of samples analyzed. The values of colony forming units (CFU) were higher in feed factories than in farms, with a maximum of 7.10 Log CFU/m<sup>2</sup>. In samples of respirable dust, average values were 3.12 Log CFU/m<sup>3</sup> in animal feed factories and 2.95 Log CFU/m<sup>3</sup> in farms. The study demonstrated that the levels of mycotoxin-producing fungi found in the work environments studied may be of concern to the health of the workers.

**Keywords:** Fungi; Occurrence; Workplace; Breathable Dust; Settled Dust

### Abbreviations

CFU: Colony-Forming Unit; NSF: Non-Sporulating Fungus

### Introduction

Fungi contamination of workplaces has attracted considerable attention in the last decades, considering the health concerns regarding human exposure to pathogenic and toxigenic species. Previous studies have shown the occurrence of occupational exposure to mycotoxigenic fungi in rural workers from several countries, mainly among individuals involved in animal production, especially pig farmers and poultry breeders, evidencing high rates of asthma and inflammatory manifestations in the respiratory system [1-4]. In recent years, animal management techniques start to involve closed facilities, densely packed with animals, resulting in

indoor air with high levels of dust, gases, microbes, and microbial metabolites, which increase potential health risks [5,6]. Cote., et al. [7] presented experimental evidence that, in humans, inhalation of fungal spores can cause asthma and rhinitis, while dust contaminated with mycotoxins had immunosuppressive effects. In addition, epidemiological studies have revealed high frequency of liver, lung, and biliary tumors in workers handling food contaminated with mycotoxins [8]. Thus, inhaling dust contaminated with certain mycotoxins can be very harmful to the health of workers and can potentiate adverse effects by means of combined exposure to multiple mycotoxins [9].

The main genera of mycotoxin-producing fungi are *Aspergillus* sp., *Fusarium* sp., and *Penicillium* sp. [10]. Aflatoxins are mainly

produced by *A. flavus*, *A. parasiticus*, and *A. nomius* [8]. Fumonisin are produced by *F. verticillioides*, *F. proliferatum*, and *A. niger*, usually [11]. Trichothecenes and zearalenone are produced by *F. graminearum* and *F. culmorum* [12], and ochratoxin may be produced by *A. ochraceus*, *A. niger*, *A. carbonarius*, and *P. verrucosum* [13]. Recently, studies have demonstrated possible occupational exposure to mycotoxins resulting from the inhalation of contaminated dust in work environments in several countries [14-17]. However, to our knowledge, this issue has never been addressed in rural working environments in Brazil.

### Aim of the Study

The present study aimed to conduct a preliminary investigation on the presence of mycotoxin-forming fungi genera in respirable air and surface dust from Brazilian farms and feed factories.

### Material and Methods

Four sampling procedures were conducted in 7 locations in the state of São Paulo - Brazil, totaling 4 farms (2 poultry-producing farms, 1 pig-producing farm, and 1 dairy farm), and 3 feed factories. Sample collections started in October 2020 and ended in May of the following year. A total of 78 samples were collected, 26 of respirable dust and 52 surface swabs. Collection of breathable dust samples was performed with particles of size from 0.5 to 10  $\mu\text{m}$  (PM<sub>0.5</sub> a PM<sub>10</sub>), as described by Viegas, *et al.* (2013a, 2013b), with modifications. This is in accordance with the sampling procedures as recommended by Vincent and Mark [18], who indicated that respirable dust is a fraction of the dust transported by air that reaches gas exchange regions in the lungs, with aerodynamic diameter of less than 7  $\mu\text{m}$ .

For collection of breathable dust, an Accura-2 Criffer air sampling pump was used, accompanied by a support for a membrane cassette with a flow rate of 2.50 L/min placed at a height close to the nasal region of the workers. Collection was carried out continuously for 8 hours. Fungal contamination of samples was analyzed on Dicloran Rose Bengal Chloramphenicol (DRBC) agar plates at dilutions of  $10^{-1}$  and  $10^{-2}$ . Surface dust samples were collected by rubbing the surfaces using a sterile dehydrated cellulose sponge with buffer solution, in accordance with the International Standard ISO 18593-2004 (ISO, 2004). Smears were placed in DRBC agar plates; 100 mL of peptone water were added to the sponge and stirred for a minute in a homogenizer. Dilutions of  $10^{-2}$ ,  $10^{-3}$  and  $10^{-4}$  were made for plate analysis.

All DRBC agar plates of collected breathable and surface dust samples were incubated at 27°C for 10 days. After incubation, results were determined, both quantitatively (in colony forming units, CFU/m<sup>3</sup>, CFU/m<sup>2</sup>) and qualitatively, with identification of the fungi genera of fungus. All colonies were subcultured on potato dextrose agar (Oxoid, Basingstoke, UK) at 25°C for 7 days. Fungus identification was performed by the classical method using macroscopic and microscopic characteristics [19]. Hyaline fungi were colored by lactophenol cotton blue and observed in a light microscope (400x).

### Results and Discussion

Table 1 presents the main fungi identified in the four samplings from farms and feed factories located in the state of São Paulo, Brazil. High concentrations of fungi were detected on surface and breathable dust from both sampling sites. Average count found on the surface dust of feed factories (5.99 Log CFU/m<sup>2</sup>) was higher than in farms (5.26 Log CFU/m<sup>2</sup>). However, maximum values detected were very similar, with 7.10 Log CFU/m<sup>2</sup> and 7.04 Log CFU/m<sup>2</sup> in feed factories and farms, respectively. In samples of respirable dust, mean values of CFU observed in factories (3.12 Log CFU/m<sup>3</sup>) were slightly higher than in farms (2.95 Log CFU/m<sup>3</sup>). However, maximum counts presented in Farms were greater, with 4.73 Log CFU/m<sup>3</sup>. The main fungi found throughout the samples were from the genera *Aspergillus* sp., *Fusarium* sp., *Penicillium* sp., *Cladosporium* sp., as well as yeasts.

The high frequency of *Cladosporium* sp. found in the present study was also observed by Despot and Klarić [20] in air samples from grain mills in Croatia. In a study on occupational exposure in Irish bakeries and a pizza restaurant, the most common genera identified were similar to those found in the present study: personal air samples showed *Penicillium* sp. (91.1%), *Aspergillus* sp. (4.8%), and *Cladosporium* sp. (4.1%), and settled dust samples showed *Penicillium* sp. (88.3%), *Aspergillus* sp. (7.8%), and *Chrysosporium* sp. (2.9%) [17].

The highest frequency of the genus *Aspergillus* in farms and feed factories observed in our study is agreement with the findings by Viegas, *et al.* [21], who also most frequently identified the genus *Aspergillus* in pig farms (40.5%), at high concentrations (> 2,000 CFU/m<sup>3</sup>). A study of respiratory filters of protection devices among

Type of fungus	Percentage of positive samples			
	Feed factories (N = 3)		Farms (N = 4)	
	Respirable dust samples (N = 12)	Surface dust samples (N = 24)	Respirable dust samples (N = 14)	Surface dust samples (N = 28)
<i>Aspergillus</i> sp.	67	67	64	79
<i>Fusarium</i> sp.	50	54	36	79
<i>Cladosporium</i> sp.	58	75	64	82
<i>Penicillium</i> sp.	16	29	29	39
<i>Rhizopus</i> sp.	16	38	7	21
<i>Trichothecium</i> sp.	0	4	0	7
<i>Trichoderma</i> sp.	0	0	0	7
<i>Mucor</i> sp.	0	0	0	4
NSF	67	38	28	21
Yeasts	16	71	21	64
Average count (range), Log <sub>10</sub> CFU/m <sup>3</sup>	3.12 (2.00 - 3.98)	5.99 (4.04 - 7.10) <sup>1</sup>	2.95 (2.30 - 4.73)	5.26 (3.30 - 7.04) <sup>1</sup>

**Table 1:** Main fungi detected in respirable dust and surface dust samples in four samplings conducted in feed factories and farms in the state of São Paulo, Brazil.

<sup>1</sup>Log<sub>10</sub> CFU/m<sup>2</sup>.

N: Number of Total Samples Collected; CFU: Colony-Forming Units; NSF: Non-Sporulating Fungus.

workers from a waste sorting industry detected *Aspergillus* sp. most frequently, with maximum counts of 208,500 CFU/m<sup>2</sup> (5.32 LogCFU/m<sup>2</sup>) [22]. Air samples from poultry farms in Portugal showed 40.5% *Scopulariopsis brevicaulis*, 30% *Rhizopus* sp., 10.1% *Penicillium* sp., and 9.7% *Aspergillus* sp. and samples from surfaces in the same places showed 47.9% *Aspergillus* sp., 15.3% *Fusarium* sp., and 8.8% *Penicillium* sp. [23].

High frequency of *Aspergillus* sp. is a health concern for the workers, since this genus is the most common producer of aflatoxins [10], which is among the most toxic mycotoxins for humans due to its genotoxic and carcinogenic potential [24]. In addition, species from the genus *Aspergillus* can produce ochratoxin A and fumonisins, both classified as possible carcinogens to humans [24]. Furthermore, *Fusarium* sp. and *Penicillium* sp. are also important producers of mycotoxins, such as fumonisins, ochratoxin A, zearalenone and deoxynivalenol, among other mycotoxins [8,11-13].

Fungi counts reported in the present work were well below those reported in environmental samples collected in places that

do not handle grains or animal feed, including Irish bakeries [17], a veterinary practice in Krakow [25], a waste-sorting plant in Poland [26], a flour mill in Egypt [27], and apartments and basements in Croatia [20]. However, in air samples from grain mills in Croatia, fungi counts were close to those found in the present study, ranging from 1,700 to 40,000 CFU/m<sup>3</sup> (3.23 to 4.6 Log CFU/m<sup>3</sup>) [20]. Counts above 3.30 Log CFU/m<sup>3</sup> or m<sup>2</sup> were also identified in pig farms and waste sorting industries [21,22]. These findings, taken together, confirm the presence of large amounts of fungi in environments where grains and animal feed are handled in Brazil, raising concerns about the risk posed to the health of workers.

## Conclusion

The present study shows a high presence of fungi producers of mycotoxins on farms and feed factories in São Paulo, Brazil. However, the presence of fungi does not necessarily mean the presence of mycotoxins in the environment, thus this is exactly the reason further studies are necessary. Further studies should evaluate the levels of mycotoxins produced by the commonly occurring species from the fungi genera detected in the environments of the studied

workplaces. Furthermore, in these workplaces, the use of protective equipment should be mandatory to avoid possible contamination and inflammatory reactions in the respiratory system.

### Acknowledgements

This research was funded by the São Paulo Research of Foundation (FAPESP), grant #2019/00990-7, for financial support and fellowship.

### Competing interests

The authors declare that there is no conflict of interest.

### Bibliography

1. Karjalainen A., *et al.* "Incidence of occupational asthma by occupation and industry in Finland". *American Journal of Industrial Medicine* 37 (2000): 451-458.
2. Burch JB., *et al.* "Endotoxin exposure and inflammation markers among agricultural workers in Colorado and Nebraska". *Journal of Toxicology and Environmental Health Part A* 73 (2010): 5-22.
3. Viegas C., *et al.* "Fungal contamination in swine: A potential occupational health threat". *Journal of Toxicology and Environmental Health Part A* 76.4-5 (2013): 272-280.
4. Viegas C., *et al.* "Environmental impact caused by fungal and particles contamination of Portuguese swine". *Transactions on Biomedicine and Health* 16 (2013): 11-23.
5. Cleave J., *et al.* "Fractionation of swine barn dust and assessment of its impact on the respiratory tract following repeated airway exposure". *Journal of Toxicology and Environmental Health Part A* 73 (2010): 1090-1101.
6. Harting JR., *et al.* "Chronic obstructive pulmonary disease patients have greater systemic responsiveness to *ex vivo* stimulation with swine dust extract and its components versus healthy volunteers". *Journal of Toxicology and Environmental Health Part A* 75 (2012): 1456-1470.
7. Cote J *et al.* "Occupational asthma caused by exposure to neurospora in plywood factory worker". *American Journal of Industrial Medicine* 48 (1991): 279-282.
8. Richard JL. "Some major mycotoxins and their mycotoxicoses -an overview". *International Journal of Food Microbiology* 119.1-2 (2007): 3-10.
9. Degen GH. "Tools for investigating workplace-related risks from mycotoxin exposure". *World Mycotoxin Journal* 4 (2011): 315-327.
10. Oliveira CAF and Corrêa B. "Interactive effects between mycotoxins in livestock". In: *Mycotoxins in animals economically important*. New York: Nova Science Publishers (2010): 117-129.
11. Frisvad JC., *et al.* "Fumonisin and ochratoxin production in industrial *Aspergillus niger* strains". *Plos One* 6.8 (2011): e23496.
12. Ostry V., *et al.* "Mycotoxins as human carcinogens-the IARC Monographs classification". *Mycotoxin Research* 33.1 (2017): 65-73.
13. Kőszegi T and Poór M. "Ochratoxin A: Molecular interactions, mechanisms of toxicity and prevention at the molecular level". *Toxins* 8.4 (2016): 111-125.
14. Franco LT and Oliveira CAF. "Assessment of occupational and dietary exposures of feed handling workers to mycotoxins in rural areas from São Paulo, Brazil". *Science of the Total Environment* 837 (2022): 155763.
15. Mohamad Asri AA., *et al.* "Exposure of aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) in inhalable dust and its respiratory effects among rice millers". *World Journal of Translational Medicine* 23 (2020): 60-71.
16. Ndaw S., *et al.* "Investigating multi-mycotoxin exposure in occupational settings: a biomonitoring and airborne measurement approach". *Toxins* 13.1 (2021): 429.
17. Viegas C., *et al.* "Occupational exposures to organic dust in Irish bakeries and a pizzeria restaurant". *Microorganisms* 8.1 (2020): 118.
18. Vincent JH and Mark D. "The basis of dust sampling in occupational hygiene: a critical review". *The Annals of Occupational Hygiene* 24.4 (1981): 375-390.

19. De Hoog GS., *et al.* "Atlas of clinical fungi. 2<sup>nd</sup> edition". Centraalbureau voor Schimmelcultures, Utrecht, the Netherlands (2000).
20. Despot DJ and Klarić MS. "A year-round investigation of indoor airborne fungi in Croatia". *Archives of Industrial Hygiene and Toxicology* 65 (2014): 209-218.
21. Viegas S., *et al.* "Occupational exposure to aflatoxin B<sub>1</sub> in swine production and possible contamination sources". *Journal of Toxicology and Environmental Health Part A* 76.15 (2013): 944-951.
22. Viegas C., *et al.* "Are workers from waste sorting industry really protected by wearing filtering respiratory protective devices? The gap between the myth and reality". *Waste Management* 102 (2020): 856-867.
23. Viegas C., *et al.* "Assessing indoor fungal contamination using conventional and molecular methods in Portuguese poultries". *Environmental Monitoring and Assessment* 186.3 (2014): 1951-1959.
24. IARC. International Agency for Research on Cancer. "IARC Monograph on the Evaluation of Carcinogenic Risk to Humans" 82.171 (2022).
25. Bulski K and Frączek K. "Mycological air quality at animal veterinary practice". *Rocznik Ochrona Srodowiska* 23 (2021): 168-179.
26. Bragoszewska E. "The dose of fungal aerosol inhaled by workers in a waste-sorting plant in Poland: A case study". *International Journal of Environmental Research and Public Health* 17 (2019): 177.
27. Saad-Hussein A., *et al.* "Effects of airborne *Aspergillus* on serum aflatoxin B<sub>1</sub> and liver enzymes in workers handling wheat flour". *Human and Experimental Toxicology* 35.1 (2016): 3-9.