



## Antioxidant Properties of Nine Commonly Used Medicinal Plants in Arabs Region

**Abuelgassim Omer Abuelgassim\***

Department of Biochemistry, King Saud University, Riyadh, Kingdom of Saudi Arabia

\*Corresponding Author: Abuelgassim Omer Abuelgassim, Department of Biochemistry, King Saud University, Riyadh, Kingdom of Saudi Arabia.

DOI: 10.31080/ASNH.2022.06.1149

Received: October 25, 2022

Published: November 07, 2022

© All rights are reserved by **Abuelgassim Omer Abuelgassim**.

### Abstract

The present study was conducted to investigate the total phenolics, total flavonoid, and Trolox equivalent antioxidant capacity of 9 commonly used medicinal plants and herbs among Arabs nation namely, *Trachyspermum ammi* (*Apiaceae*) seeds, *Ferula assa-foetida* (*Apiaceae*) dried latex, *Elettaria cardamomum* (*Zingiberaceae*) fruit 1, *Senna alexandrina* (*Caesalpinioideae*) *Leguminosae* leaves, *Adenium obesum* (*Apocynaceae*) leaves, *Elettaria cardamomum* (*Zingiberaceae*) fruit 2, *Trigonella foenum-graecum* (*Leguminosae*) seeds, *Lepidium sativum* (*Brassicaceae*) seeds, *Ziziphus spina-christi* (*Rhamnaceae*) fruit, and *Nigella sativa* (*Ranunculaceae*) seeds.

The methanol extract of each plant was obtained by dissolving 2g of plant powder in 20 ml methanol-water (4:1v/v) to overnight at room temperature. The supernatants were concentrated by using a rotary evaporator. Residues of each extract were dissolved in distilled water and the final volume was recorded. A serial dilution of 10, 100, 1000, and 5000 were made.

The mean values of the total phenolics content were  $535.24 \pm 70.68$ ,  $150.28 \pm 36.89$ ,  $69.05 \pm 14.18$ ,  $1358.7 \pm 118.88$ ,  $605.83 \pm 117.67$ ,  $228.28 \pm 53.8$ ,  $2171.57 \pm 296.2$ ,  $726.65 \pm 130.73$ ,  $702.63 \pm 32.93$ , and  $132.4 \pm 30.77$  (mg GAE/100g) in *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum-graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* respectively, whereas the mean values of the total flavonoid content were  $143.96 \pm 6.44$ ,  $0.0$ ,  $9.16 \pm 3.14$ ,  $761.66 \pm 14.71$ ,  $210.09 \pm 7.61$ ,  $12.83 \pm 0.88$ ,  $347 \pm 60.15$ ,  $94.84 \pm 10.04$ ,  $94.07 \pm 6.26$ , and  $30.49 \pm 0.33$  (mg QE/100g) respectively. The determined scavenging activity against ABTS radical of the aqueous methanolic extracts of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum-graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* were  $1271.32 \pm 273.35$ ,  $234.89 \pm 42.2$ ,  $135 \pm 21.45$ ,  $2551.83 \pm 305.35$ ,  $2101.1 \pm 298.97$ ,  $794.9 \pm 175.72$ ,  $794.73 \pm 149.49$ ,  $2219.74 \pm 382.12$ ,  $3093.11 \pm 445.93$ , and  $483.07 \pm 98.17$  ( $\mu$ mole TEAC/100g) respectively. Among studied plants *T. foenum-graecum* showed the highest phenolic content ( $2171.57 \pm 296.2$  mg GAE/100g) whereas *S. alexandrina* showed the highest total flavonoid content ( $761.66 \pm 14.71$  mg QE/100g). The highest  $IC_{50s}$  were noticed in *T. ammi*, *S. alexandrina*, *Z. spina-christi*, and *A. obesum* extracts ( $100.035 \pm 19.38$ ,  $583.38 \pm 45.11$ ,  $666.87 \pm 79.62$  and  $720.51 \pm 92.24$   $\mu$ g respectively). The hydromethanolic extracts of *T. foenum-graecum*, *S. alexandrina*, *A. obesum*, *L. sativum* and *Z. spina-christi* showed a significant antioxidant property, further research is ongoing to identify their bioactive compounds, in particular phenolic acids. Our findings suggest the possibility of applying *T. foenum-graecum*, *S. alexandrina*, *A. obesum*, *L. sativum* and *Z. spina-christi* in food and/or pharmaceutical industries.

**Keywords:** Total Phenolics; Total Flavonoid; Scavenging Activity; *T. Foenum-Graecum*; *S. Alexandrina*; *Z. Spina-Christi*

## Introduction

In developing countries, herbal remedies are used for treatment of human diseases such as atherosclerosis, cancer, diabetes, aging, various skin diseases and other degenerative diseases [1-5]. In recent years an increasing attention has been focused on the study of substances with antioxidant potential and protective effects against free radicals, and reactive oxygen and nitrogen species. Free radicals, and reactive oxygen and nitrogen species can cause oxidative stress resulting in damage to several biomolecules, and therefore leads to chronic disease and serious illness [6]. Plants, including vegetables, fruits, herbs, and cereals, have various bioactive compounds that have anti-diabetic, antibacterial, anti-cancer, anti-atherogenic, and anti-inflammatory effects [7]. The administration of *Senna alata* (*S. alata*) extract reported to improve the impaired lipid metabolism in HFD-induced obese mice by the reducing of serum lipid concentrations and hepatic triglycerides content [8]. Mercury chloride - induced nephrotoxicity rats when administered *Ziziphus spina-christi* leaf extract at a concentration of 300mg/Kg showed significant reduction in serum creatinine and urea, also kidney lipid peroxidation was significantly reduced as malondialdehyde levels diminished [9]. *A. obesum* ethanolic seeds extract reported to have a cytotoxic potential against MCF-7 breast cancer cells [10]. Serum levels of glucose, lipid profile, kidney, and liver enzyme were significantly reduced after treatment of streptozotocin-induced diabetic rats with oral administration of *L. sativum* extract at the dosage of 100 and 200 mg/kg [11]. Supplementation of *T. foenum-graecum* leaves in the diet to streptozotocin-induced diabetic rats daily for 45 days reduced serum and tissue concentration of total cholesterol, triglycerides, and free fatty acids [12]. *T. foenum-graecum* leaf extract also showed a significant level of total phenolics and antioxidant properties and significantly reduced the lipid oxidation in chicken meat during refrigerated storage [13].

Here, we studied the total phenolics, total flavonoid, and ABTS radical scavenging activity of *T. ammi*, *F. assa-foetida*, *E. cardamomum*1, *S. alexandrina*, *A. obesum*, *E. cardamomum*2, *T. foenum-graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa*, a commonly used medicinal plants in Arabs region. The possibility of applying extracts with potential antioxidant properties in food and/or pharmaceutical industries was also a focus.

## Materials and Methods

### Chemicals and reagents

Methanol and other chemicals were of analytical grade and were obtained from BDH (Poole, UK). Gallic acid, Folin-Ciocalteu reagent, quercetin, 2,2-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) di-

ammonium salt (ABTS), and potassium persulfate were purchased from Sigma-Aldrich (St. Louis, MO, USA). Trolox (6-hydroxy-2, 5, 7, 8-tetramethylchroman-2-carboxylic acid) was obtained from Fluka Chemie AG (Buchs, Switzerland).

### Plant materials and extraction

*Trachyspermum ammi* (*Apiaceae*) seeds, *Ferula assa-foetida* (*Apiaceae*) dried latex, *Elettaria cardamomum* (*Zingiberaceae*) Fruit 1, *Senna alexandrina* (*Caesalpinioideae*) *Leguminosae* leaves, *Adenium obesum* (*Apocynaceae*) leaves, *Elettaria cardamomum* (*Zingiberaceae*) Fruit 2, *Trigonella foenum-graecum* (*Leguminosae*) seeds, *Lepidium sativum* (*Brassicaceae*) seeds *Ziziphus spina-christi* (*Rhamnaceae*) Fruit, and *Nigella sativa* (*Ranunculaceae*) seeds. All plants were obtained from the local market, Jeddah, the western region of Kingdom of Saudi Arabia except *Adenium obesum* was obtained from Yemen.

Plants were identified by staff members of the Botany Department, College of Science, King Saud University, Saudi Arabia. Voucher specimens were deposited at the herbarium of the Botany Department. Powdered extracts were obtained by use of a mill and they were stored in amber bottles to prevent degradation. The methanol extract of each plant was obtained by dissolving 2g of each plant powder in 20 ml methanol-water (4:1v/v) to overnight at room temperature using an orbital shaker. Filtrates were collected and residues were extracted again with 20 ml solvent. Appropriate filtrates were pooled and centrifuged at 5000 rpm for 10 min, and the supernatants were concentrated under reduced pressure at 40°C using a rotary evaporator. After solvent evaporation, residues of each extract were dissolved in distilled water and the final volume was recorded. A serial dilutions of 10, 100, 1000, and 5000 were made for each plant extract and stored at -20 °C until analysis.

### Determination of total phenolics (TP)

Total phenolics content was determined by Folin-Ciocalteu's reagent method [14]. 0.5 ml of each diluted extract and 0.1 ml Folin-Ciocalteu's reagent were mixed and incubated at room temperature for 15 min. Then 2.5 ml of 7% sodium carbonate solution was added, the tubes were incubated for 30 min at room temperature and the absorbance was measured at 760 nm. Gallic acid standard curve was constructed (0 -50 µg). Total phenolics content of plant extracts were expressed as mg gallic acid equivalent (GAE)/100 g.

### Determination of total flavonoid (TF)

Total flavonoid content was determined by aluminum chloride colorimetric method [15]. In brief, 0.5ml of each diluted extract

was mixed with 1.5 ml of 95% ethanol, 0.5 ml of 1.2% aluminum chloride, 0.5 ml of 120 mM potassium acetate were added in order. After incubation at room temperature for 30 min, the absorbance was measured at 415 nm against reagent blank. Quercetin standard curve was constructed (0- 50  $\mu\text{g}$ ). Total flavonoid content of plant extracts was expressed as milligram quercetin antioxidant equivalents (QE/100g).

### ABTS radical cation scavenging activity

The ABTS free radical decolorization assay was performed [16]. ABTS radical cation was generated by reacting 7 mM ABTS and 2.45 mM potassium persulfate after incubation in dark for 16hr at room temperature. The ABTS solution was diluted with 80% ethanol to an absorbance of  $0.700 \pm 0.050$  at 734 nm. 0.1 mL of each diluted extract and 3.9 ml ABTS solution were mixed thoroughly, the mixture was allowed to stand at room temperature for 6 min and the absorbance was immediately recorded at 734 nm. Trolox standard curve (0-15  $\mu\text{M}$ ) in 80% ethanol was prepared. Results were expressed in terms of Trolox equivalent antioxidant capacity (TEAC,  $\mu\text{mol}$  Trolox equivalents per 100g sample).

### Data analysis

Results were presented as means  $\pm$  standard deviation. All samples were analyzed in three replications.

## Results and Discussion

The calibration curve of gallic acid for determining total phenolics content is shown in figure 1. The total phenolics content of the aqueous methanolic extracts of *T. ammi*, *F. assa- foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum- graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* were  $535.24 \pm 70.68$ ,  $150.28 \pm 36.89$ ,  $69.05 \pm 14.18$ ,  $1358.7 \pm 118.88$ ,  $605.83 \pm 117.67$ ,  $228.28 \pm 53.8$ ,  $2171.57 \pm 296.2$ ,  $726.65 \pm 130.73$ ,  $702.63 \pm 32.93$ , and  $132.4 \pm 30.77$  (mg GAE/100g) respectively (Figure 2). *T. foenum- graecum* showed the highest phenolics content ( $2171.57 \pm 296.2$ mg GAE/100g). A high phenolics content of the freeze-dried *Trigonella foenum-graecum* seeds was reported [17]. Another study also reported that fenugreek seeds have a high level of total polyphenol content (85.88 mg GAE/g) [18]. *S. alexandrina* also showed a considerable total phenolics content ( $1358.70 \pm 118.88$  mg GAE/100g). The association between the presence of phenolic compounds and the biological activity of the acetone extract of *Senna italica* has been reported [19]. *T. ammi* showed a total phenolics content of ( $535.24 \pm 70.68$  mg GAE/100 g), however,




Figure 1: Standard curve of Gallic acid.




Figure 2: Total phenolics content as mg gallic acid (GAE) /100g of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum-graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa*. Data are expressed as means  $\pm$  standard deviation and calculated from three determinations.

an early study suggested that phenolic compounds are important components of *Trachyspermum ammi*, and the pharmacological effects of the plant could be attributed to the presence of these polyphenols [20]. Our finding regarding the total phenolics content of *A. obesum* ( $605.83 \pm 117.67$  mg GAE/100g) is coincided with the report concluded that *A. obesum* extract collected from Yemen is a rich source of anthocyanins and possesses a significant antioxidant activity [21]. The total phenolics content of *L. sativum* seeds and *Z. spina-christi* flower was  $726.65 \pm 130.73$  and  $702.63 \pm 32.93$  mg GAE/100g respectively.

The total phenolics content of *L. sativum* is in a full agreement with the reported concentration range between 6.33 - 7.40 mg GAE/g [22]. A total phenolics content of  $1644.00 \pm 3.20$  mg GAE/100g in *Z. spina-christi* flower has been reported [23], however, the discrepancy between the current investigation and that report, concerning the total phenolics content in *Z. spina-christi* flower could be attributed to the source of the plant and the extraction protocol. The observed total phenolics content in *E. cardamomum1* and *E. cardamomum2* were  $69.05 \pm 14.18$  and  $228.28 \pm 53.8$  mg GAE/100g respectively. The total phenolics components of the *E. cardamomum* at different methanol concentrations has been studied and found to range from 0.317 to 1.66 g GAE/100g [24]. Our finding regarding the total phenolics content in *F. assa-foetida* is in contrast to an earlier reported data of the plant, this contrast could be attributed to extraction protocol [25]. Our obtained result regarding the total phenolics content of *N. sativa*, ( $132.40 \pm 30.77$  mg GAE/100g) is consistent with the report stated that the phenol content of *N. sativa* is  $122.67 \pm 3.03$  mg GAE/100g [26].

The calibration curve of quercetin for determining total flavonoid content is shown in figure 3. The total flavonoid content of the aqueous methanolic extracts of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum- graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* were  $143.96 \pm 6.44, 0.0, 9.16 \pm 3.14, 761.66 \pm 14.71, 210.09 \pm 7.61, 12.83 \pm 0.88, 347 \pm 60.15, 94.84 \pm 10.04, 94.07 \pm 6.26,$  and  $30.49 \pm 0.33$  (mg QE/100g) respectively (Figure 4). The highest content of total flavonoid content was observed in *S. alexandrina* leaves ( $761.66 \pm 14.71$  mg QE/100g). The total flavonoid content in the bark of *Senna tora* which belongs to the family *Caesalpinaceae*, another species of *Senna*, was reported to be 21.58 mg QE/100g [27]. Another study showed that the hydroalcoholic extract of *Cassia acutifolia*, another *Senna* species, has a total flavonoid content of  $20.8 \pm 0.40$  mg QE/g [28].

*T. foenum- graecum* seeds content of total flavonoid was  $347 \pm 60.15$ mg QE/100g (Figure 4). However, the aqueous methanolic extract of the plant seed reported to have a total flavonoid content (as Catechin) of 20.8 mg CE/100g [29]. The total flavonoid content in *A. obesum* and *T. ammi* were  $210.09 \pm 7.61$  and  $143.96 \pm 6.44$  mg QE/100g respectively. Flavonoids have been reported to have antibacterial, antineoplastic activity, anti-inflammatory, anti-allergic and antiviral activity. Our applied extraction protocol as well as the protocol for total flavonoid content determination failed to detect flavonoid in the dried latex of *F. assa-foetida*, however, it has been




Figure 3: Standard curve of Quercetin.




Figure 4: Total flavonoid content as mg Quercetin (QEA)/100g of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum-graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa*. Data are expressed as means  $\pm$  standard deviation and calculated from three determinations.

reported that the hydroalcoholic extract of *F. assa-foetida* leaves and *F. assa-foetida* leaves essential oil have a total flavonoid content of  $12.53 \pm 3.20, 0.015 \pm 0.002$  mg QE/100g respectively [30]. *L. sativum* and *Z. spina-christi* have the same total flavonoid content (94 mg QE/100g) (Figure 4). Our finding regarding the total flavonoid content of *Z. spina-christi* is comparable with the report stated that the fruit of *Z. spina-christi* grown in Oman has a total flavonoid content of 47 mg CE/100 g [23].

The calibration curve of Trolox scavenging activity against ABTS radical is shown in figure 5. The recorded scavenging activity of the aqueous methanolic extracts of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T.*




Figure 5: Standard curve of Trolox.

*foenum- graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* were  $1271.32 \pm 273.35$ ,  $234.89 \pm 42.2$ ,  $135 \pm 21.45$ ,  $2551.83 \pm 305.35$ ,  $2101.1 \pm 298.97$ ,  $794.9 \pm 175.72$ ,  $794.73 \pm 149.49$ ,  $2219.74 \pm 382.12$ ,  $3093.11 \pm 445.93$ , and  $483.07 \pm 98.17$  ( $\mu\text{mole TEAC}/100\text{g}$ ) respectively (Figure 6).




Figure 6: Trolox equivalent antioxidant capacity (TEAC) as  $\mu\text{mole}/100\text{g}$  of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum- graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa*.

Data are expressed as means  $\pm$  standard deviation and calculated from three determinations.

The obtained  $\text{IC}_{50}$  of Trolox standard against ABTS radical was determined to be  $11.63 \mu\text{M}$  (Figure 5), whereas the calculated  $\text{IC}_{50}$ s of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum- graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* were  $100.035 \pm 19.38$ ,  $3471.5 \pm 213.76$ ,

$1015.52 \pm 166.04$ ,  $583.38 \pm 45.11$ ,  $720.51 \pm 92.24$ ,  $1569.4 \pm 217.94$ ,  $1558.96 \pm 206.24$ ,  $817.01 \pm 81.21$ ,  $666.87 \pm 79.62$ , and  $2673.52 \pm 241.59 \mu\text{g}$  respectively (Figure 7). The highest  $\text{IC}_{50}$ s were noticed in *T. ammi*, *S. alexandrina*, *Z. spina-christi*, and *A. obesum* extracts ( $100.035 \pm 19.38$ ,  $583.38 \pm 45.11$ ,  $666.87 \pm 79.62$  and  $720.51 \pm 92.24 \mu\text{g}$  respectively). A recent report showed that the aerial part of *S. alexandrina* that undergo ultrasound-assisted extraction exhibited a powerful antioxidant activity against DPPH and ABTS when ascorbic acid is used as standard [31]. *Alternaria alternata* that had been isolated from *Ziziphus spina-christi* reported to have as potential antioxidant activity against DPPH as the  $\text{IC}_{50}$  reached  $409 \mu\text{g}/\text{ml}$  [32]. Our finding that *T. ammi* seed extract has a strong ABTS scavenging activity as the  $\text{IC}_{50}$  reached  $100.035 \pm 19.38 \mu\text{g}$  is in full agreement with the report that studied 24 commonly used medicinal plants from Nepal and Japan and stated that *T. ammi* seed showed a strong activity against DPPH as the recorded % of inhibition was  $90.67 \pm 0.58$  [33].




Figure 7: Micrograms required of *T. ammi*, *F. assa-foetida*, *E. cardamomum1*, *S. alexandrina*, *A. obesum*, *E. cardamomum2*, *T. foenum- graecum*, *L. sativum*, *Z. spina-christi* and *N. sativa* extracts to give  $\text{IC}_{50}$ s against ABTS radical.

Data are expressed as means  $\pm$  standard deviation and calculated from three determinations.

### Conclusion

The hydromethanolic extracts of *T. foenum-graecum*, *S. alexandrina*, *A. obesum*, *L. sativum* and *Z. spina-christi* showed a significant antioxidant property, further research is ongoing to identify their bioactive compounds, in particular phenolic acids. Our findings suggest the possibility of applying *T. foenum-graecum*, *S. alexandrina*, *A. obesum*, *L. sativum*, and *Z. spina-christi* in food and/or pharmaceutical industries.



## Acknowledgements

The author would like to thank the research center, college of science for supporting this project. The author also thanks the Deanship of Scientific Research and RSSU at King Saud University for their technical support.

## Conflict of Interest

The author declares no conflict of interest.

## Bibliography

1. Sabu MC and Kuttan R. "Anti-diabetic activity of medicinal plants and its relationship with their antioxidant property". *Journal of Ethnopharmacology* 81 (2002): 155-160.
2. Naik GH., et al. "Comparative antioxidant activity of individual herbal components used in Ayurvedic medicine". *Phytochemistry* 63 (2003): 97-104.
3. Cai Y., et al. "Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anti-cancer". *Life Sciences* 74 (2004): 2157-2184.
4. Al-Mustafa AH and Al-Thunibat OY. "Antioxidant activity of some Jordanian medicinal plants used traditionally for treatment of diabetes". *Pakistan Journal of Biological Sciences* 11 (2008): 351-358.
5. Singh S., et al. "A Review on Cassia species: Pharmacological, Traditional and Medicinal Aspects in Various Countries". *American Journal of Phytomedicine and Clinical Therapeutics* 1.3 (2013): 291- 312.
6. Dalle-Donne Isabella., et al. "Biomarkers of oxidative damage in human disease". *Clinical Chemistry* 52.4 (2006): 601-623.
7. Saldana M., et al. In "Emerging Technologies Used for the Extraction of Phytochemicals from Fruits, Vegetables, and Other Natural Sources". de la Rosa, L.A., Alvarez-Parrilla, E., Gonzalez-Aguilar, G.A., Eds., Blackwell Publishing, John Wiley and Sons, USA (2010): 246.
8. Naowaboot J and Wannasiri S. "Anti-lipogenic effect of Senna alata leaf extract in high-fat diet-induced obese mice". *Asian Pacific Journal of Tropical Biomedicine* 6.3 (2016): 232-238.
9. Almeer Rafa S., et al. "Ziziphus spina-christi Leaf Extract Suppressed Mercury Chloride-Induced Nephrotoxicity via Nrf2-Antioxidant Pathway Activation and Inhibition of Inflammatory and Apoptotic Signaling". *Oxidative Medicine and Cellular Longevity* 2019 (2019): 5634685.
10. Ali Ahmed Qasem., et al. "Cytogenotoxic effects of Adenium obesum seeds extracts on breast cancer cells". *Saudi Journal of Biological Sciences* 26.3 (2019): 547-553.
11. Ramadan Shimaa., et al. "Comparison of the Efficiency of Lepidium sativum, Ficus carica, and Punica granatum Methanolic Extracts in Relieving Hyperglycemia and Hyperlipidemia of Streptozotocin-Induced Diabetic Rats". *Journal of Diabetes Research* 2021 (2021): 6018835.
12. Annida B and P Stanely Mainzen Prince. "Supplementation of fenugreek leaves lower lipid profile in streptozotocin-induced diabetic rats". *Journal of Medicinal Food* 7.2 (2004): 153-156.
13. Devatkal Suresh K., et al. "Comparative antioxidant effect of aqueous extracts of curry leaves, fenugreek leaves and butylated hydroxytoluene in raw chicken patties". *Journal of Food Science and Technology* 49.6 (2012): 781-785.
14. McDonald Suzy., et al. "Phenolic content and antioxidant activity of olive extracts". *Food Chemistry* 73. (2001): 73-84.
15. Chang C., et al. "Estimation of total flavonoid content in propolis by two complementary colorimetric methods". *Journal of Food and Drug Analysis* 10. (2002): 178-182.
16. Re R., et al. "Antioxidant activity applying an improved ABTS radical cation decolorization assay". *Free Radical Biology and Medicine* 26.9-10 (1999): 1231-1237.
17. Kenny O., et al. "Antioxidant properties and quantitative UPLC-MS analysis of phenolic compounds from extracts of fenugreek (*Trigonella foenum-graecum*) seeds and bitter melon (*Momordica charantia*) fruit". *Food chemistry* 141.4 (2013): 4295-4302.
18. Naidu MM., et al. "Chemical composition and antioxidant activity of the husk and endosperm of fenugreek seeds". *LWT - Food Science and Technology* 44 (2011): 451- 456.
19. Masoko P., et al. "Evaluation of the antioxidant, antibacterial, and antiproliferative activities of the acetone extract of the roots of Senna italica (Fabaceae)". *African Journal of Traditional, Complementary, and Alternative Medicines : AJTCAM* 7.2 (2009): 138-148.
20. Chatterjee S., et al. "Estimation of Phenolic Components and in vitro Antioxidant Activity of Fennel (*Foeniculum vulgare*) and Ajwain (*Trachyspermum ammi*) seeds". *Advances in Bioresearch* 3.2 (2012): 109-118.

21. Ebrahim N., *et al.* "Free radical scavenging activity and anthocyanin in flower of *Adenium obesum* collected from Yemen". *Journal of Pharmacy and Phytotherapeutics* 1 (2013): 5-7.
22. Sat IG., *et al.* "Antioxidant and nutritional characteristics of garden cress (*Lepidium sativum*)". *Acta Scientiarum Polonorum Hortorum Cultus* 12.4 (2013): 173-179.
23. Singh V., *et al.* "In vitro Antioxidant Activities of *Ziziphus spina-christi* Fruits (Red Date) Grown in Oman". *Biotechnology* 11.4 (2012): 209-216.
24. Bhatti HN., *et al.* "Evaluation of Phenolic Contents and Antioxidant Potential of Methanolic Extracts of Green Cardamom (*Elettaria cardamomum*)". *Asian Journal of Chemistry* 22.6 (2010): 4787- 4794.
25. Ebrahimzadeh MA., *et al.* "Correlation between the *in vitro* iron chelating activity and poly phenol and flavonoid contents of some medicinal plants". *Pakistan Journal of Biological Sciences* 12.12 (2009): 934-938.
26. Souri E., *et al.* "Screening of antioxidant activity and phenolic content of 24 medicinal plant extracts". *DARU Journal of Pharmaceutical Sciences* 16.2 (2008 ): 83-87.
27. Sulaiman CT and Balachandran I. "Total Phenolics and Total Flavonoids in Selected Indian Medicinal Plants". *Indian Journal of Pharmaceutical Sciences* 74.3 (2012): 258-260.
28. Al-Dabbagh Bayan., *et al.* "Antioxidant and anticancer activities of *Trigonella foenum-graecum*, *Cassia acutifolia* and *Rhazya stricta*". *BMC Complementary and Alternative Medicine* 18.1 (2018): 240.V
29. Belguith-Hadriche O., *et al.* "Comparative study on hypocholesterolemic and antioxidant activities of various extracts of fenugreek seeds". *Food Chemistry* 138 (2013): 1448-1453.
30. Ahmadvand H., *et al.* "Chemical Composition and Antioxidant Properties of *Ferula-assa-foetida* Leaves Essential Oil". *Iranian Journal of Pharmacology and Therapeutics* 12 .2 (2013): 52-57.
31. Alam Perwez., *et al.* "Response Surface Methodology (RSM)-Based Optimization of Ultrasound-Assisted Extraction of Sennoside A, Sennoside B, Aloe-Emodin, Emodin, and Chrysophanol from *Senna alexandrina* (Aerial Parts): HPLC-UV and Antioxidant Analysis". *Molecules (Basel, Switzerland)* 27.1 (2022): 298.
32. Elghaffar Rasha Y Abd., *et al.* "Promising Endophytic *Alternaria alternata* from Leaves of *Ziziphus spina-christi*: Phytochemical Analyses, Antimicrobial and Antioxidant Activities". *Applied Biochemistry and Biotechnology* 194.9 (2022): 3984-4001.
33. Ghimir BK., *et al.* "A comparative evaluation of the antioxidant activity of some medicinal plants popularly used in Nepal". *Journal of Medicinal Plants Research* 5.10 (2011): 1884-1891.