



## Long Covid (SARS-CoV-2/Covid-19) Infection Related Self-inflicted oral Trauma to Right-Side of Tongue Due to Jaw-Cheek-Tongue Dynamic Motor Incoordination with Recurrent Right-Sided Anterior Bleeding Tongue-Bite Crush Lesions and Right-Sided Lower-Lip Bruising Only During Chewing Food

**Vinod Kumar Gupta\***

General Medicine, Medical Director-Physician, Migraine-Headache Institute Gupta Medical Centre, India

\***Corresponding Author:** Vinod Kumar Gupta, General Medicine, Medical Director-Physician, Migraine-Headache Institute Gupta Medical Centre, India.

**Received:** September 03, 2024

**Published:** October 14, 2024

© All rights are reserved by **Vinod Kumar Gupta**.

**Keywords:** Fifth Cranial Nerve; Mandibular Division Motor Fibres; Facial Nerve; Hypoglossal Nerve; Bleeding Tongue-Bite; Lip-Bruising; Chewing; MRI

I report a prolonged case of SARS-CoV-2 (Covid-19) infection, now known as Long Covid, in a 67 years old man. This patient had recurrent tongue bites during eating (four episodes), particularly with stacking of food, leading to bleeding crush-lesions with ulcers that would spontaneously heal in 4-6 weeks. He has also bruised his lower lip frequently on the same side. His family members have noted recurrent but transient (30-60 seconds) deviation of angle of mouth to the left, indicating intermittent short-lasting right-sided facial nerve paresis with subclinical weakening of cheek muscles of right side of the face. He has now to take care while chewing food, to eat with greater care and more slowly, and restrain himself from talking during meals. All cranial nerves are otherwise normal on clinical examination as well as on MRI. Such self-inflicted events did not occur prior to the Covid-19 infection that has been established in this patient. This is the first report of intermittent subclinical trigeminal nerve (motor component of anterior division of right mandibular nerve, V3), transient subclinical paresis of right-sided VII cranial (facial) nerve, and recurrent traumatic involvement of right side of anterior tongue as sequela in Covid-19, expanding the clinical picture and the immuno-pathological profile of the disorder. With care during eating solid foods, with specific instruction not to stack food in the oral cavity, there has been no recurrence over the last three years. (see photograph, published with permission) [1-5].

### Bibliography

1. Paszynska E., *et al.* "Prevalence of oral complications in the course of severe SARS-CoV-2 infection under mechanical non-invasive ventilation". *European Journal of Medical Research* 28.1 (2023): 293.
2. Lin W., *et al.* "The oral manifestations and related mechanisms of COVID-19 caused by SARS-CoV-2 infection". *Frontiers in Cell Neuroscience* 16 (2023): 1006977.
3. Ferreira MD., *et al.* "COVID-19 hospitalized patients and oral changes: a case-control study". *Clinical Oral Investigation* 27.8 (2023): 4481-4491.
4. Erbaş GS., *et al.* "COVID-19-related oral mucosa lesions among confirmed SARS-CoV-2 patients: a systematic review". *International Journal of Dermatology* 61.1 (2021): 20-32.
5. Merad M., *et al.* "The immunology and immunopathology of COVID-19". *Science* 375.6585 (2022): 1122-1127.