



## Dry Needling and Osteoarthritis Knee

**Subhanjan Das\***

Associate Professor, Department of Physiotherapy, Garden City University, India

**\*Corresponding Author:** Subhanjan Das, Associate Professor, Department of Physiotherapy, Garden City University, India.

**Received:** May 01, 2019; **Published:** May 24, 2019

**DOI:** 10.31080/ASOR.2019.02.0058

### Abstract

Osteoarthritis is the most common form of joint disease and its most common location is in knee [1]. Symptomatic knee OA occurs in 10% men and 13% in women aged 60 years or older [2]. Osteoarthritis is the leading cause of disability globally [3] and ranked among ten noncommunicable diseases for disability adjusted life years across world [4].

**Keywords:** Dry Needling; Knee; Osteoarthritis

### Introduction

Dry needling is a skilled intervention that uses a thin filiform needle to penetrate the skin and stimulate underlying myofascial trigger points, muscular, and connective tissues for the management of neuromusculoskeletal pain and movement impairments [5]. Physiotherapists use dry needling to stimulate neural, muscular and connective tissues [6]. Dry needling in various forms has been used in the physiotherapeutic treatment of osteoarthritis.

### Clinical features of osteoarthritis knee

The primary clinical feature of osteoarthritis is pain [7]. Persistent knee pain, limited morning stiffness, and reduced function are the three symptoms that are recommended for the diagnosis of knee OA by the EULAR [8].

Tibiofemoral joint OA is characterised by joint effusion, osteophytes, diminished flexion range, mediolateral instability and varus deformity. On the other hand in patellofemoral osteoarthritis, valgus deformity, grossly reduced quadriceps strength, pain on patellofemoral joint compression difficulty descending stairs and coarse crepitus is seen with a history of marked swelling [9]. Malalignment, like genu valgus or varus can be seen in advanced cases [10].

Muscle impairment is another well known clinical feature of knee osteoarthritis. Quadriceps, hamstrings and hip musculatures are most commonly involved [11]. This inability to completely activate a muscle secondary to joint dysfunction is known as arthrogenic muscle inhibition [12].

There has also been findings of altered hamstring quadriceps muscle balance in OA knee, with increased activation of hamstrings compared to the normal population [13].

Myofascial trigger point is prevalent in knee osteoarthritis especially in quadriceps, vastus medialis and lateralis are the usual areas [14].

Together all these factors contribute to reduction in the functional capacity and impaired activities of daily living.

### Dry needling in osteoarthritis

The effects of trigger point dry needling in the treatment of knee osteoarthritis has been well documented [15,16].

Dry needling has been found to be superior to acupuncture and sham acupuncture [17]. Needling by physiotherapists have been found as a cost effective use of health care resources [18].

Percutaneous electrical nerve stimulation is a more recent form of dry needling used by the physiotherapists. PENS has been associated with no morbidity, good pain relief, and increased function in patients with knee osteoarthritis [19]. A randomised sham controlled trial found PENS to be significantly better in reducing pain and stiffness and in improving function [20].

### Dry needling: Schools and types

Dry needling has been developed in various geographical locations and therefore there are various schools of needling [21].

Initial use of dry needling almost exclusively targeted myofascial trigger point (Trigger point dry needling). One of the pioneer of deep dry needling (needling into the trigger point) was Czech physician Karel Lewit in 1979. Peter Baldry started another type of dry needling called superficial dry needling. In the early 1980 [22]. Dr Yun Tao Ma, in his book biomedical acupuncture talks about primary and secondary reflex points [23]. The spinal segment sensitization technique of Chan Gunn stresses on needling the small spinal musculatures for peripheral conditions. Percutaneous electrical nerve stimulation is yet another variation of needling which incorporates TENS with needling.

### Proposed mechanism of action of dry needling in osteoarthritis knee

#### Neurophysiological Effects: Effects on central sensitization

The most likely mechanism of pain relief through needle stimulation is hyper stimulation analgesia [24]. Needling around knee is purported to release a number of neurotransmitters including endorphins, serotonin and cortisol.

#### Segmental inhibition/Gate control

When a needle is rapidly thrust into the soft tissue around knee, the sensory response evoked leads to a large diameter sensory afferent proprioceptive input into the spinal cord. This could have a "gate-controlling" effect of blocking the intra-dorsal horn passage of noxious information generated in the nociceptors [25].

#### Release of endogenous opioids

Pain following the insertion of the needle around knee releases opioids which is mediated through the PAG in the brainstem [26].

#### Effect on the release of neurotransmitters: Serotonin and noradrenaline

Stimulation of A $\delta$  nerve fibers may also activate the serotonergic and noradrenergic descending inhibitory system [26].

#### Conditioned pain modulation (CPM)

Patients with chronic knee pain have impaired CPM. Depressed CPM will lead to a reduction of endogenous pain inhibition and can contribute to a chronic pain state [27]. Several reviews have hypothesized that needling may affect CPM positively and can produce a reduction in chronic pain.

#### Effects on peripheral sensitization

The concentrations of SP and CGRP are found to be higher in the vicinity of active myofascial trigger points compared to latent

ones or normal muscle tissue. After Dry Needling of trigger points such as those present in quadriceps in osteoarthritis knee, SP and CGRP concentrations reduce significantly compared to their pre-DN values [27].

#### Effects on the taut band

DN to a trigger point region could effectively suppress spontaneous electrical activity which is a hallmark of trigger point. The process also produces a local twitch response, which normalizes the chemical environment [28].

#### Percutaneous electrical nerve stimulation (PENS)

PENS is a relatively newer modality used almost exclusively for pain relief and it has shown a lot of potential in reducing osteoarthritis pain. Studies have shown PENS provide significantly better and lasting analgesia with minimal side effects [29].

It is a combination of Dry needling and TENS. There are two primary pain relief mechanisms which can be stimulated: the Pain gate mechanism and the Endogenous Opioid System. Pain relief by means of the pain gate mechanism involves activation (excitation) of the A beta (A $\beta$ ) sensory fibers, and, reduces the transmission of the noxious stimulus from the 'C' fibers, through the spinal cord and hence on to the higher centers. The A $\beta$  fibers appear to appreciate being stimulated at a relatively high rate (in the order of 90 - 130 Hz or pps). Clinically it is important to enable the patient to find their optimal treatment frequency – which will almost certainly vary between individuals [30].

An alternative approach is to stimulate the A delta (A $\delta$ ) fibers which respond preferentially to much lower rate of stimulation (in the order of 2 - 5 Hz), which will activate the opioid mechanisms, and provide pain relief by causing the release of an endogenous opiate (encephalin) in the spinal cord which will reduce the activation of the noxious sensory pathways [5].

The released opioids may generate their analgesic action at peripheral, spinal and supraspinal sites. However, other neurochemicals like GABA have been implicated in TENS analgesia [31].

#### Conclusion

Osteoarthritis is a widely prevalent degenerative condition which produces considerable morbidity. Dry needling in various forms has been administered in the management of osteoarthritis. While the exact mechanism of action is hard to predict it appears dry needling has reasonable usefulness in the treatment of osteoarthritis knee.

## Bibliography

1. Creamer P and Hochberg Mc. "Osteoarthritis". *Lancet* 350.9076 (1997): 503-508.
2. Zhang Y and Jordan JM. "Epidemiology of osteoarthritis". *Clinics in Geriatric Medicine* 26.3 (2010): 355-369.
3. Vos T, *et al.* "Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010". *Lancet* 380.9859 (2012): 2163-2196.
4. Murray CJ, *et al.* "Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010". *Lancet* 380.9859 (2012): 2197-2223.
5. Alexandria VA. "APTA Public Policy, Practice, and Professional Affairs Unit. Description of dry needling in clinical practice: An education resource paper". *American Physical Therapy Association* (2013): 2.
6. James Dunning, *et al.* "Dry needling: a literature review with implications for clinical practice guidelines". *Physical Therapy Reviews* 19.4 (2014): 252-265.
7. Neogi T. "The epidemiology and impact of pain in osteoarthritis". *Osteoarthritis Cartilage* 21.9 (2013): 1145-1153.
8. Zhang W, *et al.* "EULAR evidence-based recommendations for the diagnosis of knee osteoarthritis". *Annals of the Rheumatic Diseases* 69.3 (2010): 483-489.
9. Peat G, *et al.* "Clinical features of symptomatic patellofemoral joint osteoarthritis". *Arthritis Research and Therapy* 14.2 (2012): R63.
10. Heidari B. *Rheumatic diseases*. 1st ed. Babol: Iran Babol university of medical sciences publication; (2002).
11. Ali H, *et al.* "Zeni, Lynn Snyder. Muscle Impairments in Patients With Knee Osteoarthritis". *Sports Health* 4.4 (2012): 284-292.
12. Rice DA and McNair PJ. "Quadriceps arthrogenic muscle inhibition: neural mechanisms and treatment perspectives". *Seminars in Arthritis and Rheumatism* 40.3 (2010): 250-266.
13. Hortobágyi T, *et al.* "Altered hamstring-quadriceps muscle balance in patients with knee osteoarthritis". *Clinical Biomech (Bristol, Avon)* 20.1 (2005): 97-104.
14. Sánchez-Romero EA, *et al.* "Clinical features and myofascial pain syndrome in older adults with knee osteoarthritis by sex and age distribution: A cross-sectional study". *Knee* 26.1 (2019): 165-173.
15. Woehrle J, *et al.* "Dry Needling and its Use in Health Care - A Treatment Modality and Adjunct for Pain Management". *Journal of Pain and Relief* 4 (2015): 194.
16. James Dunning, *et al.* "Dry needling: a literature review with implications for clinical practice guidelines". *Physical Therapy Reviews* 19.4 (2014): 252-265.
17. Itoh K, *et al.* "Trigger point acupuncture for treatment of knee osteoarthritis--a preliminary RCT for a pragmatic trial". *Acupuncture in Medicine* 26.1 (2008): 17-26.
18. David GT, *et al.* "Cost-Effectiveness of Acupuncture Care as an Adjunct to Exercise-Based Physical Therapy for Osteoarthritis of the Knee". *Foster Physical Therapy* 91.5 (2011): 630-641.
19. Kang RW, *et al.* "Prospective randomized single-blinded controlled clinical trial of percutaneous neuromodulation pain therapy device versus sham for the osteoarthritic knee: a pilot study". *Orthopedics* 2007 30.6 (2008):439-445.
20. a-peng He; Jun Zhang; Zheng-fa Bai, Percutaneous Electrical Nerve Stimulation for Chronic Knee Pain: A Randomized, Sham-controlled Trial". *Alternative Therapies in Health and Medicine* 23.7 (2017): 233-237.
21. Jan Dommerholt. "Dry needling — peripheral and central considerations". *Journal of Manual and Manipulative Therapy* 19.4 (2011): 223-227.
22. Baldry P. "Superficial Versus Deep Dry Needling". *Acupuncture in Medicine* 20.2-3 (2002): 78-81.
23. Biomedical Acupuncture for Sports and Trauma Rehabilitation: Dry Needling Techniques Elsevier - Health Sciences Division.
24. Li-Wei Chou, *et al.* "Probable Mechanisms of Needling Therapies for Myofascial Pain Control," *Evidence-Based Complementary and Alternative Medicine* (2012): 11.
25. Chou LW, *et al.* "Needling therapy for myofascial pain: recommended technique with multiple rapid needle insertion". *Bio-medicine* 4.2 (2015): 13.
26. David M, *et al.* "Central modulation of pain evoked from myofascial trigger point". *The Clinical Journal of Pain* 23.5 (2007): 440-448.
27. Wiebke Grashorn, *et al.* "Age-Dependent Decline of Endogenous Pain Control: Exploring the Effect of Expectation and Depression". *PLoS One* 8.9 (2013): e75629.
28. Hsieh YL, *et al.* "Spinal cord mechanism involving the remote effects of dry needling on the irritability of myofascial trigger spots in rabbit skeletal muscle". *Archives of Physical Medicine and Rehabilitation* 92.7 (2011):1098-1105.

29. Ossipov MH. "The perception and endogenous modulation of pain". *Scientifica* (2012).
30. Melzack R and P Wall. "Pain mechanisms: A new theory". *Science* 150 (1965): 971-979.
31. Josimari M DeSantana, *et al.* Effectiveness of Transcutaneous Electrical Nerve Stimulation for Treatment of Hyperalgesia and Pain.

**Volume 2 Issue 6 June 2019**

**© All rights are reserved by Subhanjan Das.**