



## Impact of Maternal and Early Life Undernutrition/Anemia on Mental Functions

**Kailash Nath Agarwal\* and Dev Kumari Agarwal**

*Health Care and Research Association for Adolescents, Noida, India*

**\*Corresponding Author:** Kailash Nath Agarwal, Health Care and Research Association for Adolescents, Noida, India.

**Received:** December 14, 2018; **Published:** January 22, 2019

### Abstract

Malnutrition refers to deficiencies, excesses or imbalances in intake of energy, protein and/or other nutrients and refers to both under nutrition and over nutrition. New classification scheme proposed by American Society for Parenteral and Enteral Nutrition incorporates chronicity, etiology, mechanisms of nutrient imbalance, severity of malnutrition and its impact on outcomes. In growing economies both under nutrition as well as over nutrition are observed with associated health consequences.

Intrauterine and early life under nutrition result in impairment of growth and development, intelligence, behavioral, conceptual and sensory motor development in preschool years. A child is stunted if the body is proportionate yet when compared to a normal child of the same age, he/she is shorter. In a wasted child, body fat and muscle reserves are broken down to maintain essential functions leading to weight loss. A child who is stunted and wasted is both short and thin compared to that of a normal child. Malnutrition in childhood affects IQ, cognitive function, persistence of soft neurological signs and affects higher-order thinking skills in adolescents. The changes that occur early in iron deficiency/anemia (pregnancy anemia in Bangladesh 77% and India 86%) may account for much of the long-term impact on decreased cognitive abilities. As anemia or iron deficiency affect brain iron content and neurotransmitters, irreversibly.

**Keywords:** Malnutrition; Definition; Maternal; Low Birth Weight; Intrauterine Growth Retardation; Child; Stunted; Wasted; Anemia; Iron Deficiency; Brain Development

### Malnutrition

The World Bank estimates that India has highest number of children suffering from malnutrition in the world. The prevalence of underweight children in India is among the highest in the world, and is nearly double that of Sub-Saharan Africa with dire consequences for morbidity like retardation of physical growth and mental development, mortality, productivity and economic growth. The 2015 Global Health Index (GHI) Report [1] ranked India 20<sup>th</sup> amongst leading countries with a serious hunger situation. Despite of economic growth in India, India's hunger is still worse than North Korea or Sudan. And a child raised in India is more likely to be malnourished than Somalia. Various studies suggest that the biggest reason for India's malnutrition is poor sanitation, polluted water, unclean vegetables, adulteration and illiteracy. Because of poor sanitation situation, more children in India than North Ko-

rea, Sudan and Somalia are exposed to bacteria. The bacteria sicken them, and make it hard for children to consume nutrients, which results in malnutrition. 620 million people in India don't have a toilet in their house and they use public toilet or defecate in fields or any open space (Prime Minister of India has initiated a national program "Swachh Bharat" to meet the much needed in sanitation). In addition, the air quality in India is among the worst in the world. UNICEF is also recognizing the poor sanitation as one of the reasons for malnutrition. In addition, cost of food, no easy affordability of pulses (source of protein for vegetarians) and vegetables/fruits make it worse. Women who suffer malnutrition are less likely to have healthy babies. Further they generally lack proper knowledge in nurturing/feeding children. Consequently, newborns/infants in early life are unable to get adequate amount of nutrition from their mothers.

**The chapter on ‘Malnutrition’ will cover aspects of maternal/ childhood nutrition and associated health consequences:**

- Definition of Malnutrition
- Nutrition during Pre-pregnancy, Pregnancy and Lactation Stages
- Health Consequences of Intrauterine and Childhood Malnutrition
- Effects of Malnutrition and/or Iron Deficiency on Brain

**Definition of Malnutrition**

**Malnutrition**

According to the World Health Organization [2], malnutrition refers to deficiencies, excesses or imbalances in intakes of energy, protein and/or other nutrients.

Malnutrition refers to deficiencies, excesses or imbalances in the intake of, energy, protein and/or other nutrients

Malnutrition includes both Under nutrition and Over nutrition

# Under nutrition is the result of food intake that is Continuously insufficient to meet dietary energy requirements, poor absorption and/or poor biological use of nutrients consumed.  
# Usually results in loss of body weight and/or obesity

# over nutrition refers to a chronic condition where intake of food is in excess of dietary energy requirements.  
# Usually results in overweight and/or obesity

**Figure a:** DWHO. WHO Child Growth Standards. Available at: [http://www.who.int/nutrition/media\\_page/backgrounders\\_4\\_en.pdf](http://www.who.int/nutrition/media_page/backgrounders_4_en.pdf). Accessed on: August 5<sup>th</sup> 2014.

Contrary to the common usage, the term ‘malnutrition’ correctly includes both under nutrition and over nutrition.

- Under nutrition is defined as the result of food intake that is continuously insufficient to meet dietary energy requirements, poor absorption and/or poor biological use of nutrients consumed. It usually results in loss of body weight.
- Over nutrition refers to a chronic condition where the intake of food is in excess of dietary energy requirements. Over nutrition usually results in overweight and/or obesity.

**Redefining Pediatric Malnutrition: Paradigm Shift towards Etiology-based Definition-American Society for Parenteral and Enteral Nutrition (ASPEN) [3] has proposed:**

New classification scheme incorporates chronicity, etiology, mechanisms of nutrient imbalance, severity of malnutrition, and its impact on outcomes. Based on its etiology, malnutrition is either:

- **Illness-related:** Secondary to one or more diseases/injury or
- **Non-illness-related:** Caused by environmental/behavioral factors, or both

‘Pediatric malnutrition (under nutrition) is defined as an imbalance between nutrient requirements and the intake that results in cumulative deficits of energy, protein, or micronutrients that may negatively affect growth, development and other relevant outcomes’.

**Malnutrition Statistics**

India with surplus food continues to see malnutrition: manifesting as stunting, wasting, underweight and anemia. Unfortunately, these indices are on a rise as observed in National Family Health Surveys (NFHS) II vs III. According to the 2011 Hunger and Malnutrition survey conducted by the Nandi Foundation, 42 percent of Indian children under five years old are underweight - almost double the rate of sub-Saharan Africa. This is in spite of the fact that today India is one of the major food producers in the world. The food sector contributes to about 18% of India’s GDP.

**In terms of world ranking, India stands**

- FIRST in the production of cereals, milk and livestock population
- SECOND in producing fruits and vegetables
- Ranks among in the top 5 in producing rice, wheat, groundnuts, tea, coffee, tobacco, spices, sugar and oilseeds [4].

**Comparison of Maternal and Child Health Indices**

Indicators	India <sup>a</sup>	Sri Lanka <sup>b</sup>	Bangladesh <sup>c</sup>
Neonatal Mortality Rate	29	6	24
Infant Mortality Rate	42	8	33
Maternal Mortality Rate (per 10,000 live births)	210	39	220
Institutional Delivery	46	98	28

**Table a:** It is estimated that 67,000 women die of puerperal causes each year in India<sup>d</sup>

a. UNICEF, UNICEF - India - Statistics, 2014. Available at: [http://www.unicef.org/infoby country /india\\_statistics.html](http://www.unicef.org/infoby country /india_statistics.html).

b. UNICEF, UNICEF - Sri Lanka - Statistics, 2014. Available at: [http://www.unicef.org/infoby country /sri\\_lank a\\_statistics.html](http://www.unicef.org/infoby country /sri_lank a_statistics.html).

3c UNICEF, UNICEF - Bangladesh - Statistics, 2014. Available at: [http://www.unicef.org/infoby country /bangladesh\\_bangladesh\\_statistics.html](http://www.unicef.org/infoby country /bangladesh_bangladesh_statistics.html).

d. Bhowmik S. Child mortality: Will India achieve the 2015 target? *BMJ*. 2013;346:1502.

## Nutrition during Pre-pregnancy, Pregnancy and Lactation Stages

We will discuss community-based case studies, which outline the importance of maternal nutrition on maternal health and child outcome [5].

In the case study- 1 the impact of maternal under nutrition can be observed from a prospective study conducted in rural setting of eastern Uttar Pradesh, Varanasi. The study evaluated the birth weight pattern in chronic as well as currently undernourished pregnant women. In the study, anthropometry, hemoglobin, dietary intake, birth weight, fundal height and abdominal girth data of 3700 eligible pregnant women at  $16 \pm 2$ ,  $28 \pm 2$  and  $36 \pm 2$  weeks of gestation were recorded. Outcome measure was birth weight pattern of newborns.

Results indicated that out of 3700 births, 34.6% were of low birth weight and only 8.2% weighed more than 3000 g. Fundal height was  $<24.5$  cm at 28 weeks of gestation (1368 women) and was associated with higher low-birth-weight deliveries. Fundal height did not increase during 35-39 weeks of gestation (was lower by 5 cm as compared to normal).

With respect to the maternal weight gain, women in later pregnancy (during 35 - 43 weeks of gestation), showed weekly weight gain of 15 - 53 g, only. In contrast, healthy women gained 400 g/week in the second and third trimesters. Total weight gained during the entire period of pregnancy was about 6 kg only, whereas ideally the gain should be between 13 and 18 kg [5].

For understanding the importance of nutrition in pregnancy, it is useful to additionally note the data of World War II-famine in Leningrad (USSR) where women were already undernourished. About 50% of the children who were born were low birth weight (LBW) and birth weight fell by 530g. In contrast, during the Dutch famine, only 9% of the infants born to previously well-nourished Dutch women had LBW and birth weight fell by 327g. The Dutch famine was imposed on a previously well-nourished population and the official daily food for the general adult population gradually decreased from 1800 calories in December 1943 to 1400 calories in October 1944 to below 1000 calories in the late November 1944. December 1944 to April 1945 was the peak of the famine during which the official daily ration fell abruptly to about 400-800 calories. Even though pregnant and lactating women had extra food during the famine, these extra supplies could no longer be provided at the height of the famine. What is unique about Dutch Famine is that it was imposed on a previously well-nourished population [6].

These two historical examples illustrate the importance of pre-pregnancy nutrition and its impact on pregnancy outcome.

Let us now look at the community-based case study - 2.

We will discuss the impact of ICDS (Integrated Child Development Services) program on maternal nutrition and birth weight in rural Varanasi [7]. India's ICDS (nutrition + education + child care program) has been adapted from the Women Infants and Children (WIC) in USA.

5289 pregnancies were registered during 1987-1993. All live births were seen within 48 hr and their weight was recorded.

- In the ICDS, 916 pregnant women were nutritionally supplemented and 1453 were un supplemented, both groups received healthcare and nutrition education.
- There were 1748 pregnancies from the non-ICDS villages, and they received simple healthcare.

The ICDS-supplemented mothers gained 100 g more in pregnancy, birth weight increased by 58 g, preterm's and low birth weights reduced by 12.9% and 29.4%, respectively, as compared to the un supplemented mothers (ICDS). In the un supplemented ICDS area, preterm and low-birth-weight births reduced by 44.2% and 22.4%, respectively, as compared to non-ICDS area pregnancies. The un supplemented (ICDS) had higher birth weight by 25 g as compared to the non-ICDS. There were 16.2%, 11.0% and 12.9% having birth weight over 3000 g in these three areas, respectively. The multiple regression analysis showed that the increased weight gain in pregnancy, length of gestation, caloric intake and term hemoglobin were significantly associated with birth weight.

Thus, nutritional supplementation can benefit undernourished pregnant women and lead to better maternal and fetal outcomes.

### Maternal anemia and its consequences

- The anemia prevalence in three countries, India, Bangladesh and Sri Lanka are [8,9] shown in table 1. The prevalence of anemia in pregnancy is 86% and 81% in lactation in India and is responsible for 20-40% maternal deaths (MMR > 200). Indian Council Medical Research 2001 showed that 84.7% of mothers are anemic (hemoglobin < 11.0 gm/dl), of these 9.9% are with severe anemia (hemoglobin < 7.0 gm/dl).
- Agarwal, *et al.* [8] showed in 7 states pregnancy anemia was in 86.1% (severe anemia in 9.7%); anemia during lactation being 81.7% (severe anemia - 7%). Thus, pregnancy/lactation anemia is a serious health hazard for mother and child in India. Therefore, we must discuss ef-

ffects of pregnancy anemia on fetal outcome including brain growth.

- Anemia is a serious health hazard for mother and child in India. Therefore, we must discuss affects of pregnancy anemia on fetal outcome including brain growth. Fetal growth depends, to a large extent, on the availability of iron from the mother. Placenta formation needs iron in the first 2 weeks. It continues to trap iron at a gradient even when fetus is removed in animals. Normally, placental iron transfer to fetus increases 3-4 times during 20-37 weeks of gestation.

Countries	Anemia in pregnancy	Anemia in lactation
India [8]	86.1%	81.7 %
Sri Lanka [9]	16.0%	—
Bangladesh [8]	77.0%	77.0%

**Table 1:** Prevalence of anemia in India, Srilanka and Bangladesh.

In maternal anemia, fetal liver iron is low resulting in early-onset anemia. There is an increased gradient in the presence of maternal iron deficiency for transport of iron from mother to fetus but the transport remains proportionate to the degree of maternal hypoferrremia. Fetal Liver iron stores are reduced significantly in maternal hypoferrremia, and fetal brain iron content and neurotransmitters are reduced irreversibly, and their receptors are altered [10].

**Health Consequences of Intrauterine and Childhood Malnutrition**

Maternal suboptimal nutrition during pregnancy results in Infants born with IUGR (Intra Uterine Growth Retardation-Birth weight less than 10<sup>th</sup> centile for gestational age) are at an increased risk of prenatal morbidity and mortality [11,12]. Furthermore, when compared to infants with appropriate intrauterine growth, infants with IUGR have greater rates of physical (Stunting), neurological and mental impairment [13]. Globally, about 30 million infants suffer from IUGR every year, and nearly 75% of all affected infants are born in Asia [13].

Low birth-weight infants are at an increased risk of morbidity, impaired mental development and mortality. The risk of neonatal death is reported to be four times higher in infants who weigh 2000-2499 g at birth when compared to those who weigh 2500-3499 g. The greater the severity of growth restriction within the low birth-weight category, the higher the risk of mortality [14].

Low birth weight is a strong predictor of size in later life since infants born with IUGR seldom catch-up to normal size during childhood [3]. Low birth weight increases the risk for development of adult metabolic syndrome. Low-birth-weight babies have an increased risk of later life diseases such as coronary heart disease, hypertension, type 2 diabetes and other chronic diseases [15-18].

**Maternal anemia and its consequences**

Intrauterine Growth Retardation [15,16]	Low Birth Weight
Increased risk of perinatal morbidity and mortality [15]	Increased risk of morbidity, impaired mental development and mortality [17]
Greater incidence of physical, neurological and mental impairment when compared to infants with appropriate intrauterine growth [16]	Greater the severity of growth restriction, higher the mortality risk [18]
Global incidence: 30 million infants every year (nearly 75% of these are born in Asia) [16]	Increased risk for development of adult metabolic syndrome [17,18]
Maternal suboptimal nutrition during pregnancy results in intrauterine growth restriction and the birth of newborns with low birth weight [1]	

**Table b**

**Childhood Undernutrition**

India and Bangladesh have the highest proportion of children who are underweight. In India, 20% of children are wasted (Table 2). National Family Health Survey in India indicated that 1 in 6 children aged between 38 and 57 months are wasted. It is encouraging to note that the corresponding figures in Sri Lanka are on the lower side [11,12].

Indicator	India	Sri Lanka	Bangladesh
<b>Underweight (moderate and severe) (weight for age)</b>	43.0% (29.4%)	29.0%	48.0%
<b>Stunting (height for age)</b>	48.0% (38.8%)	14.0%	43.0%
<b>Wasting (weight for height)</b>	20.0% (15.0%) (1 in 6 Children 38-57 month) [11,12]	14.0%	13.0%

**Table 2:** Underweight, stunted and wasted children in India, Sri Lanka and Bangladesh.


In parenthesis recent “Rapid survey on children”, UNICEF and Government of India figures are given 2015.

An underweight child has a weight-for-age Z-score that is at least 2 SD below the median for the National Child Growth Standards. This condition can result from either chronic or acute malnutrition, or both.

A stunted child has a height-for-age Z-score that is at least 2 standard deviations (SD) below the median for the National Child

Growth Standards. Chronic malnutrition is an indicator of linear growth retardation that results from failure to receive adequate nutrition over a long period and may be exacerbated by recurrent and chronic illness [14]. These chronic malnourished have serious health consequences i.e. short stature with poor strength and structural and functional brain damage.

### Stunting Consequences



Evidence from developing countries indicates that being stunted between 12 and 36 months of age is associated with poorer cognitive performance and lower school achievement in middle childhood.

A wasted child has a weight-for-height Z-score that is at least 2 SD below the median for the National Child Growth Standards. Wasting represents a recent failure to receive adequate nutrition and may be affected by recent episodes of diarrhea and other acute illnesses.

### Effects of Malnutrition and/or Iron Deficiency on Brain

The intrauterine growth retarded offspring of undernourished mothers [19,20]:

- Showed hypotonia in 72% and hypo excitability in 56%.
- Limp posture, poor recoil of limbs, incomplete Moro's and crossed extensor responses.
- Their EEG had shortening of sleep cycle (REM and NERM), the reduction was marked for REM babies weighing < 2000g. There was some inter and intra hemispheric asymmetry and abnormal paroxysmal discharges; suggesting dysmaturity of brain
- Malnutrition in childhood [21-30].
- Affects IQ, cognitive function and persistence of soft neurological signs, impaired repetitive speed movements with higher degree of overflow and dysrhythmia with EEG changes supporting abnormalities in frontal lobe.

- The deficit in higher mental abilities and prolonged reaction time persisted in later years during adolescence.
- Brain MRI: Both frontal lobes showed reduction in size anteriorly as well as posteriorly with loss of asymmetry.
- P3 latency was normal, but the P2 and P3 amplitudes were higher suggesting neuronal compensation.
- Nutrition supplementation in pregnancy, lactation, infancy childhood showed improvement in mental functions [28,30].

### Iron depletion and brain

Neonatal Period: Infants with cord serum ferritin concentrations < 35 mcg/L have abnormal auditory recognition memory, infants do not discriminate a familiar stimulus (e.g. maternal voice) from a stranger's voice [31].

- Term infants born to iron deficient anemic mothers also exhibit alterations in temperament and activity [32].
- Mental Functions in Rural Anemic Primary School Children (nutrition controlled) Mental functions in nutrition controlled 388 (6-8 yr of age), matched for social and educational status were studied by WISC and arithmetic test to assess "Intelligence, Attention and Concentration". Anemia did not affect intelligence, except subtest-digit span. In Arithmetic test 'Attention and Concentration' was poor in anemic children [33].
- Those anemic in infancy in spite of iron therapy in early life remain poor in school achievements and have presence of soft neurological signs. Central conduction time for auditory brain stem responses at 6, 12, 18 months were slower. Heart rate variability during sleep/wakefulness was less mature [34].

### Anemia and brain iron [35,36]

Studies in children 8-12 years of age having anemia (nutritional, serum ferritin < 20ng) or due to thalassemia (serum ferritin > 1000ng), on brain MRI spectroscopy showed similar iron content in globus pallidus, caudate and dentate nucleus in both the conditions. There was an increase in creatinine and aspartate and reduction in choline concentration. Such changes are also observed in Huntington's chorea and Alzheimer's disease. Reduction of choline, in latent iron deficiency is a significant effect. Choline is synthesized in the brain in very small amounts; its uptake is Na<sup>+</sup> dependent, which requires oxygen.

### Latent Iron deficiency in rat model

Dietary iron depletion IN PREGNANT RATS- reduced fetal hepatic iron and selectively brain iron content.

### Iron content decreased [37]

In Cerebral cortex 17%, Cerebellum 18%, Hypothalamus 19%, Mid brain 21%, and Corpus striatum 32%

- No change in – medulla oblongata and pons.
- Fetal brain iron content did not change after maternal “Fe” supplementation.

Low brain “Fe” content was associated with significant alteration in brain Cu, Zn, Ca, Mn, Pb and Cd levels.

### Fetal brain iron reduction and latent iron deficiency in post weanling rats altered neurotransmitters, irreversibly [38,39]

- Glutamate metabolism- GAD, GDH, GABA-T and their receptors-Binding of H3 Muscimol at pH 7.5 and 1mg protein/assay (GABA receptor) increased by 143%, but glutamate receptor binding decreased in the vesicular membranes of latent iron deficient rats by 63%. The significant effects on neurotransmitter receptors (glutamate mediators) during early stages of “Fe” deficiency clearly indicate the irreversible deficit in both excitatory and inhibitory pathways of the CNS.
- TCA-cycle enzymes-mitochondrial NAD+ linked dehydrogenase
- Catecholamine metabolism- Whole brain-dopamine, nor-epinephrine, tyrosine and TAT; Corpus striatum – same as in whole brain, except TAT increased.
- 5-HT metabolism- Tryptophan, 5-HT, 5-HIAA.

The neurotransmitter changes were specific to “Fe” deficiency as in under nutrition these get corrected partially or completely on rehabilitation [35].

Iron is essential for myelination - the process by which the brain produces a fatty insulation around the nerves. Iron has clear effects on the function of dopamine and probably serotonin, two brain chemicals that help send and receive signals (neurotransmitters) and have many roles in the brain.

### Conclusions

- Traditionally, malnutrition refers to deficiencies, excesses or imbalances in intake of energy, protein and/or other nutrients and refers to both under nutrition and over nutrition. New classification scheme proposed by ASPEN incorporates chronicity, etiology, mechanisms of nutrient imbalance, severity of malnutrition and its impact on out-

comes.

- Nutritional supplementation can benefit undernourished pregnant women and lead to better maternal and fetal outcomes as shown by the Varanasi rural study. Intrauterine and early life under nutrition results in impairment of growth and development, intelligence, behavioral, conceptual and sensory motor development in preschool years. Malnutrition in childhood affects IQ, cognitive function, persistence of soft neurological signs and affects higher-order thinking skills in adolescents.
- Changes that occur early in the course of iron deficiency may account for much of the long-term impact on decreased cognitive abilities, due to irreversible brain iron and neurotransmitter changes.

### Bibliography

1. International Food Policy Research Institute, Global Hunger Index, (2015).
2. WHO. WHO Child Growth Standards (2104).
3. Mehta NM., *et al.* “Defining pediatric malnutrition: A paradigm shift toward etiology-related definitions”. *Journal of Parenteral and Enteral Nutrition* 37 (2013): 460-481.
4. Indian National Science Academy, Pursuit and Promotion of Science: The Indian Experience, (2001).
5. Agarwal S., *et al.* “Birth weight pattern in rural undernourished pregnant women”. *Indian Pediatric* 39 (2002): 244-253.
6. Stein Z., *et al.* “Famine and Human Development”. The Dutch Hunger Winter of 1944-1945. New York, (1975).
7. Agarwal KN., *et al.* “Impact of the Integrated Child Development Services (ICDS) on maternal nutrition and birth weight in rural Varanasi”. *Indian Pediatric* 37 (2000): 1321-1327.
8. Agarwal KN., *et al.* “Anemia in pregnancy interstate differences”. *Indian Journal of Medical Research* 124.2 (2006): 173-184.
9. Agampodi S. “Screening for anemia during pregnancy: what evidence suggests. Maternal Health Task Force: 2011” (2014).
10. Agarwal KN., *et al.* “Effect of maternal iron status on placenta, fetus and newborn”. *Indian Journal of Medical science* 5.9 (2013): 391-395.
11. UNICEF. Child Survival. The State of the World’s children (2008).
12. Arnold F., *et al.* “National Family Health Survey (NFHS-3) India”. Nutrition in India, 2005-2006.

13. Nutrition in Emergencies WHO.
14. Dewey KG., et al. "Long-term consequences of stunting in early life". *Maternal and Child Nutrition* 7 (2011): 5-18.
15. Belkacemi L., et al. "Maternal under nutrition influences placental-fetal development". *Biology of Reproduction* 83 (2010): 325-331.
16. Muhammad T., et al. "Maternal factors associated with intrauterine growth restriction". *Journal of Ayub Medical College Abbottabad* 22.4 (2010): 64-69.
17. Menon PSN. "Adult health consequences of being born with intrauterine growth retardation". In: *The Growth Infancy to Adolescence* 3<sup>rd</sup> ed. Chap 7 ed K N Agarwal. (2015): 87-92.
18. Barker DJ., et al. "Fetal origins of adult". *The Journal of Clinical Endocrinology and Metabolism* 87 (2002): 4657-4661.
19. Bhatia VP., et al. "Effect of intrauterine nutritional deprivation on neuromotor behavior of the newborn". *Acta Paediatrica Scandinavica* 68 (1979): 561-566.
20. Bhatia VP., et al. "Sleep cycle studies in babies of undernourished mothers". *Archives of Disease in Childhood* 55 (1980): 134-138.
21. Agarwal KN., et al. "Soft neurological signs and EEG pattern in rural malnourished children". *Acta Paediatrica Scandinavica* 78.6 (1989): 873-878.
22. Agarwal KN., et al. "Impact of chronic under nutrition on higher mental functions in Indian boys aged 10-12 years". *Acta Paediatrica* 84.12 (1995): 1357-1361.
23. Agarwal DK., et al. "Influence of malnutrition on cognitive development assessed by Piagetian tasks". *Acta Paediatrica Scandinavica* 78.1 (1989): 115-122.
24. Upadhyay S., et al. "Persistence of soft neurological signs in chronic undernourished children". *Nutrition Research* 15 (1995): 193-199.
25. Agarwal KN., et al. "Sequelae of early under nutrition on reaction time of rural children at 11-14 years". *Indian Journal of Medical Research* 107 (1998): 98-102.
26. Misra UK., et al. "Brain MRI and cognitive evoked potentials in rural chronically undernourished children". *Nutrition Research* 16 (1996): 1147-1451.
27. Agarwal DK., et al. "Effect of mid-day meal programme on physical growth and mental function". *Indian Journal of Medical Research* 90 (1989): 163-174.
28. Grantham-McGregor S. "A review of studies of the effect of severe malnutrition on mental development". *Journal of Nutrition* 125 (1995): 2233S-2238S.
29. Kar BR., et al. "Cognitive development in children with chronic protein energy malnutrition". *Behavior of Brain Function* 4 (2008): 31.
30. Grantham-McGregor SM., et al. "Effects of early childhood supplementation with and without stimulation on later development in stunted Jamaican children". *The American Journal of Clinical Nutrition* 66 (1997): 247-253.
31. Siddappa AM., et al. "Iron deficiency alters auditory recognition memory in newborn infants of diabetic mothers". *Pediatric Research* 55 (2004): 1034-1041.
32. Wachs TD., et al. "Relation of neonatal iron status to individual variability in neonatal temperament". *Development of Psychobiology* 46 (2005): 141-153.
33. Agarwal DK., et al. "Anaemia and mental functions in rural primary school children". *Annals of Tropical Paediatrics* 9 (1989): 194-198.
34. Walter T. "Effect of iron-deficiency anemia on cognitive skills and neuromaturation in infancy and childhood". *Food Nutrition Bulletin* 24 (2003): S104-S110.
35. Agarwal KN. "Iron and the brain: neurotransmitter receptors and magnetic resonance spectroscopy". 85 (2001): s147-s150.
36. Mittal RD., et al. "Effect of latent iron deficiency on gaba and glutamate neuroreceptors in rat brain". *Indian Journal of Clinical Biochemistry* 17 (2002): 1-6.
37. Shukla A., et al. "Effect of latent iron deficiency on metal levels of rat brain regions". *Biological Trace Element Research* 22.2 (1989): 141-152.
38. Shukla A., et al. "Latent iron deficiency alters gamma aminobutyric acid and glutamate metabolism in rat brain". *Experientia* 198945: 343-345.
39. Shukla A., et al. "Effect of latent iron deficiency on 5-hydroxytryptamine metabolism in rat brain". *Journal of Neurochemistry* 52 (1989): 730-735.

## Volume 2 Issue 2 February 2019

© All rights are reserved by Kailash Nath Agarwal and Dev Kumari Agarwal .