



Learning to Crack the Cleft Aesthetic Code in Unilateral Cleft Lip Surgery by Younger Cleft Trainees: Using Nuances and the Innovative Taipei Pear Analogy for Inferior Turbinate-flap Floor of Nose Reconstruction

Lotha B^{1*}, Bergonzani M² and Zeinalddin M³

¹Craniofacial Cleft Surgeon, Yemen Global Smiles, Sanaa, Yemen

²MD, Maxillo-Facial Surgery Division, Parma University Hospital, Italy

³Craniofacial Orthodontist Mohammad Orthodontic Center - Senior Lecturer Oman Dental College, Oman

*Corresponding Author: Lotha B, Craniofacial Surgery Fellowship, Yemen Global Smiles, Sanaa, Yemen.

Received: July 24, 2020

Published: August 18, 2020

© All rights are reserved by Lotha B., et al.

Abstract

Unilateral cleft lip surgery is part and parcel of every trainee cleft surgeon's armamentarium. As their cleft expertise is founded on solid principles and anthropometric measurements, the chances of error and iatrogenic deformity are significantly lower than that of the untrained surgeon who attempts cleft surgery. In this communication we would like to mention the possible dimensions of cleft surgery, and how one can improve outcomes by looking at different ways to ease the psychological burden of cleft children. The sequential repair makes for better aesthetic functional results and patients will be satisfied with the post-operative results when the surgery is properly executed, taking into consideration all the rules of engagement in tackling cleft challenges.

Keywords: Cleft Lip; Inferior Turbinate Flap; Median Tubercle; Cleft Training

Introduction

Unilateral cleft lip surgery is the commonest procedure performed at all smile centres across developing world smile centres. Every cleft surgeon starts with repair of the unilateral cleft lip before moving on to other more advanced surgeries. Accuracy of repair is always a concern and proper training is the need of the hour [1].

It is not so much about comparisons or picture perfect results, but using correct principles as a launching pad for future improvements even as younger cleft surgery trainees start their careers as primary cleft surgeons.

At the outset, it cannot be stressed enough that anthropometric measurements are absolutely essential for every cleft trainee. As advised by our dear mentor John Mulliken of Childrens MA,

we make sure trainees everywhere are familiar with the measurements and also practice the speciality at home before embarking on overseas missions [2]. Earlier this season, John had shared with us that the typical cleft lip repair of an untrained surgeon manifests itself as a poorly understood and badly executed repair.

Getting it right the first time would serve as the perfect launching pad for any cleft service as primary unilateral cleft surgery is the first test of a surgeon's skill at clefts. Failure to launch will inevitably lead to iatrogenic deformities where one deformity is replaced by another, as the unenviable process of serial expertise continues due to poor understanding of the basic principles [3].

Since the majority of the world's population with clefts live in developing countries, the need of the hour is to market the AAR-affordable, applicable and reproducible, take-home message for all

who follow this sub speciality with a passion for underprivileged cleft populations.

One of the key strategies in understanding the anatomy of change in unilateral cleft lip surgery, lies in a good knowledge of the orbicularis oris muscle-dermis complex [4]. Our focus is on the perfect simulation of a normal muscle dermis complex with the usual add-ons that, arguably create more happiness for our patients over the years.

Materials and Methods

We would like to propose five possible dimensions in unilateral cleft lip surgery. In poor countries with sparse resources, the 1st to 5th Dimension of cleft care is more or less a 1st to 5th Illusion of cleft care, as all of us are sadly aware. The first and second dimension are part and parcel of what we would term “Functional and Aesthetic Correction” during primary the cleft lip surgery in well-developed units, where the plastic surgeon is well versed with the different aspects of the procedure and is not trying to divide the two into separate entities.

The First dimension- reconstructive element which corrects the functional aspects and leads to proper function of repaired the cleft lip; this is what is so commonly encountered in developing world smile missions where this aspect alone is addressed without the equally important second step.


The Second dimension includes the small z and cheiloplasty procedures with Tajima primary rhinoplasty [5], which are aesthetic in nature, thereby adding an attractive aesthetic appearance after the correction; the usual aesthetic lip repairs in well-developed units.

The others are ancillary procedures involving scar modulation, gleaned from the interphase between Aesthetic Medicine and Plastic Surgery.

The Third dimension is offered by Aesthetic Medicine where the accidental discovery of “Aesthetic Botox” became a game changer for millions of aesthetic procedures [6]. It makes the aesthetic appearance even more attractive and supple by controlled muscle relaxation and improved scar outcomes.

The Fourth dimension includes Aesthetic Medicine enhancement procedures using Aesthetic Medicine in stepwise progression, using autologous fat filler to improve scar appearance [7], skin tone and volume correction.

The Fifth dimension involves the use of advanced lasers including nano second, fractional or picosecond lasers [8]. Combined with the fourth dimension, the overall results will be enhanced even further. There is an overlap between four and five, because a combined approach may be even more beneficial.



Cracking the Cleft Aesthetic Code

- Code A and B : Cupid's bow, lip vermilion, white roll, orbicularis oris, floor of nose, deviated caudal septum, alar cartilages
- Code C- post operative wound tension and scar
- Code D- post operative pigmentation and scar not improved by botox
- E- finer enhancement of scar and pigment

Dimension 1	Dimension 2	Dimension 3	Dimension 4	Dimension 5
Code A – NAM and Reconstruction Code A and B= one code	Code B- aesthetic recon One code for functional aesthetic recon	Code C- Botox Scar modulation providing supple scar without wound tension	Code D- fat graft, dermaneedling, tyrosinase pigment lighte Improvement of skin pigment and prevention of hypertrophic scar	Code E- lasers nano/pico second Enhancing skin appearance for selective photo thermomolysis by specifically targeting melanosomes
Cracked the code A by NAM, muscle dermal recon	Cracked the code B by fine adjustments of vermillion, philtrum, septal cartilage and alar cartilages	Cracked the code C by preventing wound tension and improved healing	Cracked the code D by taking care of pigmentation and scarring not addressed by botox	Cracked the code E by further improvement of overall scar and pigment by more advanced methods

Figure 1: Cracking the cleft aesthetic code (Courtesy L Bona 2020).

Steps proposed for cracking the cleft code:

- The C flap and modifications
- Inferior turbinate flap, L and M flap using the pear analogy and our innovative slinky S incision
- Vermillion and baby zee flaps
- Muscle dermis complex pars peripheralis to contralateral side to simulate philtrum
- Pars marginalis to prevent a lip notch
- Long muscle flap, without tension
- Upper orbicularis muscle fibres hitch to columella to prevent a long lip
- Caudal septum shift to midline by stitching muscle to septum causing cleft side medialization
- Tajima over correction 15-20% on the cleft side to intentionally keep the ala cartilage at higher level
- Cupid’s bow and philtral heights of muscle and skin made symmetrical
- Labial elements create the median tubercle of lip [9] with an attractive pout. Michela suggested the marginalis be dissected off the mucosa for about one mm and everting horizontal sutures used to increase the lip pout. We call this median tubercle modification, the M Parma code of the lip
- The difficult code involves reconstruction of the nasal floor using mucosal flaps with an inferior turbinate flap.

For the purpose of simplicity, one can try to make this a user friendly code by using the analogy of a continuous “skin of the pear”, a concept designed in Taipei during a casual discussion of the inferior turbinate flap [10]. Since we use the lateral L flap in nasal floor reconstruction, the concept uses AI (augmented imagination) and imagines that the mucosal and vestibular surfaces are like the skin of a pear; this is true for the continuous inferior turbinate surface facing the nasal cavity (Figure 2,3).

- The incisional approach to the inferior pole of the inferior turbinate flap is easy- trace along the Noordhoff point and incise between skin and mucosa up the vestibule and cut along inferior part of the turbinate-just follow the pear all around and connect to the superior pole
- A small transverse stab between the lower lateral cartilages is the superior entry point
- Next, we connect the pear and after removing the bone to prevent curling of the flap, rotate it down by 90 degrees; The superior pole edge is fixed to the pyriform edge to provide lining for the nose
- The inferior edge joins the L flap and is sutured as a turn over flap to the turned over CM flap
- This completes the floor of nose correction. The “Taipei pear analogy” has made floor of nose reconstruction much easier to understand!




Figure 2: Code M Parma for the lip (courtesy Bergonzani M, Parma).




Figure 3: Analogy of the Taipei slinky S pear incision and its progressions (courtesy L Bona).




Figure 4

Our usual sequence of closure

Muscle closure: CINCH SUTURE FIRST. The first stitch to be tied is the cinch suture which has to be tied, with slight over correction. We use internal and not external cinching sutures:

- The marginalis muscle below the Cupids bow is approximated to prevent a lip notch
- We create a dimple above 1 with muscle to muscle stitch at unequal heights –one thicker bite at non cleft side and thinner bite on the cleft side
- Above this we align muscle to muscle at unequal heights, similarly, including dermis on lateral side to simulate a dynamic philtrum
- We also fix the muscle to the columella base or caudal septum with slight overcorrection to medialize the repair and bring the deviated columella to the midline with slight over correction
- Using an internal cinch for the lateral ala base of cleft side to opposite columella soft tissue or caudal septum, we prefer overcorrection in all cases.

For easy sequencing, we perform the 5th internal cinching suture first, followed by 1 to 4. We prefer to use our innovative “skin hook” rule by downward traction to suture at equal distances without discrepancy. This ensures symmetrical closure of muscle and skin flaps.

Next, we perform the Tajima rim incision with slight over correction of 1mm on the cleft side ala. Releasing the domes adequately, we then transfix with the cleft dome by an inverted U suture using 4 0 nylon, where the cleft cartilage is kept 1 mm higher. Not every cleft surgeon is familiar with this technique and may use other methods or repair the nose later (older sample pics). Post operatively, a nasal conformer is inserted, with one side 1 mm higher. This is kept for 6 months to reshape the hypoplastic alar cartilage. We also tape the lip edges every 2 to 3 days with 3M tape, without involving the cheeks for 6 months.

Results

Observing the rules of engagement in cleft lip repair, the challenges are managed quite well using the anthropometric measurements and proper sequencing. The step-wise progression is very useful in getting a symmetric lip repair without deformity. The corresponding author’s results have been consistently uniform using these methods on many cleft children during mission outreach with the US Smile Train humanitarian programmes in Yemen and elsewhere.

Pic samples: Smile Train NYC and Yemen Global Smiles outreach pics 2010 - 2015 (from smile mission database) (Courtesy L Bona).




Picture 1: Left cleft lip (Tajima not done).



Picture 2: Left Cleft Lip (Tajima not done).



Picture 3: Left cleft lip (Tajima not done).



Picture 4: Microform left cleft lip.



Picture 5: Left cleft lip.

Discussion and Conclusion

Cracking the cleft aesthetic code is easy to learn by cleft trainees because the anatomical landmarks remain constant for every case. The results will not always be picture perfect because no two deformities are the same. Our aim is not to create a flawless designer lip, but show younger cleft surgeons the importance of anthropometric measurements and the principles of accurate lip repair by our long time mentor John Mulliken. Getting it reasonably good the first time round makes subsequent repairs a lot easier to manage. We hope this brief communication will be of help to trainee cleft surgeons even as they embark upon smile missions to lessen the burden of cleft children across the world.

Bibliography

1. Matthew Louis BS, *et al.* "Making the Grade in Global Cleft Care: Smile Train". *Craniofacial Trauma Reconstruction* 11 (2018): 1-5.
2. John B Mulliken., *et al.* "The Changing Faces of Children with Cleft Lip and Palate". *The New England Journal of Medicine* 351 (2004): 8.
3. Christian C Dupuis. "Humanitarian Missions in the Third World: A Polite Dissent". *Plastic and Reconstructive Surgery* 113.1 (2004).
4. Carolyn R Rogers., *et al.* "The Philtrum in Cleft Lip: Review of Anatomy and Techniques for Construction". *The Journal of Craniofacial Surgery* 25.1 (2014) 457-461.
5. BA Coglán Boorman. "Objective evaluation of Tajema secondary cleft lip nose correction". *BJPS* 49(1996): 457-461.
6. CC Chang and Philip KT Chen. "Botulinum Toxin to improve results in cleft lip repair: a double blinded, randomized vehicle controlled clinical trial". *PLOS One* (2014).
7. Christian Jones, Donald Mackay. "Autologous fat grafting in cleft lip and palate". *The Journal of Craniofacial Surgery* 30.3 (2019).
8. Nasrin Saki. "Pico second lasers in aesthetic dermatology". *Lasers in Dermatology Journal of Surgical Dermatology* 2 (2019): 1-3.
9. Noordhoff MS. "Reconstruction of vermilion in unilateral and bilateral cleft lips". *Plastic and Reconstructive Surgery* 73.1 (1984): 52-61.
10. The Surgical technique for the unilateral cleft lip-nasal. Deformity Noordhoff MS Taipei: Noordhoff Craniofacial Foundation (1997).

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667