

BCG Adverse Effects in Infants - Case Series

Chakradhar Maddela*

Neonatologist, Department of Neonatology, ESIC Medical College, Sanath Nagar, Hyderabad, India

***Corresponding Author:** Chakradhar Maddela, Neonatologist, Department of Neonatology, ESIC Medical College, Sanath Nagar, Hyderabad, India.

DOI: 10.31080/ASPE.2022.05.0548

Received: June 28, 2022

Published: July 27, 2022

© All rights are reserved by **Chakradhar Maddela.**

Abstract

Bacillus Calmette-Guerin (BCG) is widely using vaccination for prevention of tuberculosis in children globally. BCG adverse effects are uncommon. Common adverse effects include BCG abscess, non-suppurative and suppurative lymphadenitis and rarely BCG disseminated disease. Common therapeutic strategies for BCG complications include wait and watch policy, needle aspiration, antibiotic therapy and surgery. Correct guidelines when and how to manage BCG complications are not available. We came across four cases of BCG adverse effects recently. All four cases were noted, followed up and presenting as case series with literature review on how to treat.

Keywords: BCG; Vaccination; Complications; Management

Abbreviations

ATT: Antituberculous Treatment; BCG: Bacillus Calmette-Guerin; HCF: Health Care Facility; HCP: Health Care Personnel; LSCS: Lower Segment Caesarean Section; OPD: Outpatient Department; USS: Ultrasound Scan

Introduction

Bacillus Calmetti Gereine (BCG) is one of the most widely used vaccine for immunization in infants globally. It's use intended to protect against tuberculosis infection and its dissemination. BCG vaccine protection against tuberculosis widely varies and argued by many. Its role expected in protecting miliary tuberculosis, neuro-tuberculosis [1] and shown extended protection in children from all-cause mortality and from leprosy [2]. Commonly used BCG strains for vaccination purpose are Danish 1331, Japanese and Russian strains. The abnormal adverse BCG response with large cold abscess formation depends on technique of vaccine administration and BCG strain used.

Methods

At our health care facility outpatient clinic, we came across four cases of large BCG abscesses in infants on different days and from different places recently. All four cases were noted with case details and followed up.

Results

All four cases were infants, of which one infant was female child of 9 months old and other 3 cases were male infants.

Case 1

Nine month old female baby brought for consultation for a swelling at BCG injection site. Baby delivered at full term by spontaneous vaginal delivery at a health care facility (HCF). It was a non consanguineous marriage of young couple. Antenatal, intranatal and postnatal history was uneventful. Baby vaccinated at birth by healthcare personnel (HCP). Baby had vaccination up-to-date. Baby developed large swelling at the site of BCG injection on left arm at 1

month. The swelling was progressed and attained 3 × 3 cm and not regressed or burst to heal up even after 9 months of age. An ipsilateral axillary lymphadenitis, single, non-suppurative, measuring 2.5 × 2.5^{cm}2 was present. No adverse reactions occurred to any other vaccinations given at later date as per immunization schedule to the child. There was no history of recurrent infections or hospitalization prior to consultation. No family history of immunodeficiency or allergy. Baby was on breast feeds and weaning introduced in 6th month of life. Baby weighed 8.2kg at 9 months and her birth weight was 2.9kg. Keeping in mind the parental apprehension and unresolved abscess and local non-suppurative lymphadenitis, infant given initially oral amoxicillin-clavulanate for a week without any response. Infant responded to oral INH and Rifampicin therapy with complete resolution.

Case 2

Seventy-two days old male baby brought to our outpatient clinic for abnormal swelling at BCG injection site. Baby born to non-consanguineous young couple and delivered by lower segment caesarean section (LSCS) at a health care facility. Baby vaccinated at birth with BCG, oral polio vaccine and hepatitis B at HCF by HCP. Antenatal, intranatal and post-natal history was uneventful. There was no history of recurrent infections or hospitalization prior to consulta-

tion. Baby was on exclusive breast feeds. Swelling at BCG injection site noticed after 1 month of injection, gradually increased in size and attained present size of 3.4 × 2 cm, soft, nontender, fluctuant, well-defined borders and negative translucency test. Ultrasound scan revealed circumscribed lesion with anechoic central area and internal echogenic spots and showed no increased vascularity on colour Doppler. Baby was on follow up for wait and watch policy. The abscess was spontaneously drained after 3 days at home and later resulted in formation of large scar.

Case 3

Sixty-four days old male baby brought to our OPD for large swelling at BCG injection site on left arm by attendants. It was a full term LSCS institutional delivery to a non-consanguineously married young mother. Pregnancy was uneventful. Baby was on top feeds for first 3 days followed by on breast feeds. Baby was vaccinated with BCG injection of left arm, OPV and Hepatitis B. Baby was well and up-to-date vaccinated. BCG abscess noticed on left arm measuring 2.7 × 1.8 cm. The abscess was drained with needle and syringe by HCP at her village 3 days prior to consultation and refilled partially. Swelling was showing portal of entry into abscess with needle, margins well marked, non-tender, fluctuant and translucency test negative. Mother revealed the aspirated content was pearly white liquid.

Figure 1: Abnormal BCG response in three infants. All three cases are showing large lesions with cold abscesses. Case 1 measured 3 × 3 cm Case 2 measured 3.4 × 2.3cm and Case 3 -partially drained by local HCW and measured 2.7 × 1.8 cm.

Case 4

Four-month-old male child brought to our outpatient clinic with history of drained BCG abscess by health care personnel outside and showing scar of 2.9 × 2.6 cm on left arm towards acromion instead of towards deltoid insertion and ipsilateral, solitary, axillary lymphadenitis measuring 3.7 × 2.3 × 1.57cm and wall thickness of 1.05cm with increased peripheral vascularity on colour Doppler ultrasound scan. It is non-suppurative with no inflammatory signs

clinically. BCG was given at birth by HCP. No other significant findings in history and on examination. Child's total leukocyte count was 9,800 per cu.mm with neutrophils 49%, lymphocytes 44% and 'C' reactive protein 9.1mg/L. Child prescribed antibiotics (amoxicillin-clavulanate) in view of outside needle drainage and peripheral rim of increased vascularity of axillary lymph node on colour Doppler USS. Child is still on follow up.

Figure 2: Showing case 3 ultrasound scan image and large BCG scar and case 4 showing solitary axillary lymphadenitis (left top), BCG scar (right top), ultrasound scan images and translucency test of lymphadenitis.

Discussion

All neonatal infants receive BCG vaccine at birth in India under Universal Immunization Programme and globally in most of the countries as well. Normal BCG response includes formation of a wheal of > 5 mm immediately after the vaccination followed by papule formation by 2 weeks, further enlargement results in formation of nodule by 4 weeks. This is followed by pustule formation by 6 weeks and later self drains and results in healing with scar formation between 6-12weeks. Abnormal attainment in size with large abscess formation and local lymphadenitis can be caused by improper technique of administration of vaccine, type of vaccine strain used, age and immune status of the infant [3]. BCG lymphadenitis develops by 8weeks after injection administration [4]. Adverse reactions are more often seen with Danish strain (more effective) and least common with Russian strain. Danish strain previously widely used in India and is now replaced by Russian strain by the vaccine makers Serum Institute of India. All the four cases were vaccinated with BCG Russian strain and manufactured by the Serum institute of India. Common BCG injection complications include abscess formation at injection site (1/3 of complications),

local/axillary lymphadenitis (2/3 of complications) with or without suppuration, and BCG disease (disseminated) including osteomyelitis. These complications present usually within 6months of injection (Venkatraman A 2015).

Our case 1 and 4 had BCG abscesses and axillary lymphadenitis whereas case 2 and 3 were presented with BCG abscesses. Case 2 spontaneously ruptured and healed with large scar. Case 3 and case 4 were drained outside by HCP and were prescribed antibiotics. Case 3 partially drained outside. Only case 1 was treated with ATT and resolved completely. Case 4 had BCG scar but having non-tender, non-suppurative axillary lymphadenitis and on follow up.

Literature Review

BCG vaccination - In newborns, BCG should be given by the trained vaccinator with correct dose of 0.05ml (0.1ml in children) by intradermal route with 26G needle at insertion of deltoid on left upper arm.

Natural course of BCG response and its variants: Normal BCG response after injection was already discussed above in discussion section. BCG injection has got both therapeutic and diagnostic value. BCG reaction is said to be accelerated if an induration of > 5mm at the injection site occurs within 24 - 72 hours, hyper-accelerated if the same induration noticed between 6-12 hours and delayed response if same induration response occurs between 72hours to 2weeks [5]. These reactions may indicate latent tuberculous infection or active tuberculous infection and warrants further evaluation. BCG adverse reactions includes 1. local BCG abscess, 2. local non-suppurative lymphadenitis, 3. local suppurative lymphadenitis and 4. disseminated BCG disease [6].

BCG adverse reactions common in the following circumstances - 1. When BCG wrongly administered by subcutaneously 2. When given on upper 1/3 of arm 3. When BCG given to newborn on day one or within 1st week of life 4. When BCG vaccine vial contains less live bacteria and more killed BCG bacteria 5. When the antigen dose increased 6. Preterm infants < 37weeks gestational age experience 6times more risk of developing local lymphadenitis 7. Three times increased risk of lymphadenitis was noticed when BCG administered on right arm instead of on left arm [7].

- **BCG abscess:** Most BCG abscesses start forming before 1month after injection. They are fluctuant, nontender and without systemic features. Routine laboratory evaluation is not necessary unless a secondary bacterial infection is suspected. Most isolated BCG abscesses resolve and heal spontaneously after spontaneous rupture and needs wait and watch policy. Suspected BCG abscesses with secondary bacterial infection need laboratory work up for evidence of secondary infection and antibiotic therapy (Amoxicillin-clavulanate × 7days or Erythromycin) and needle drainage.
- **Non-suppurative BCG lymphadenitis:** Ipsilateral, axillary, solitary, non-suppurative lymphadenitis < 3CM² usually resolve spontaneously over time and needs to adopt wait and watch policy. Lymphadenitis > 3CM² may undergo suppuration (> 30% cases Goraya J S 2002) [8] and may need needle aspiration.
- **Suppurative BCG lymphadenitis:** These cases should be treated with needle aspiration – single or repeated and antibiotic therapy with either INH or INH plus Rifampicin for 3 months to prevent sinus and keloid formation and can be

managed on out patient department [9]. Needle drainage facilitates regression and prevents further complications. In rare cases, excisional biopsy is needed and should be reserved for repeated recollections after needle drainage and matted or multilocular lymphadenitis [10]. Surgical management needs hospitalization and general anaesthesia administration.

- **Disseminated BCG disease:** This condition common in immunodeficiency states like severe combined immunodeficiency (SCID), chronic granulomatous disease (CGD) and Mendilian susceptibility to mycobacterial disease (MSMD). Osteomyelitis and disseminated BCG tuberculosis need aggressive management with hospitalization, supportive therapy and complete anti tuberculosis therapy. Pyrazinamide do not work in BCG disease because usage of Bovine strain in BCG vaccine.

Key points

- **What is already known about this topic:** 1. Normal course of BCG response after its intradermal administration. 2. BCG adverse reactions are related to strain used, technique of administration, age and immune status of the child.
- **What is not known about this topic:** Less evidence is available regarding the correct approach to BCG complications. When to treat and how to treat?
- **What this paper adds:** Clinical and therapeutic approach to common BCG complications at low resource setting.

Conclusions

- Majority BCG abscesses need wait and watch policy for spontaneous resolution.
- BCG Abscess with secondary bacterial infection needs antibiotic therapy and needle drainage.
- Non-suppurative BCG lymphadenitis needs wait and watch policy for spontaneous resolution.
- Suppurative BCG lymphadenitis needs needle drainage and ATT and rarely excisional biopsy.
- Disseminated BCG disease indicates underlying immunodeficiency state.

Conflict of Interest

There are no conflicts of interest for this study.

Sponsors

There are no sponsors for this study.

ORCHID Number

0000-0001-7475-0777.

Bibliography

1. Rodrigues LC., *et al.* "Protective effect of BCG against tuberculosis meningitis and miliary tuberculosis: a meta-analysis". *International Journal of Epidemiology* 22 (1993): 1154-1158.
2. Venkatraman A., *et al.* "Management and outcome of Bacille Calmette - Guerin vaccine adverse reactions". *Vaccine* 33 (2016): 5470-5474.
3. Al-Mousa Hamond., *et al.* "An infant with disseminated bacillus Calmette - Buerin infection (BCGitis)". *International Journal of Pediatrics and Adolescent Medicine* 1 (2014): 89-92.
4. Alfawaz TS., *et al.* "BCG related complications: A single centre prospective observational study". *International Journal of Pediatrics and Adolescent Medicine* 2.2 (2015): 75-78.
5. Vikkanueva P., *et al.* "What is the significance of an accelerated BCG reaction in children?" *Archives of Disease in Childhood* 107.5 (2022).
6. Nissen TN., *et al.* "Adverse reactions to the Bacillus Calmette-Guerin (BCG) vaccine in new-born infants - an evaluation of the Danish strain 1331 SSI in a randomized clinical trial". *Vaccine* 34 (2016): 2477-2482.
7. Mostaan S., *et al.* "Adverse effects of BCG vaccine 1173 P2 in Iran: A meta-analysis". *Advanced Biomedical Research* 5 (2016): 99.
8. Goraya JS and Viridi VS. "Bacille Calmette-Guerin lymphadenitis". *Postgraduate Medical Journal* 78.920 (2002): 327-329.
9. Aoki Y., *et al.* "Unusual wound healing following suppurative BCG lymphadenitis". *BMJ Case Reports* (2018).
10. Banani SA and Alborzi A. "Needle aspiration for suppuration post-BCG adenitis". *Archives of Disease in Childhood* 71 (1994): 446-447.