



Broad Antibacterial Spectrum and High Performance Liquid Chromatography Profiles of *Ocimum sanctum* Leaf Extract

Malay Kr Das¹, Manisha Mandal² and Shyamapada Mandal^{1*}

¹Department of Zoology, University of Gour Banga, Malda, West Bengal, India

²Department of Physiology, MGM Medical College and LSK Hospital, Kishanganj, India

*Corresponding Author: Shyamapada Mandal, Laboratory of Microbiology and Experimental Medicine, Department of Zoology, University of Gour Banga, Malda, West Bengal, India. E-mail: samtropmed@gmail.com

Received: September 23, 2017; Published: December 06, 2017

Abstract

Medicinal plants are the promising source of bioactive compounds which are well known for their several biological activities including antibacterial property. The current study assesses the antibacterial activity of two indigenous varieties (Dark and Bright) of *Ocimum sanctum* leaf extracts against gram-positive and gram-negative clinical bacteria, and determines the HPLC profiles of the extracts. Following disc diffusion method, the antibacterial activity of ethanolic leaf extracts (DOSE and BOSE) and aqueous leaf extracts (AqDOS and AqBOS) of both varieties *O. sanctum*, was determined and their phytochemical analysis was performed by HPLC. The HPLC chromatogram showed the presence of 11, in DOSE, and 9, in BOSE, major compounds with the RTs 1.77 - 5.29 min and 1.92 - 6.27 min, respectively, while both AqDOS and AqBOS had 4 major compounds, with the respective RTs of 1.66 - 6.17 min and 1.64 - 2.51 min. The ZDI values, due to the action of DOSE (6.25 mg), were recorded as 20 mm, for gram-positive bacteria, and 20 - 24 mm, for gram-negative bacteria, while BOSE (6.25 mg) had ZDIs 17 - 19 mm and 14 - 24 mm, respectively, for gram-positive and gram-negative bacteria. The DOSE and BOSE were more active compared to the AqDOS and AqBOS, in terms of the antibacterial activity by disc diffusion. Thus, the *O. sanctum* leaves might be utilized as the source of biotherapeutic agents to be administered against broad range of bacterial infections.

Keywords: *Ocimum sanctum*; Broad Spectrum Antibacterial Activity; Zone Diameter of Inhibition; Phytocomponents; High Performance Liquid Chromatography

Abbreviations

AqBOS: Aqueous Leaf Extract of *O. sanctum* Bright Variety; AqDOS: Aqueous Leaf Extract of *O. sanctum* Dark Variety; BOSE: Ethanolic Leaf Extract of *O. sanctum* Bright Variety; DOSE: Ethanolic Leaf Extract of *O. sanctum* Dark Variety; HPLC: High Performance Liquid Chromatography; MDR: Multi Drug Resistant; MTCC: Microbial Type Culture Collection; RT: Retention Time; SD: Standard Deviation; ZDI: Zone Diameter of Inhibition

Introduction

The bacterial pathogens, such as *Staphylococcus aureus* (*S. aureus*), *Acinetobacter baumannii* (*A. baumannii*), *Klebsiella pneumonia* (*K. pneumoniae*), and *Pseudomonas aeruginosa* (*P. aeruginosa*) are the important cause of nosocomial infections [1]. The antibiotic resistances among such human pathogenic bacteria have been reported worldwide [2-5]. In developing countries, including India, the burden of the infectious diseases has been recorded high because of the emergence of MDR bacterial pathogens resulting from the poor health-care facilities, and over-the-counter use of antimicrobial agents [6]. This fact has directed the clinicians and the scientists around the world to search for novel affordable nontoxic antimicrobials, alternative to the synthetic antibiotics, with promising therapeutic potential to overcome the situation of MDR bacterial infections.

Various parts of different plants have been shown in several studies to be one of the most promising sources for obtaining antimicrobial compounds [7-9]. Among different medicinal plants, *Ocimum sanctum* L. (= *Ocimum tenuiflorum*) is a unique one, and is known as tulsi in Bengali: the holy basil, belonging to the fam-

ily Lamiaceae [10]. It is an herbaceous aromatic plant that grows abundantly in the tropical and subtropical regions of the Indian subcontinents, and widespread as a cultivated plant throughout the world, due to its great therapeutic importance to cure various infectious and non-infectious diseases [11,12]. There are commonly two varieties of *O. sanctum* found in India, which include Dark variety or the 'Krishna tulsi' and 'Bright variety or the 'Rama tulsi' [12-14]. The earlier authors from different parts of the globe documented the presence of various bioactive compounds in *O. sanctum* leaf extracts by HPLC analysis [15-17], and validated their antibacterial efficacy against gram-negative as well as gram-positive pathogenic bacteria [18-21].

In our part of the globe, though both varieties (Dark and Bright) of *O. sanctum* are found enormously, the extensive scientific studies remain to be explored in terms of the antibacterial activity or phytochemical analysis of the plant. Therefore, the above mentioned background prompted us to explore the broad spectrum antibacterial activity of *O. sanctum* leaf extracts along with HPLC profiles of phytochemical contained in them, in order to validate the traditional usage of *O. sanctum* leaves against bacterial infection to humans.

Materials and Methods

Bacterial strains

The gram-positive, such as *S. aureus* (n = 2), and gram-negative, such as *A. baumannii*, *K. pneumonia* and *P. aeruginosa* (n = 2), clinical bacteria were utilized as the indicator strains in testing the antibacterial activity of *O. sanctum* leaf extracts.

Preparation of *O. sanctum* leaf extracts

The ethanolic and aqueous extracts of *O. sanctum* leaves were prepared for the present study. The collection, processing and preparation of ethanolic and aqueous *O. sanctum* leaf extracts have been described in the previous publication [11]. The concentrations of the ethanolic (DOSE and BOSE) and aqueous (AqDOS and AqBOS) leaf extracts were 125 µg/µl.

Bacterial growth inhibition property of *O. sanctum* leaf extracts

The bacterial growth inhibition property of the plant extracts were assessed by disk diffusion method [22], and the details of the method have been mentioned elsewhere [2]. Briefly, two different concentrations (3.75 and 6.25 mg) of each, of the prepared extracts, were dropped on sterile blank paper disc, placed on the surface of the already inoculated nutrient agar plates, with test bacterial fresh broth culture, by swabbing. The sensitivity of the test bacterial isolates, to the plant extracts after incubation for 24h at 37°C, was considered with ZDIs ≥ 7 mm, obtained around the discs loaded with the extracts [23].

HPLC analysis

The HPLC analysis of the ethanolic (DOSE and BOSE) and aqueous (AqDOS and AqBOS) *O. sanctum* leaf extracts was done in the YL 9100 HPLC system associated with C₁₈ column (5 µm; 100Å; 4.6 × 250 mm). The sample volume used for injection into the column was 10 µl, and flow rate was 1.0 ml/min. The mobile phase used in this system comprised of acetonitrile and water (2:3 ratio). The detection of the eluting compounds was done at 230 nm, at 35°C.

Statistical analysis

The results were expressed as the mean ± SD, and evaluated by 't' test in comparing the data (ZDI values from the action of *O. sanctum* leaf extracts), using MS Excel 2010 software; a 'p' value of ≤ 0.05 was considered for statistical significance.

Results

The antibacterial activity of *O. sanctum* leaf ethanolic (DOSE and BOSE) and aqueous (AqDOS and AqBOS) extracts, following disc diffusion technique, is depicted in Figure 1.




Figure 1: Antibacterial activity of *O. sanctum* leaf extracts; A-C: dark variety ethanolic extract; D-F: bright variety ethanolic extract. a: 3.75 mg; b: 6.25 mg. The clear halos around each disc on the plates are indicative of growth inhibitory action of the extracts.

The ZDI values due to the action of ethanolic and aqueous extracts of *O. sanctum* leaf are represented in Table 1. The ZDIs of DOSE and BOSE, at the concentration of 3.75 mg, for the gram-positive bacteria ranged 17 - 19 mm and 11 - 17 mm, respectively, while for the gram-negative bacteria the ZDIs ranged 14 - 20 mm and 9 - 15 mm, respectively. At the concentration of 6.25 mg the ZDIs of DOSE and BOSE for the gram-positive bacteria were 20 mm and 17 - 19 mm, respectively, and for the gram-negative bacteria, the ZDIs ranged 20 - 24 mm and 14 - 24 mm, respectively. The overall ZDIs recorded were 17.5 ± 2.07 mm and 21 ± 1.67 mm for DOSE (at 3.5 and 6.25 mg, respectively), and 12.5 ± 2.94 mm and 18.67 ± 3.33


mm for BOSE (at 3.5 and 6.25 mg, respectively); the AqDOS had no activity against the test bacteria, while AqBOS showed activity against *A. baumannii* (ZDIs: 9 - 11 mm), only (Table 1).

The HPLC chromatograms revealed the presence of 11 and 9 major compounds, respectively, for DOSE and BOSE (Figure 2), and in cases of AqDOS and AqBOS, 4 major compounds were displayed (Figure 3). The retention times (RTs) within which the compounds were detected ranged 1.77 - 5.29 min and 1.92 - 6.27 min, respectively, for DOSE and BOSE, and 1.66 - 6.17 min and 1.64 - 2.51 min, respectively, for AqDOS and AqBOS.


Bacterial strain	ZDI (mm) at different extract concentration (mg)							
	3.75		6.25		3.75		6.25	
	DOSE*	BOSE**	DOSE ^α	BOSE ^{αα}	AqDOS*	AqBOS**	AqDOS ^α	AqBOS ^{αα}
<i>S. aureus</i> ZM1	17	17	20	19	6	6	7	6
<i>S. aureus</i> ZM2	19	11	20	17	6	6	6	6
<i>A. baumannii</i> ZM1	14	9	24	14	6	9	6	11
<i>K. pneumonia</i> ZM1	17	15	20	24	6	6	6	6
<i>P. aeruginosa</i> ZM1	18	11	20	18	6	6	6	6
<i>P. aeruginosa</i> ZM2	20	12	22	20	6	6	6	6
Mean value	17.5	12.5	21	18.67	6	6.5	6.17	6.83
SD	±2.07	±2.94	±1.67	±3.33	0	±1.23	±0.41	±2.04
P value	0.003		0.083		0.181		0.232	

Table 1: The ZDI (Zone diameter of inhibition) values due to the action of *O. sanctum* leaf extracts at two different concentrations against pathogenic bacteria

DOSE: Ethanolic Leaf Extract of *O. sanctum*, Dark Variety; BOSE: Ethanolic Leaf Extract of *O. sanctum*, Bright Variety; AqDOS: Aqueous Leaf Extract of *O. sanctum*, Dark Variety; AqBOS: Aqueous Leaf Extract of *O. sanctum*, Bright Variety. *p value: <0.01; ** p value: < 0.01; α p value: 0.01; αα p value: < 0.01.




DOSE




BOSE

Figure 2: The HPLC chromatograms of *O. sanctum* ethanolic extracts. DOSE: ethanolic leaf extract of *O. sanctum*, dark variety; BOSE: ethanolic leaf extract of *O. sanctum*, bright variety.



AqDOS



AqBOS

Figure 3: The HPLC chromatograms of *O. sanctum* aqueous extracts. AqDOS: aqueous leaf extract of *O. sanctum*, dark variety; AqBOS: aqueous leaf extract of *O. sanctum*, bright variety.

Discussion

The *O. sanctum* is regarded as the queen of herbs and has been used since the antiquity in traditional medicinal system to diverse range of therapeutic activities, for alleviating many diseases including bacterial infections [10,24]. The antibacterial activity of *O. sanctum* has been reported by earlier authors. Geeta *et al.* [25] reported the growth inhibitory activity of *O. sanctum* alcoholic leaf extract against *Vibrio cholerae*. Antibacterial efficacy of *O. sanctum* ethanolic leaf extract against gram-positive (*S. aureus* and *Bacillus cereus*) and gram-negative (*Escherichia coli*, *Proteus vulgaris*, *Salmonella typhi*, *K. pneumonia*, *P. aeruginosa* and *A. baumannii*) bacteria has been reported by Das *et al.* [11]. Auil *et al.* [26] reported the antibacterial activity of *O. sanctum* leaf extract against *S. aureus* having resistance to β -lactams. As per the report of Singh *et al.* [27], due to higher content of linolenic acid, the *O. sanctum* fixed oil was found to be active against *Bacillus pumilus*, *P. aeruginosa* and *S. aureus*. In the current study, ethanolic leaf extract of both varieties of *O. sanctum*, showed concentration dependant antibacterial activity against all the gram-positive and gram-negative bacteria tested, while most of the bacterial isolates had resistance (ZDI = 6 mm) to the aqueous extracts: AqDOS and AqBOS. A significant difference ($P < 0.01$) was observed in between the antibacterial activity of ethanolic and aqueous extracts of both varieties of *O. sanctum* leaf. At 3.75 mg extract concentration, the ZDIs of DOSE recorded were 17 - 19 mm and 14 - 20 mm, respectively, for the gram-positive and gram-negative bacteria tested; at the same concentration, BOSE displayed respective ZDIs of 11 - 17 mm and 9 - 15 mm, for the gram-positive and gram-negative test bacteria. The ZDI values, due to the action of DOSE (at 6.25 mg) were recorded as 20 mm, for gram-positive bacteria, and 20 - 24 mm, for gram-negative bacteria, while BOSE at the same concentration had ZDIs 17 - 19 mm and 14 - 24 mm, re-

spectively, for the gram-positive and gram-negative test bacteria. Therefore, except at 3.75 mg concentration ($p: 0.003$), there was no significant differences in antibacterial action between DOSE and BOSE, and between AqDOS and AqBOS ($p: 0.083 - 0.232$). Similar to the current study, Sadul *et al.* [28], demonstrated the antimicrobial activity of *O. sanctum* leaf extract, and found ethanolic extract more efficacious than the aqueous *O. sanctum* leaf extracts.

The HPLC is an important tool that is utilized in the identification, quantification and purification of the individual components from the mixture of samples [29]. Several authors from different parts of the world using HPLC have identified medicinally important bioactive compounds from *O. sanctum* leaf extracts. Among the different biologically active components present in *O. sanctum* leaf, eugenol is a major one, which is most widely used as flavouring agents in foods and beverages; it has been detected, in HPLC system, at different RTs: 11.96 - 14.41 min, using methanol-water-acetonitrile mobile phases [16,30,31]. Tamilselvan *et al.* [32] identified three phenolic components (RT ranged: 3.33 - 12.57 min) in *O. gratissimum* chloroform extract using acetonitrile and water (60:40) as the mobile phase. In HPLC system, with the mobile phase of 10 mM hexane -1- sulphonic acid sodium salt containing 1% acetic acid and 0.13 % triethyl amine, and a flow rate of 1.0 ml/min, the presence of antibacterial components: ursolic acid and oleanolic acid, in *O. sanctum* leaf extracts at respective RT of 14.06 min and 14.43 min has been reported by Ali *et al.* [33]. Herein, the HPLC chromatograms demonstrated the presence of 11 and 9 major compounds, respectively in DOSE and BOSE, and 4 major compounds both in AqDOS and AqBOS. The RTs for each of the test extract ranged 1.77 - 5.29 min and 1.92 - 6.27 min, respectively for DOSE and BOSE, whereas for the AqDOS and AqBOS

the RTs ranged 1.66 - 6.17 min and 1.64 - 2.51 min, respectively, in detecting the compounds. Similar to the present study, Deo *et al.* [17] found the presence of 10 compounds (RTs: 1.78 - 11.55 min) in methanolic extract and 3 compounds (RTs: 1.72 - 3.45 min) in aqueous extract of *O. sanctum* leaf. Shanaida *et al.* [34] detected various amino acids: aspartic acid (RT: 1.92 min), asparagine (RT: 2.89 min), serine (RT: 3.12 min), arginine (RT: 3.28 min), alanine (RT: 3.62 min), proline (RT: 3.71 min) and isoleucine (RT: 4.81 min), in *O. americanum* seed; the RTs were closely similar to the compounds detected in DOSE and BOSE, in HPLC system. The HPLC analysis of *O. basilicum* methanolic leaf extract revealed the presence of seven phenolic components such as gallic acid (RT: 1.84 min), protocatechuic acid (RT: 2.43 min), caffeic acid (RT: 4.20 min), rutin (RT: 6.04 min), p-coumaric acid (RT: 6.66 min) and rosmarinic acid (RT: 10.53); among such components, antibacterial potentiality of caffeic acid and rosmarinic acid, against different gram-positive and gram-negative bacterial isolates, had been recorded [35]. Shafqatullah *et al.* [15] analysed the ethanolic *O. sanctum* leaf extract by HPLC, and detected 10 compounds at 210 nm, of which three compounds were identified as gallic acid (RT: 2.61 min), chlorogenic acid (RT: 6.40 min) and vanillic acid (RT: 21.02 min). Depending on the earlier findings [15,34,35], the 7 plausible components detected in DOSE were: gallic acid (compound 1), aspartic acid (compound 2), asparagine (compound 6), serine (compound 7), arginine (compound 8), proline (compound 10) and caffeic acid (compound 11), while in BOSE, the possible detected components included aspartic acid (compound 1), protocatechuic acid (compound 2), gallic acid (compound 3), arginine (compound 5), alanine (compound 6), isoleucine (compound 8) and chlorogenic acid (compound 9). In the aqueous *O. sanctum* leaf extracts gallic acid (compound 1), aspartic acid (compound 2) and chlorogenic acid (compound 4) were detected (as in AqDOS), and gallic acid (compound 1), aspartic acid (compound 2) and protocatechuic acid (compound 3) (as in AqBOS). The phytochemicals, detected in the *O. sanctum* leaf extracts, might act alone or in combination in exerting growth inhibition activity against test pathogenic gram-positive and gram-negative bacteria.

Conclusion

The current study authenticated the presence of an array of phytochemicals in *O. sanctum* leaf ethanolic and aqueous extracts, attributing the excellence of the extracts in bacterial growth inhibition activity. The plant might be utilised in the preparation of non-antibiotic biotherapeutic agents in order to combat the life-threatening illnesses, caused due to the infection of different gram-positive and gram-negative pathogenic bacteria (at least in our part of the globe), thus having broad spectrum of antibacterial activity.

However, further studies are warranted in reverence to the dose determination, and safety and toxicity profiling of the *O. sanctum* plant (not studied before) available in the local niches.

Acknowledgement

The corresponding author gratefully acknowledges the help of Dr. MA Mondal, Assistant Professor of Chemistry, University of Gour Banga, Malda, in determining the HPLC chromatography of the plant studied.

Bibliography

1. Khan HA., *et al.* "Nosocomial infections and their control strategies". *Asian Pacific Journal of Tropical Biomedicine* 5.7 (2015): 509-514.

2. Das MK., *et al.* "Syzygium cumini and Mangifera indica seed extracts: in vitro assessment for antibacterial activity alone and in combination with antibiotics against clinical bacteria". *Journal of Infectious Diseases and Preventive Medicine* 4.1 (2016): 129.
3. Cunha MA., *et al.* "Antibiotic resistance patterns of urinary tract infections in a Northeastern Brazilian capital". *Revista do Instituto de Medicina Tropical de Sao Paulo* 58 (2016): 2.
4. Muhammad SM., *et al.* "Antimicrobial susceptibility of *Acinetobacter* clinical isolates and emerging antibiogram trends for nosocomial infection management". *Revista da Sociedade Brasileira de Medicina Tropical* 49.3 (2016): 300-304.
5. Sircar B., *et al.* "High performance liquid chromatography analysis and anti-methicillin resistant *Staphylococcus aureus* activity of Olive fruit ethanolic extract". *International Research Journal Pharmacy* 8.7 (2017): 1-5.
6. Farooqui A., *et al.* "Synergistic antimicrobial activity of *Camellia sinensis* and *Juglans regia* against multidrug-resistant bacteria". *Plos One* 10.2 (2015): e0118431.
7. Boakye YD., *et al.* "Anti-infective properties and time-kill kinetics of *Phyllanthus muellerianus* and its major constituent, Geraniin". *Medicinal Chemistry* 6.2 (2016): 95-104.
8. Sircar B., *et al.* "Antibacterial activity of *Mimusops elengi* leaf, seed and bark extracts alone and in combination with antibiotics against human pathogenic bacteria". *Translational Medicine* 6.4 (2016): 188.
9. Mandal S., *et al.* "Antibacterial potential of *Azadirachta indica* seed and *Bacopa monniera* leaf extracts against multidrug resistant *Salmonella enterica* serovar Typhi Isolates". *Archives of Medical Science* 3.1 (2007): 14-18.
10. Luthra D. "Ocimum sanctum (Tulsi): A potent medicinal herb". *WebmedCentral* 1.11 (2010): 1-12.
11. Das MK., *et al.* "Assessment of bacterial growth inhibition property and phytochemical analysis of *Ocimum sanctum* L. leaf extract". *International Research Journal of Pharmacy* 8.7 (2017): 1-6.
12. Kumar PK., *et al.* "Pharmacological actions of *Ocimum sanctum*- review article". *International Journal of Advances in Pharmacy Biology and Chemistry* 1.3 (2012): 406-414.
13. Shahedur R., *et al.* "*Ocimum sanctum* L.: A review of phytochemical and pharmacological profile". *American Journal of Drug Discovery and Development* (2011).
14. Bhatt MK., *et al.* "Evaluation of anti-microbial activity of *Ocimum sanctum* methanolic extract". *Journal of Pharmaceutical and Scientific Innovation* 1.4 (2012): 39-41.
15. Shafqatullah., *et al.* "Development of HPLC method by UV-VIS detection for the quantification of phenolic acids in different *Ocimum sanctum* Linn. extracts". *Pakistan Journal of Pharmaceutical Science* 27.5 (2014): 1271-1275.
16. Joshi A., *et al.* "Analytical method development for determination of eugenol concentrations in selected species of *Ocimum*". *World Journal of Pharmacy and Pharmaceutical Sciences* 3.10 (2014): 1358-1368.
17. Deo SS., *et al.* "Antimicrobial activity and HPLC fingerprinting of crude *Ocimum* extracts". *E-Journal of Chemistry* 8.3 (2011): 1430-1437.

18. Naik LS., et al. "Antimicrobial activity and phytochemical analysis of *Ocimum tenuiflorum* leaf extract". *International Journal of PharmTech Research* 8.1 (2015): 88-95.
19. Mandal S., et al. "Enhancing chloramphenicol and trimethoprim *in vitro* activity by *Ocimum sanctum* Linn. (Lamiaceae) leaf extract against *Salmonella enterica* serovar Typhi". *Asian Pacific Journal of Tropical Medicine* 5.3 (2012): 220-224.
20. Mishra P., et al. "Study of antibacterial activity of *Ocimum sanctum* extract against gram positive and gram negative bacteria". *American Journal of Food Technology* 6.4 (2011): 336-341.
21. Rahman MS., et al. "Anti-bacterial evaluation and minimum inhibitory concentration analysis of *Oxalis corniculata* and *Ocimum sanctum* against bacterial pathogens". *Biotechnology* 9.4 (2010): 533-536.
22. Bauer AW., et al. "Antibiotic susceptibility testing by a standardized single disc method". *American Journal of Clinical Pathology* 45.4 (1996): 493-496.
23. Mandal S., et al. "Synergistic anti-*Staphylococcus aureus* activity of Amoxicillin in combination with *Embllica officinalis* and *Nymphae odorata* extracts". *Asian Pacific Journal of Tropical Medicine* 3.9 (2010): 711-714.
24. Tewari D., et al. "A review on phytoconstituents of *Ocimum* (Tulsi)". *International Journal of Ayurvedic Medicine* 3.1 (2012): 1-9.
25. Geeta DMV., et al. "Activity of *Ocimum sanctum* (the traditional medicinal plant) against the enteric pathogens". *Indian Journal Medical Science* 55.8 (2001): 434-438.
26. Auil F., et al. "Effect of certain bioactive plant extracts on clinical isolates of beta lactamase producing Methicillin resistant *Staphylococcus aureus*". *Journal of Basic Microbiology* 45.2 (2005): 106-114.
27. Singh S., et al. "Antibacterial activity of *Ocimum sanctum* L. fixed oil". *Indian Journal of Experimental Biology* 43.9 (2005): 835-837.
28. Sadul RR., et al. "Comparative study of antimicrobial activity and phytochemical screening of aqueous and alcoholic leaf extract of *Ocimum sanctum* on *E. coli* (faecal indicator of water pollution)". *Journal of Environmental Research and Development* 7.1 (2012): 312-320.
29. Malviya R., et al. "High performance liquid chromatography: a short review". *Journal of Global Pharma Technology* 2.5 (2010): 22-26.
30. Joshi A., et al. "Development of an analytical method to determine eugenol concentrations in alcoholic extracts of different species of *Ocimum*'s". *International Journal of Phytopharmacy* 1.2 (2011): 35-42.
31. Inam F., et al. "HPLC -UV method development and quantification of eugenol from methanolic extracts of some spices". *International Journal of Chemical and Physical Sciences* 3.6 (2014): 96-102.
32. Tamilselvan P., et al. "Larvicidal, pupicidal and adulticidal potential of *Ocimum gratissimum* plant leaf extracts against filariasis inducing vector". *International Journal of Mosquito Research* 2.2 (2015): 1-8.
33. Ali R., et al. "In-vitro analysis of antibacterial activity of *Ocimum sanctum* against pathogenic bacteria and quantification of ursolic acid and oleanolic acid". *International Journal of Pharmaceutical Sciences Review and Research* 25.2 (2014): 13-17.
34. Shanaida M., et al. "Chromatographic analysis of organic acids, amino acids, and sugars in *Ocimum americanum* L". *Acta Poloniae Pharmaceutica - Drug Research* 74.2 (2017): 729-734.
35. Khattab OKH., et al. "Biological activity of extracts from Olive and Basil leaves against pathogenic microbial isolates". *Egyptian Journal of Medical Microbiology* 24.2 (2015): 1-9.

Volume 1 Issue 6 December 2017

© All rights are reserved by Shyamapada Mandal, et al.