



Effect of O⁶-Methylguanine-DNA Methyltransferase Resistance to Temozolomide in Gliomas or Brain Metastases of Melanoma

Emad Y Moawad*

Researcher Graduated, Department of Engineering, Ain Shams University, Cairo, Egypt

*Corresponding Author: Emad Y Moawad, Researcher Graduated, Department of Engineering, Ain Shams University, Cairo, Egypt.

Received: July 19, 2018; Published: August 02, 2018

Abstract

The aim of this study is to investigate the resistance of O⁶-methylguanine-DNA methyltransferase (MGMT) in gliomas or brain metastases of melanoma to Temozolomide (TMZ) therapy. The in-vivo effects of standard, metronomic and dose-dense TMZ regimens in animal models were monitored to identify efficacy of those regimens with respect to suppression of MGMT-resistance in tumor cells. MGMT-resistance is dependent mainly on the TMZ dose/day of the applied regimen, observed when applying standard regimens of 2.45 mg/kg bw/day for 5 days every 28 days and maximized by increasing dose/day to 6.74 mg/kg bw/day for 5 days every 28 days. Afterwards, MGMT-resistance decreased gradually until being suppressed by increasing the received dose/day in standard regimen to 8.88 mg/kg bw/day for 5 days every 28 days. Standard regimen of dose less than 4.37 mg/kg bw/day for 5 days every 28 days is more efficient than the metronomic one of dose/day less than 0.78 mg/kg bw/day for 28 days in early stages of primary tumors. While the metronomic regimen of dose/day lies between 0.78 and 1.6 mg/kg bw/day for 28 days is more efficient than the standard one of administered dose lies between 4.37 and 8.96 mg/kg bw/day for 5 days every 28 days in the moderate stages of recurrent tumors of higher MGMT-resistance. Dose-dense regimens with standard schedule of dose/day higher than 8.96 mg/kg bw/day for 5 days every 28 days or metronomic schedule of dose/day higher than 1.6 mg/kg bw/day for 28 days suppress MGMT-mediated resistance in advanced stages of high-grade tumors by depleting MGMT in tumor cells.

Keywords: Temozolomide; O⁶-Methylguanine-DNA Methyltransferase; Gliomas; Brain Metastases of Melanoma

Introduction

Temozolomide (TMZ) is an imidazotetrazine derivative of the oral alkylating agent dacarbazine with established antitumor activity in patients with primary and recurrent brain tumors and brain metastases of melanoma [1]. It undergoes rapid chemical conversion in the systemic circulation at physiological pH to the active compound, MTIC (monomethyl triazeno imidazole carboxamide) [2]. In a recent randomized trial, concomitant and adjuvant TMZ chemotherapy with radiation significantly improves, from 12.1 months to 14.6 months, progression free survival and overall survival in glioblastoma Multiforme (GBM) patients [3]. The therapeutic benefit of TMZ depends on its ability to alkylate/methylate DNA, which most often occurs at the N⁷ or O⁶ positions of guanine residues. This methylation damages the DNA and triggers the death of tumor cells [4]. Unfortunately, TMZ is not always effective in patients with gliomas; in some cases tumor cells are able to repair this type of DNA damage, and therefore diminish the therapeutic efficacy of TMZ due to an enzyme called O⁶-methylguanine-DNA methyltransferase (MGMT) or O⁶-alkylguanine-DNA alkyltransferase (AGT or AGAT) [5-7]. While in other cases of responding tu-

mors to TMZ therapy, epigenetic silencing of the MGMT/AGT gene prevents the synthesis of this enzyme, and as a consequence such tumors are more sensitive to killing by TMZ [8]. Thus, the increase of MGMT in brain tumors during TMZ therapy predicts poor response and these patients receive little benefit from chemotherapy with TMZ [9]. Such unsatisfactory outcome of chemotherapy with TMZ is mainly defined by the intrinsic or acquired MGMT-resistance repairing DNA of tumor cells. Several trails have been conducted to overcome TMZ resistance acquired by recurrent GBM or High Grade Glioma (HGG) including reversed approaches and the use of therapeutic agents to suppress the activity of MGMT in tumor cells. However those trails had adverse effects as impaired hepatic and renal function or myelosuppression [10,11]. Since TMZ exhibits schedule-dependent antineoplastic activity by interfering with DNA replication, then a better understanding of the relationship between the MGMT-resistance and the efficacy of the applied TMZ regimen could help to design a new therapeutic regimen to overcome MGMT-resistance to TMZ. The originally approved TMZ dosing regimen is 150 to 200 mg/m² per day on days 1 to 5 of each 28-day cycle (5 of 28 days) as a clinically active and generally

well-tolerated regimen [12-17]. Tolcher, *et al.* showed that MGMT enzyme activity consumed during treatment with TMZ for either 7 consecutive days every 14 days (7 of 14-day regimen) or 21 consecutive days every 28 days (21 of 28-day regimen). These extended dosing regimens allow for administration of a higher cumulative dose per cycle and have been shown to deplete O⁶-methylguanine-DNA methyltransferase, which may enhance cytotoxic activity [18]. Thus, these schedules of dose-dense regimens that deplete MGMT were suggested to improve antitumor activity against MGMT-resistance in tumor cells. On the other hand, the antitumor activity of low dose metronomic (LDM) regimens in the TMZ therapy has gradually been elucidated by preclinical and clinical research suggesting that the resistance of recurrent tumors to TMZ can be overcome by synchronizing metronomic TMZ regimen with MGMT resistance. However, the extent of MGMT depletion in tumor cells in each type of the proposed regimens has not been directly demonstrated yet. Current thesis identifies the dosing limits of TMZ to overcome MGMT-resistance, investigates the efficacy of those regimens and recommendations to all stages of primary or recurrent tumors for optimizing therapy.

Materials and Methods

Standard and LDM TMZ regimens in rats bearing C6/LacZ rat glioma cells and mice bearing U-87MG human glioblastoma cells

As conducted and described by Kim, *et al.* [19]; Rat glioma cell line, C6/LacZ, and U-87MG human glioblastoma cells (1×10^5 cells/10 μ l of Hanks balanced salt solution) were injected at a depth of 5 mm for male Sprague-Dawley (SD) rats (200 - 250g) and male Balb/c-nu mice (6 weeks) respectively, through a 10- μ l Hamilton syringe connected to the manipulating arm of the produced 7-stereotatic device. All injections consisted of a total volume of 10 μ l delivered over 12.5 min by a micro infusion pump. Before treatment of TMZ, the rats were randomized into four groups (n = 10 per group). One experimental group of rats was administered orally with 7 mg/kg TMZ for 5 days (between day 7 and 11 after intracranial implantation) to represent a tolerated synchronized standard regimen for TMZ therapy. Two groups of rats were respectively administered 1 or 2 mg/kg TMZ via per os (p.o.) everyday for 16 days to represent a tolerated LDM regimen for TMZ therapy. The control group of rats was treated by injection via p.o. with 10% DMSO. Human U-87MG at nude mice brain is so sensitive to TMZ then the starting dose of TMZ was minimized to show the antitumor activity. The mice were randomized into five groups (n = 15 per group). Two conventionally-treated groups of mice were administered orally with 2.5 or 1.25 mg/kg TMZ for 5 days, respectively (between day 14 and 18 after intracranial implantation) to represent a tolerated standard regimen for TMZ therapy. Two groups of mice were treated orally with each dose of TMZ (0.5 and 0.25 mg/kg) daily for 25 days to

represent a tolerated LDM regimen for TMZ therapy. The control group of mice was treated by p.o. with 10% DMSO. Seventeen and twenty five days after the inoculation of tumor cells, rats and mice were anesthetized and sacrificed respectively. The tumor volume was calculated by measuring the section with the largest tumor portion and applying the formula: Length \times Width² \times 0.5.

As conducted and described by Son, *et al.* [20]; briefly, male SD rats (200 - 250g) were anesthetized and shaved. They were secured in a rodent stereotactic frame, and a hollow guide screw was implanted into a small drill hole made 3 mm left lateral and 1 mm anterior to the bregma [21]. C6/LacZ rat glioma cells (1×10^5 cells/10 μ l) were injected through this guide screw into the white matter at a depth of 5 mm. TMZ (7.5 mg/kg) was administered intraperitoneally for 5 days, between the 7th and 11th day after tumor cell inoculation to represent a synchronized tolerated standard regimen for TMZ therapy.

Dose-dense regimen in mice bearing human melanoma UACC903 tumor model

As conducted and described by Zhang L., *et al.* [22], Melanoma mouse model: Nude mice (swissnu/nu) were inoculated s.c. with UACC903 human melanoma cells (1×10^6 cells/100 μ l/mouse). When the tumors reached 100 mm³ in volume, the mice were treated i.p. with TMZ (15 mg/kg) on days 1, 3, 5, 7 and 9 to represent a dose-dense regimen for TMZ therapy. The control group of mice was treated by p.o. with 10% DMSO. Tumor sizes and body weight of the animals were measured every other day. The differences between treated and control groups were analyzed using a two-sample t-test. The survival curves of the treated tumor-bearing mice were estimated using Kaplan-Meier method and compared by log-rank statistic analysis.

Monitoring the mechanical behavior of the tumor response to therapy

Comparing the mechanical behavior of tumor response of the treated groups by that of the control groups is assessed by determining the growth/or shrinkage constants of those tumors of different volumes along the corresponding periods [23,24]. The tumor growth/or shrinkage constant at a certain time expresses the rate of the difference between Mitosis and Apoptosis with respect to the total number of the tumor cells (M - A) that characterize the tumor response at that time [23-27]. If rate of mitosis is greater than that of apoptosis, tumor grows by growth constant of $\frac{\ln 2}{t_D}$ and vice versa if rate of mitosis was less than that of apoptosis, tumor shrinks by shrinkage constant of $-\frac{\ln 2}{t_{1/2}}$ where $t_{1/2}$ is the tumor half-life time [23,27].

i.e. $(M - A) = \frac{\ln 2}{t_D} S^{-1}$ in case of tumor growth,

and

$(A - M) = \frac{\ln 2}{t_{1/2}} S^{-1}$ in case of tumor shrinkage,

where t_D and $t_{1/2}$ in seconds Eq (1).

The clinical staging model presented by Moawad showed that the tumor histologic grade (H_G) that expresses tumor response can be identified by using Emad formula [28-46] as follows:

In case of tumor growth

$$H_G = \ln \left(\ln \frac{\ln 2}{t_D} \right)^2 \times C_0 \times h \times 23234.59 \text{ MeV} \text{ Eq (2)},$$

where $C_0 \times h$ is number of the hypoxic cells in the tumor or number of the inoculated cells in the transplanted tumor in xenografted models [25-40].

In case of tumor shrinkage

The chemotherapeutic drugs affect the tumor cells such that the more the drug dose the less of mitotic cells or the more of apoptotic cells. Since the portion of tumor cells underwent apoptosis due to chemotherapeutic growth inhibitors had been prevented first from mitosis. Thus to apply equation 2 in the shrinkage case, the apoptotic tumor portion of half-life time ($t_{1/2}$) would be replaced by the growth portion of doubling time (t_D) which had been prevented first from mitosis whose rate of growth is inversely proportional to the rate of the shrinkage of the apoptotic portion as follows:

$$\left(\frac{V_{\text{Initial}} - V_{\text{Final}}}{V_{\text{Initial}}} \right)_{\text{Shrinkage}} = \left(\frac{V_{\text{Initial}}}{V_{\text{Final}} - V_{\text{Initial}}} \right)_{\text{Growth}} \text{ Eq (3) [25-27,40]}$$

Accordingly, from equations (1) and (2) the alteration in the treated tumor HG from that of the control tumor induced by the drug dose would be equivalent to the energy yield by the drug dose according to the following model:

$$E_{\text{Dose}} = \left[\ln \left(\ln (M - A)_{\text{Treated}} \right)^2 - \ln \left(\ln (M - A)_{\text{Control}} \right)^2 \right] \times C_0 \times h \times 23234.59 \text{ MeV} \text{ Eq (4) [25-27,35-40]}$$

Results and Analysis

To evaluate the *in vivo* activity of TMZ in the LDM regimens

In rats: LDM regimens of TMZ inhibit the tumor growth in C6/LacZ rat glioma orthotopic model in a dose-dependent fashion after daily dosing for 16 days; the average tumor size of control group

grew from 10 mm³ at the implantation to 100 mm³ ($n = 10$, $P < 0.001$) in 17 days [19] with doubling time (t_D) of 5.12 day. While, the average tumor size of the two treated groups by daily dose of 1 and 2 mg/kg for 16 days grew from 10 mm³ to 44.09 and 29.73 mm³ ($n = 10$, $P < 0.001$) in 17 days [19] with t_D of 7.94 and 10.82 days respectively. TMZ doses of 1 and 2 mg/kg/d for 16 days in human (70 kg, 2.5L plasma) are equivalent to ([1 or 2] 16 70 mg/2.5L) 448 and 896 g/mL respectively. Thus from Eqts 1 and 4, the energy yield by 448 and 896 g/mL of TMZ in LDM regimens ($E_{\text{TMZ in LDM}}$) in tumor xenograft of intravenously transplanted 1×10^5 C6/LacZ rat glioma cells were equivalent to:

$$E_{\text{TMZ (448 } \mu\text{g/mL) in LDM}} = \left[\ln \left(\ln \left[\frac{\ln 2}{7.94 \times 24 \times 60 \times 60} \right] \right) \right]^2 - \ln \left(\ln \left[\frac{\ln 2}{5.12 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59 = 1.50309111 \times 10^8 \text{ MeV}$$

$$E_{\text{TMZ (896 } \mu\text{g/mL) in LDM}} = \left[\ln \left(\ln \left[\frac{\ln 2}{10.82 \times 24 \times 60 \times 60} \right] \right) \right]^2 - \ln \left(\ln \left[\frac{\ln 2}{5.12 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59 = 2.53130851 \times 10^8 \text{ MeV}$$

In mice: LDM regimens of TMZ inhibit the tumor growth in U-87MG human glioblastoma model in a dose-dependent fashion after daily dosing for 25 days; the average tumor size of the control group grew from 10 mm³ at the implantation to 120 mm³ in 25 days ($n = 15$, $P < 0.001$) [19] with t_D of 6.97 days. While, the average tumor size of the two treated groups by daily dose of 0.25 and 0.5 mg/kg for 25 days grew from 10 mm³ to 130.1 and 58.77 mm³ in 25 days ($n = 15$, $P < 0.001$) [19] with t_D of 6.74 and 9.78 days respectively. TMZ doses of 0.25 and 0.5 mg/kg/d for 25 days in human (70 kg, 2.5L plasma) are equivalent to ([0.25 or 0.5] 25 70 mg/2.5L) 175 and 350 g/mL respectively. Thus from Eqts 1 and 4, the energy yield by 175 and 350 g/mL of TMZ in LDM regimens ($E_{\text{TMZ in LDM}}$) in tumor xenograft of intravenously transplanted U-87MG human glioma cells were equivalent to:

$$E_{\text{TMZ (175 } \mu\text{g/mL) in LDM}} = \left[\ln \left(\ln \left[\frac{\ln 2}{6.97 \times 24 \times 60 \times 60} \right] \right) \right]^2 - \ln \left(\ln \left[\frac{\ln 2}{6.74 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59 = 1.08680448 \times 10^7 \text{ MeV}$$

$$E_{\text{TMZ (375 } \mu\text{g/mL) in LDM}} = \left[\ln \left(\ln \left[\frac{\ln 2}{9.78 \times 24 \times 60 \times 60} \right] \right) \right]^2$$

$$- \ln \left(\ln \left[\frac{\ln 2}{6.97 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59 =$$

$$1.13689785 \times 10^8 \text{ MeV}$$

Table 1 shows the energy yield by TMZ in the applied LDM regimens.

TMZ doses in $\mu\text{g/mL}$	Energy yield by TMZ in LDM regimens
175	$1.08680448 \times 10^7 \text{ MeV}$
350	$1.13689785 \times 10^8 \text{ MeV}$
448	$1.50309111 \times 10^8 \text{ MeV}$
896	$2.53130851 \times 10^8 \text{ MeV}$

Table 1: Shows the energy yield by TMZ in the applied LDM regimens.

The perfect correlation ($r = 1$) between the logarithms of TMZ doses of the applied LDM regimens and their corresponding energy derived in the presented *in-vivo* study shown in figure 1 and table 1 boost the confidence to establish the following efficient estimation model with a perfect fit ($R^2=1$) shown in figure 2 and expressed in Eqt 5 to describe the energy yield by TMZ dose in LDM regimen.

$$E_{\text{TMZ in LDM}} = 1.48340415 \ln D - 7.55278448 \text{ MeV (Eqt 5),}$$

where D is the TMZ dose in $\mu\text{g/mL}$ applied in LDM regimen, $E_{\text{TMZ in LDM}}$ is the corresponding energy yield of that dose in MeV.




Figure 1: Shows a scatter plot expressing perfect correlation ($r = 1$) between \ln (TMZ dose) and energy yield in metronomic regimens.




Figure 2: Shows the increase of energy yield by increasing TMZ doses in metronomic regimens expressing the non-existence of MGMT-resistance.

Figure 2 shows that the function of the energy yield by TMZ doses in metronomic regimes followed a logarithmic curve of no critical points (1st derivative can't be equivalent to zero) indicating the continuous increase of $E_{\text{TMZ in metronomic regimen}}$ by the increase of the TMZ doses (D) in metronomic regimes up to 2 mg/kg/d. Accordingly, MGMT-resistance to TMZ doesn't exist in metronomic regimens of dose/day up to 2 mg/kg/d.

To evaluate the *in vivo* activity of TMZ in the standard regimens

In rats: Standard regimens of TMZ inhibit the tumor growth in C6/LacZ rat glioma orthotopic model. Kim., *et al.* showed that the average tumor size of control group grew from 10 mm³ at the implantation to 100 mm³ ($n = 10, P < 0.001$) [19] in 17 days with doubling time (t_p) of 5.12 day. Thus the average tumor size of the control group was 25.81 mm³ at the beginning of TMZ therapy on day 7. While, the average tumor size of the treated group orally with 7 mg/kg TMZ for 5 days (between day 7 and 11 after intracranial implantation) to represent a tolerated standard regimen for TMZ therapy grew from 25.81 mm³ to 75.8 mm³ ($n = 10, P < 0.001$) [19] in 10 days from the beginning of TMZ therapy on day 7 to day 17 with t_p of 7.17 days.

While Son., *et al.* showed that the average tumor size of control group grew from 10 mm³ at the implantation to 105 mm³ ($n = 14, P < 0.001$) [20] in 17 days with doubling time (t_p) of 5.01 days. Thus the average tumor size of the control group was 27.65 mm³ at the beginning of TMZ therapy on day 7. While, the average tumor size of the treated group orally with 7.5 mg/kg TMZ for 5 days (between day 7 and 11 after intracranial implantation) to represent a tolerated standard regimen for TMZ therapy grew from 27.65 mm³ to 78.16 mm³ in 10 days from the beginning of TMZ therapy on day

7 to day 17 (n = 14, P < 0.001) [20] with t_d of 6.67 days. TMZ doses of 7 and 7.5 mg/Kg/d for 5 days in human (70 kg, 2.5L plasma) are equivalent to ([7 or 7.5] 5 70 mg/2.5L) 980 and 1050 g/mL respectively. Thus from Eqts 1 and 4, the energy yield by 980 and 1050 g/mL of TMZ in standard regimens (E_{TMZ in standard regimen}) in tumor xenograft of intravenously transplanted C6/LacZ rat glioma cells were equivalent to:

$$E_{\text{TMZ (980 } \mu\text{g/mL) in standard regimen}} = \left[\ln \left(\ln \left[\frac{\ln 2}{7.17 \times 24 \times 60 \times 60} \right] \right) \right]^2 - \ln \left(\ln \left[\frac{\ln 2}{5.12 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59 = 7.89290040 \times 10^7 \text{ MeV}$$

$$E_{\text{TMZ (1050 } \mu\text{g/mL) in standard regimen}} = \left[\ln \left(\ln \left[\frac{\ln 2}{6.67 \times 24 \times 60 \times 60} \right] \right) \right]^2 - \ln \left(\ln \left[\frac{\ln 2}{5.01 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59 = 9.85200300 \times 10^7 \text{ MeV}$$

In mice: Kim., *et al.* showed that the standard regimens of TMZ inhibit the tumor growth in U-87MG human glioblastoma model in a dose-dependent fashion. The average tumor size of the control group grew from 10 mm³ at the implantation to 120 mm³ in 25 days (n = 15, P < 0.001) [19] with t_d of 6.97 days. Thus the average tumor size of the control group was 40.21 mm³ at the beginning of TMZ therapy on day 14. While, the average tumor size of the treated group orally with 1.25 mg/kg TMZ for 5 days, (between day 14 and 18 after intracranial implantation) grew from 40.21 mm³ to 70 mm³ in 11 days (from day 14 to day 25) (n = 15, P < 0.001) [19] with tD of 13.75 days. The average tumor size of the other treated group orally with 2.5 mg/kg TMZ for 5 days, (between day 14 and 18 after intracranial implantation) shrunk from 40.21 mm³ to 36.46 mm³ in 11 days (n = 15, P < 0.001) [19]. From Eq 3,

$$\frac{40.21-36.46}{40.21} = \frac{40.21}{V_0-40.21} \text{ thus the volume of the equivalent virtual grow-}$$

ing image that prevented from mitosis first of this shrinking tumor would be 471.2 mm³ in 11 days with t_d of 3.1 days. TMZ doses of 1.25 and 2.5 mg/Kg/d for 5 days in human (70 kg, 2.5L plasma) are equivalent to ([1.25 or 2.5] 5 70 mg/2.5L) 175 and 350 g/mL respectively. Thus from Eqts 1 and 4, the energy yield by 175 and 350 g/mL of TMZ in the standard regimens (E_{TMZ in standard regimen}) in tumor xenograft of intravenously transplanted 1 × 10⁵ U-87MG human glioma cells were equivalent to:

$$E_{\text{TMZ (175 } \mu\text{g/mL) in standard regimen}} = \left[\ln \left(\ln \left[\frac{\ln 2}{13.75 \times 24 \times 60 \times 60} \right] \right) \right]^2$$

$$- \ln \left(\ln \left[\frac{\ln 2}{6.97 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59$$

$$= 2.25253411 \times 10^8 \text{ MeV}$$

$$E_{\text{TMZ (375 } \mu\text{g/mL) in standard regimen}} = \left[\ln \left(\ln \left[\frac{\ln 2}{6.97 \times 24 \times 60 \times 60} \right] \right) \right]^2$$

$$- \ln \left(\ln \left[\frac{\ln 2}{3.1 \times 24 \times 60 \times 60} \right] \right)^2 \times 1 \times 10^5 \times 23234.59$$

$$= 2.84224710 \times 10^8 \text{ MeV}$$

Table 2 shows the energy yield by TMZ in the applied standard regimens.

TMZ doses in $\mu\text{g/mL}$	Energy yield by TMZ in standard regimens
175	$2.25253411 \times 10^8 \text{ MeV}$
350	$2.84224710 \times 10^8 \text{ MeV}$
980	$7.89290040 \times 10^7 \text{ MeV}$
1050	$9.85200300 \times 10^7 \text{ MeV}$

Table 2: Shows the energy yield by TMZ in the applied standard regimens.

Thus, from the given points in table 2 the following efficient dose-energy model with a perfect fit (R² = 1) was established as shown in figure 3 and expressed in Eq 6 to estimate the energy yield by TMZ dose in the standard regimens.

$$E_{\text{TMZ in standard regimen}} = 1.93 D^3 - 3728.06 D^2 + 1880466.60 D \text{ (Eq 6)}$$

,where D is the TMZ dose in $\mu\text{g/mL}$ applied in standard regimen, E_{TMZ in standard regimen} is the corresponding energy yield of that dose in MeV.

Figure 3 shows that the function of the energy yield by TMZ doses in standard regimens (E_{TMZ in standard regimen}) followed an inverted S-shaped curve having two critical points one of local maximum value (344.21 $\mu\text{g/mL}$, $2.84282515 \times 10^8 \text{ MeV}$) and the other of local minimum value (943.55 $\mu\text{g/mL}$, $7.6530142 \times 10^7 \text{ MeV}$).




Figure 3: Shows energy yield by TMZ doses and the domain of MGMT-resistance in standard regimens.

Accordingly, MGMT-resistance to TMZ was monitored at 344.21 µg/mL TMZ in a standard regimen of 2.45 mg/kg bw/d for 5 days per 28 days/cycle (Figure 4), and then increased by the increase of the standard regimen dose and had a maximum value at 943.55 µg/mL TMZ in a standard regimen of 6.74 mg/kg bw/d for 5 days per 28 days/cycle (Figure 4). Afterwards, MGMT-resistance decreased gradually for MGMT depletion in tumor cells by the increase of the standard regimen dose until the complete suppression at 1243.22 g/mL TMZ in a standard regimen of 8.88 mg/kg bw/d for 5 days per 28 days/cycle (Figure 4). At this point the energy yield by 8.88 mg/kg bw/d for 5 days per 28 days/cycle was equivalent to that yield by 2.45 mg/kg bw/d for 5 days per 28 days/cycle.




Figure 4: Shows the increase in MGMT-resistance to TMZ therapy in tumor cells to its maximum at 6.74 mg/kg bw/day (943.55 µg/mL) and its decrease by increasing TMZ dose/day until being suppressed due to MGMT depletion in tumor cells.

Predicting the *in vivo* activity of TMZ in dose-dense regimen

Dose-dense regimen of TMZ inhibits the tumor growth in human melanoma UACC903 tumor model. The average tumor size of the control group grew from 100 mm³ at the beginning of TMZ therapy at day 10 to 1200 mm³ (n = 15, P = 0.0003) in 26 days at day 36 [22] with t_d of 7.25 days. TMZ doses of 15 mg/kg/d for 5 days (on days 1, 3, 5, 7 and 9) in human (70 kg, 2.5L plasma) are equivalent to (15 × 70 mg/2.5L) 2100 g/mL. From Eq 6, the energy yield by 2100 µM is 5.38196526 × 10⁹ MeV. From Eqts 2 and 4, the histologic grade (H_c) of the treated tumor xenograft of intravenously transplanted 1 × 10⁶ human melanoma UACC903 cells by 2100 µM TMZ in dose-dense regimen of standard schedule of 15 mg/kg/d for 5 days in is supposed to be:

$$H_G = \ln \left(\ln \frac{\ln 2}{7.25 \times 24 \times 60 \times 60} \right)^2 \times 1 \times 10^6 \times 23234.59 + 5.38196526 \times 10^9 = 1.27058538 \times 10^{11} \text{ MeV.}$$

Thus from Eq 2, H_c = 1.27058538 × 10¹¹ MeV / (1 × 10⁶ cells) = 1.27058538 × 10⁵ MeV/23234.59 = 5.46850785 Emad. Accordingly, the treated human melanoma UACC903 tumor model would be predicted to grow with (t_d)_{predicted} = ln 2 × e^{√(5.46850785 Emad)} = 3.37443341 × 10⁶ sec = 39.05594229 days which is 100% identical to that reported by Zhang L., *et al.* about the actual growth of the treated human melanoma UACC903 tumor from 100 mm³ to 158.6 mm³ in 26 days (from day 10 to day 36) (n = 15, P = 0.0003) [22] with actual doubling time (t_DTreated) of $\left(\frac{\ln 2 \times 26}{\ln(158.6/100)} \right)$ 39.07466557 days. Thus, the actual response was 100% identical to the predicted one to provide a clear cut criterion to accept the hypothesis of MGMT-resistance suppression by TMZ in dose-dense regimen and strengthen the confidence in predicting the therapeutic response by dose-energy model for energy yield by TMZ doses in standard regimens shown in Eq 6 and figure 3.

Standard regimen Vs. Metronomic regimen in TMZ therapy

Figure 5 shows a comparison between efficacy of the standard regimens and the metronomic ones to differentiate between them; standard regimens of doses less than 4.37 mg/kg bw/day for 5 days/ cycle of 28 days or more than 8.96 mg/kg bw/day for 5 days/ cycle of 28 days were more efficient than the metronomic ones of equivalent accumulative dose/day along the 28 days (Figure 5). While the metronomic regimens of doses more than 0.78 mg/kg bw/day for 28 days but less than 1.6 mg/

kg bw/day for 28 days were more efficient than the standard ones of equivalent accumulative dose/day for 5 days /cycle of 28 days (Figure 5). In addition by the end of the MGMT-resistance domain, the energy yield by both types of regimens were equivalent demonstrating same efficacy (Figure 5). These findings provide a clear cut criterion to conclude that the ability of tumor cells to repair DNA in TMZ therapy is dependent mainly on dose/day that cells receive but not on the accumulated dose per the treatment cycle. Such ability was shown in the standard regimens starting from 2.45 mg/kg bw/day (344.21 µg/mL) which had decreased the effectiveness of TMZ gradually by the increase of the received dose/day to its minimum at 6.74 mg/kg bw/d (943.55 µg/mL). By increasing the received dose/day in the standard regimens more than 6.74 mg/kg bw/d gradually, the effectiveness of TMZ had been also increased gradually suppressing resistance by depleting MGMT in glioma cells. This led to support the suggestion of the extended dose-dense TMZ regimens for tumors of the advanced and recurrent stages, with the idea that they could potentially deplete MGMT in tumor cells by overwhelming the cells' ability to synthesize MGMT, which might enhance therapeutic activity.




Figure 5: Shows the energy yield in standard and metronomic regimens of TMZ therapy in the absence, presence and suppression of MGMT-resistance in tumor cells.

Discussion

Despite recent advances in the antiangiogenic therapy with TMZ for primary and recurrent gliomas, survival remains poor. The role of MGMT to inhibit the response to TMZ effectively was shown in previous studies where the epigenetic inactivation of MGMT by hypermethylation of the promoter is accompanied by tumor response improvement and survival in gliomas patients [8,47-49]. But in the same time MGMT is consumed in the process of repairing TMZ-mediated DNA damage [50]. Current approach evaluated

the resistance of MGMT expression in gliomas or melanoma to TMZ therapy. For the first time, dosing limits in metronomic or standard or extended dose-dense dosage regimen regarding the MGMT-resistance suppression were identified for all stages of primary or recurrent tumors to overcome resistance to TMZ therapy. The definition of metronomic chemotherapy has been changed several times in a manner which does not necessarily reflect the mechanism of action of the drug, but its pace and dose of administration. LDM regimen is applied on a frequent (daily, several times a week, or weekly) or continuous schedule with no extended interruptions, while dose-dense regimen has the typical schedule of the standard one with the increase in the accumulated dose per the treatment cycle. There are two oncologists' teams; one suggests that LDM regimen exerts its effects exclusively by killing the rapidly dividing endothelial cells in tumors, thus preventing angiogenesis [51,52], while the other supports the dose-dense TMZ regimens provided that MGMT would be depleted in tumors and thus MGMT-mediated resistance would be suppressed [50]. The clinical methodology for staging tumors were conducted as described in earlier studies to determine the energy yield by TMZ doses [35-40]. The consistent results confirm that MGMT-resistance is dependent mainly on the dose/day of the applied regimen. Its effect in tumor cells appears due to TMZ therapy from 2.45 mg/kg bw/day and increases gradually to its maximum at 6.74 mg/kg bw/day, and then decreases gradually by the increase of TMZ dose/day due to MGMT depletion in tumor cells -as shown in figure 4- until being suppressed at 8.88mg/kg bw/day. Accordingly, the dosage limitations of each type of regimen were identified to optimize TMZ therapy for all stages of the disease as follows; from figure 3, it becomes clear that standard TMZ regimens should be applied efficiently (overcome the effect the MGMT-resistance) on the primary tumors of early stages that is characterized by a relatively low histologic grade of administered dose/day doesn't exceed 4.37 mg/kg bw/day for 5 days/ cycle of 28 days. While, efficiency of the standard regimens decreases due to the higher MGMT-resistance in tumor cells of moderate stages of administered dose/day lies between 4.37 and 8.96 mg/kg bw/day for 5 days/ cycle of 28 days as shown in figure 3. Accordingly, LDM TMZ regimens would be more useful for those tumors of moderate stages as shown in figure 5. Thus, for tumor relapse, the accumulated dose/cycle should be increased more than that of the primary one to overcome the acquired resistance by the first-line therapy regardless to type of therapy [28]. If the required accumulated dose/5 days for tumor relapse exceed 4.37 mg/kg bw/d but less than 8.96 mg/kg bw/d, a LDM TMZ regimen should be applied to avoid the higher values of MGMT-resistance and consequently optimize the TMZ dose. The administered dose of the standard

regimen in such case would be distributed equally on all days of the LDM cycle without interruptions. With respect to tumors of the advanced stages, there are two options for choosing the optimal regimen depends on the tumor histologic grade and consequently on the required dose per cycle. If the required dose/5days for the tumor of advanced stage exceeds 8.96 mg/kg bw/d, then the required dose/day will suppress the MGMT-resistance. Accordingly, a dose-dense regimen with standard schedule should be applied taking into account the maximum tolerated dose/day. But, if the required accumulated dose/5 days for the tumor of advanced stages surpasses the maximum tolerated dose/day, then a dose-dense regimen with metronomic regimen should be applied in which the required accumulated dose should be distributed equally on daily doses. The energy yield by TMZ doses in metronomic or standard regimens derived by the two presented dose-energy models (Eqts 5, 6) was identified through *in-vivo* tumor models as conducted and described in earlier studies [25-27,35-40]. The perfect fit ($R^2 = 1$) of dose-energy model of TMZ therapy in metronomic regimen (Eq 5) strengthens the confidence to predict the therapeutic effect prior therapy as conducted and described in an earlier study [25-27,40]. Also, predicting the response of human melanoma UACC903 tumor model to 2100 M TMZ in dose-dense regimen of standard schedule of 15 mg/kg/d for 5 days was 100% identical to the actual response reported by Zhang L., *et al.* [22] to provide a clear cut criterion about accuracy of dose-energy model of TMZ therapy in standard regimen (Eq 6). Predicting the response to TMZ requires to identify each of patient's histologic grade ($H_{G\text{Control}}$) and the estimated energy yield by TMZ dose derived by dose-energy model of either regimen as presented in section of results and analysis. Accordingly to avoid treatment failure, patient's response to TMZ in either regimen should be predicted prior therapy to apply the optimal one for patient-specific H_G .

Conclusion

MGMT-resistance in TMZ therapy is dependent mainly on dose/day received by those cells. Its effect appears when applying TMZ standard regimens of 2.45mg/kg bw/day for 5 days per a cycle of 28 days and increases by the increase dose/day till 6.74 mg/kg bw/d for 5 days per a cycle of 28 days. Thereafter, MGMT depletes in tumor cells gradually by the increase in the received dose/day until suppression of MGMT-resistance. Accordingly, our data suggest that standard TMZ regimen of administered dose less than 4.37 mg/kg bw/day for 5 days every 28 days is more efficient than the metronomic one of dose/day less than 0.78 mg/kg bw/day for 28 days in early stages of primary tumors. While the metronomic TMZ regimen of dose/day lies between 0.78 and 1.6 mg/kg bw/

day for 28 days is more efficient than the standard one of administered dose lies between 4.37 and 8.96 mg/kg bw/day for 5 days every 28 days in the moderate stages of recurrent tumors to avoid the higher values of MGMT-resistance and consequently optimize the TMZ dose. Dose-dense TMZ regimens with standard schedule of dose/day higher than 8.96 mg/kg bw/day for 5 days every 28 days or metronomic schedule of dose/day higher than 1.6 mg/kg bw/day for 28 days suppress the process MGMT-mediated resistance in advanced stages of high-grade tumors by depleting MGMT in tumor cells.

Conflict of Interest

The author declares no conflict of interest.

Bibliography

1. Newlands ES., *et al.* "Temozolomide: a review of its discovery, chemical properties, pre-clinical development and clinical trials". *Cancer Treatment Reviews* 23.1 (1997): 35-61.
2. Ma J., *et al.* "Biochemical changes associated with a multi-drug-resistant phenotype of a human glioma cell line with temozolomide-acquired resistance". *Biochemical Pharmacology* 63.7 (2002): 1219-1228.
3. Friedman HS., *et al.* "Temozolomide and treatment of malignant glioma". *Clinical Cancer Research* 6.7 (2000): 2585-2597.
4. Yung WK. "Temozolomide in malignant gliomas". *Seminars in Oncology* 27 (2000): 27-34.
5. Lee SM., *et al.* "Prognostic significance of O6-methylguanine DNA methyltransferase and p57 methylation in patients with diffuse large B-cell lymphomas". *APMIS* 117.2 (2009): 87-94.
6. Glas M., *et al.* "Long-term survival of patients with glioblastoma treated with radiotherapy and lomustine plus temozolomide". *Journal of Clinical Oncology* 27.8 (2009): 1257-1261.
7. Jacinto FV and Esteller M. "MGMT hypermethylation: a prognostic foe, a predictive friend". *DNA Repair* 6.8 (2007): 1155-1160.
8. Hegi ME., *et al.* "MGMT gene silencing and benefit from temozolomide in glioblastoma". *New England Journal of Medicine* 352.10 (2005): 997-1003.
9. National Cancer Institute of Canada Clinical Trials., *et al.* "Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial". *Lancet Oncology* 10.5 (2009): 459-466.

10. Bobola MS., *et al.* "O⁶-methylguanine-DNA methyltransferase, O⁶-benzylguanine, and resistance to clinical alkylators in pediatric primary brain tumor cell lines". *Clinical Cancer Research* 11.7 (2005): 2747-2755.
11. Hermisson M., *et al.* "O⁶-methylguanine DNA methyltransferase and p53 status predict temozolomide sensitivity in human malignant glioma cells". *Journal of Neurochemistry* 96.3 (2006): 766-776.
12. Bower M., *et al.* "Multicentre CRC phase II trial of temozolomide in recurrent or progressive high-grade glioma". *Cancer Chemotherapy and Pharmacology* 40.6 (1997): 484-488.
13. Brada M., *et al.* "Multicenter phase II trial of temozolomide in patients with glioblastoma multiforme at first relapse". *Annals of Oncology* 12.2 (2001): 259-266.
14. Middleton MR., *et al.* "Randomized phase III study of temozolomide versus dacarbazine in the treatment of patients with advanced metastatic malignant melanoma". *Journal of Clinical Oncology* 18.1 (2000): 158-166.
15. Newlands ES., *et al.* "The Charing Cross Hospital experience with temozolomide in patients with gliomas". *European Journal of Cancer* 32A.13 (1996): 2236-2241.
16. Yung WK., *et al.* "A phase II study of temozolomide vs. procarbazine in patients with glioblastoma multiforme at first relapse". *British Journal of Cancer* 83.5 (2000): 588-593.
17. Yung WK., *et al.* "Multicenter phase II trial of temozolomide in patients with anaplastic astrocytoma or anaplastic oligoastrocytoma at first relapse. Temodal Brain Tumor Group". *Journal of Clinical Oncology* 17.9 (1999): 2762-2771.
18. Tolcher AW., *et al.* "Marked inactivation of O⁶-alkylguanine-DNA alkyltransferase activity with protracted temozolomide schedules". *British Journal of Cancer* 88.7 (2003): 1004-1011.
19. Kim JT., *et al.* "Metronomic treatment of temozolomide inhibits tumor cell growth through reduction of angiogenesis and augmentation of apoptosis in orthotopic models of gliomas". *Oncology Reports* 16.1 (2006): 33-39.
20. Son MJ., *et al.* "Combination treatment with temozolomide and thalidomide inhibits tumor growth and angiogenesis in an orthotopic glioma model". *International Journal of Oncology* 28.1 (2006): 53-59.
21. Lal S., *et al.* "An implantable guide-screw system for brain tumor studies in small animals". *Journal of Neurosurgery* 92.2 (2000): 326-333.
22. Cheng Y., *et al.* "Rational incorporation of selenium into temozolomide elicits superior antitumor activity associated with both apoptotic and autophagic cell death". *PLoS One* 7.4 (2012): e35104.
23. Kyprianou Natasha., *et al.* "Programmed cell death during regression of the MCF-7 human breast cancer following estrogen ablation". *Cancer Research* 51.1 (1991): 162-166.
24. Steel GG. "Growth Kinetics of Tumours. Cell Population Kinetics in Relation to the Growth and Treatment of Cancer". Oxford, United Kingdom: Clarendon Press (1977).
25. Moawad Emad Y. "Identifying and Predicting the Effectiveness of Carboplatin In Vivo and In Vitro and Evaluating its Combination with Paclitaxel". *Indian Journal of Gynecologic Oncology* 13.1 (2015): 1-9.
26. Moawad EY. "Predicting Effectiveness of Imatinib Mesylate in Tumors Expressing Platelet-Derived Growth Factors (PDGF-AA, PDGF-BB), Stem Cell Factor Ligands and Their Respective Receptors (PDGFR- α , PDGFR- β , and c-kit)". *Journal of Gastrointestinal Cancer* 46.3 (2015): 272-283.
27. Moawad EY. "Optimizing and predicting the in vivo activity of AT9283 as a monotherapy and in combination with paclitaxel". *Journal of Gastrointestinal Cancer* 46.4 (2015): 380-389.
28. Moawad Emad Y. "Mass-Energy Conversion in the Decaying System and Doubling Time-Energy Conversion in the Biological System". *Journal of Physics Research and Reviews* 1.1 (2015): 1-13.
29. Moawad E. "Isolated System Towards A Successful Radiotherapy Treatment". *Nuclear Medicine and Molecular Imaging* 44.2 (2010): 123-136.
30. Moawad EY. "Radiotherapy and risks of tumor regrowth or inducing second cancer". *Cancer Nanotechnology* 2.1-6 (2011): 81-93.
31. Moawad EY. "Clinical and pathological staging of the cancer at the nanoscale". *Cancer Nanotechnology* 3.1-6 (2012): 37-46.
32. Moawad EY. "Reconciliation between the clinical and pathological staging of cancer". *Comparative Clinical Pathology* 23.2 (2014): 255-262.
33. Emad Y Moawad. "Pathologic Cancer Staging by Measuring Cell Growth Energy". *Cancer and Oncology Research* 1.3 (2013): 69-74.
34. Emad Y Moawad. "Safe Cancer Screening for Patients after Lumpectomy, Survivors, and Healthy Subjects". *Cancer and Oncology Research* 1.2 (2013): 15-23.
35. Moawad EY. "Administering the optimum dose of l-arginine in regional tumor therapy". *Indian Journal of Clinical Biochemistry* 29.4 (2014): 442-451.

36. Moawad EY. "Induction of multiple sclerosis and response to tyrosine kinase inhibitors". *Indian Journal of Clinical Biochemistry* 29.4 (2014): 491-495.
37. Emad Y Moawad. "Induction of Rheumatoid Arthritis and Response to Tyrosine Kinase Inhibitors". *Universal Journal of Medical Science* 1.2 (2013): 50-55.
38. Emad Y Moawad. "The Mechanism by which Chronic Myeloid Leukemia Responds to Interferon- α Treatment". *Advances in Pharmacology and Pharmacy* 1.2 (2013): 88-94.
39. Moawad EY. "Identifying the optimal dose of ritonavir in the treatment of malignancies". *Metabolic Brain Disease* 29.2 (2014): 533-540.
40. Emad Y Moawad. "Optimal standard regimen and predicting response to docetaxel therapy". *Mutation Research* 770 (2014): 120-127.
41. Emad Y Moawad. "Safe Doses and Cancer Treatment Evaluation". *Cancer and Oncology Research* 1.1 (2013): 6 - 11.
42. Emad Y Moawad. "Nuclear Transmutation and Cancer in the Biological Cell". *International Journal of Biochemistry and Biophysics* 1.1 (2013): 1-8.
43. Emad Y Moawad. "Cell Growth Energy Represents a Measure for Man Health Regulates Nuclear Transmutations and Aberrant Activation in Human Cell". *Universal Journal of Medical Science* 1.2 (2013): 27-35.
44. Moawad EY. "Optimizing Bioethanol production through regulating Yeast Growth Energy". *Systems and Synthetic Biology* 6.3-4 (2012): 61-68.
45. Emad Y Moawad. "Purification of Sewage Water through the Protection of the Environment from Radioactive Contamination". *Energy and Environmental Engineering* 1.2 (2013): 55-61.
46. Emad Y Moawad. "Growth Energy of Bacteria and the Associated Electricity Generation in Fuel Cells". *Bioengineering and Bioscience* 1.1 (2013): 5-10.
47. Esteller M., *et al.* "Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to alkylating agents". *New England Journal of Medicine* 343.19 (2000): 1350-1354.
48. Herrlinger U., *et al.* "Phase II trial of lomustine plus temozolomide chemotherapy in addition to radiotherapy in newly diagnosed glioblastoma: UKT-03". *Journal of Clinical Oncology* 24.27 (2006): 4412-4417.
49. Wick W., *et al.* "Randomized phase III study of sequential radiochemotherapy of oligoastrocytic tumors of WHO-grade III with PCV or temozolomide: NOA-04". *Journal of Clinical Oncology* 26 (2008).
50. Hegi ME., *et al.* "Correlation of O⁶-methylguanine methyltransferase (MGMT) promoter methylation with clinical outcomes in glioblastoma and clinical strategies to modulate MGMT activity". *Journal of Clinical Oncology* 26.25 (2008): 4189-4199.
51. Browder T., *et al.* "Antiangiogenic scheduling of chemotherapy improves efficacy against experimental drug-resistant cancer". *Cancer Research* 60.7 (2000): 1878-1886.
52. Klement G., *et al.* "Continuous low-dose therapy with vinblastine and VEGF receptor-2 antibody induces sustained tumor regression without overt toxicity". *Journal of Clinical Investigation* 105.8 (2000): R15-R24.

Volume 2 Issue 9 September 2018

© All rights are reserved by Emad Y Moawad.