



Plant Histochemistry: A Versatile and Indispensable Tool in Localization of Gene Expression, Enzymes, Cytokines, Secondary Metabolites and Detection of Plants Infection and Pollution

Farid A Badria* and Walaa S Aboelmaaty

Department of Pharmacognosy, Mansoura University, Faculty of Pharmacy, Mansoura, Egypt

*Corresponding Author: Farid A Badria, Department of Pharmacognosy, Mansoura University, Faculty of Pharmacy, Mansoura, Egypt

Received: June 12, 2019; Published: June 20, 2019

DOI: 10.31080/ASPS.2019.03.0318

Abstract

Plants are sources of active metabolites used in medicine and primary sources for isolation of natural products. The medicinal properties of plants are due to the presence of secondary metabolites, including flavonoids, alkaloids, tannins, and saponins, which are of great importance because they possess significant biological activities and the particular active constituents of many crude drugs are still unknown. Histochemical studies are used to confirm identification of cellular and tissue chemical components (secondary metabolites). Histochemical methods are employed in the identification, density of accumulation and distribution of chemical compounds within biological cells and tissues in different organs under microscopes using the color-stain reaction technique and photographic recording. These include the preparation of fixed variably stained specimens and then the examination under the microscopic devices. It is successfully applied in detection and localization of cellular components of active cell constituents such as proteins, carbohydrates, lipids, nucleic acids, and a range of ionic elements occurring in the cell solutions, in addition to identifying the characterization of secretory structures and the chemical nature of the secreted compounds. The methods played a role in describing and tracing the ultrastructure development during different plant growth stages so as the genetic bases of plant physiological and biochemical processes could be further elucidated.

Keywords: Histochemical Localization; Histochemistry; Color-Stain; Secondary Metabolites

Introduction

Histochemistry is the branch of histology dealing with the identification of chemical components of cells and tissues. Starch deposition occurs widely in the plant body, but the particularly common places of its accumulation are seeds, the parenchyma of the secondary vascular tissues in the stem and root, tubers, rhizomes and corn [1]. Starch and proteins are the principal ergastic substances of the protoplast [2]. Tannin is the heterogeneous group of phenol derivatives, usually related to glucosides. Tannins are particularly abundant in the leaves (xylem) of many plants [3]. Saponins are the rare occurrence. Fats are widely distributed in the plant body and they probably occur in small amount in every plant cell [4]. Fats are common reserve material in seeds, spores and embryos in meristematic cells. Glucosides are the degradation product of the carbohydrates. Alkaloids are the degradation product of protein. Many plants contain medicinally important secondary product [5].

Histochemistry is devoted to study the identification and distribution of chemical compounds within and between biological cells, using stains, indicators and light and electron microscopy [6].

Histochemical analysis is essential for the study of plant secretory structures whose classification is based, at least partially, on the composition of their secretion. As each gland may produce one or more types of substances, a correct analysis of its secretion should be done using various histochemical tests to detect metabolites of different chemical classes [7].

Histochemistry is a methodological approach that allows the chemical analysis of cells and tissues in relation to their structural organization [8], but to achieve this objective for plant secretory structures, a wide histochemical analysis is necessary because the same gland and even the same glandular cell can produce several different metabolites simultaneously [9-11].

Materials and Methods

As the histochemical analysis of plant secretory structures uses reagents and dyes that are not specific, certain precautions should be taken to correctly interpret the results: (1) The natural color of the secretion should be observed *in vivo* before applying the test (avoiding the use of reagents with the same color as the

secretion), (2) Attention should be paid to the color obtained in the staining since different colors can be generated in each test but the positive staining is specific and (3) The control procedure should be carefully set up, which usually consists of removing the substance to be detected prior to the test application. The color of the test and that of the control are compared visually to interpret the result [7].

All reagents described below may be applied to fresh or fixed material. If it is necessary to use fixed material, the best fixative for hydrophilic substances is formalin-aceto-alcohol (FAA). For fixation, the material should be immersed in the FAA under vacuum for 24 h, then washed in 50% ethanol overnight and stored in 70% ethanol [12].

Some stains that are commonly used in histochemical localization and their methodology are illustrated in Table 1 for hydrophilic, lipophilic substances, phenolic compounds and alkaloids.

Detection of Hydrophilic substances	
Mucilage	
Ruthenium red staining	This method stains acidic mucilages, pectins [12, 13], and nucleic acids magenta or red (Figure 1a). 1. Apply 0.1% ruthenium red to sections for 5 min. 2. Wash sections twice in distilled water to remove surplus stain. 3. Mount the sections between slide and coverslip with glyceringelatin.
Alcian Blue Staining	This test has a similar result as ruthenium red, staining acidic mucilages, pectins [14] and nucleic acids light blue (Figure 1b). 1. Stain sections with 1% Alcian Blue for 30 min. 2. Rinse sections twice with distilled water to remove surplus stain. 3. Mount the slide with glyceringelatin.
Tannic Acid and Ferric Chloride	This method is based on the reaction of tannic acid with mucilages [15] and pectins, substances which are further revealed by the addition of ferric chloride, producing a grey to black color (Figure 1c). 1. Apply 5% tannic acid for 20 min. 2. Rinse briefly with distilled water. 3. Submerge sections in 3% ferric chloride for 5min. 4. Wash twice in distilled water to remove surplus ferric chloride. 5. Mount the sections using glyceringelatin. 6. Control: Compare the staining obtained in the test with that of sections treated only with tannic acid or with ferric chloride.
Starch	

Lugol's Reagent	This reaction highlights the starch grains in dark blue to black (Figure 1d) [12]. Almost all other structures stain yellow, but this color has no specific significance. 1. Submerge the sections in the Lugol's reagent for 10 min. 2. Rinse briefly with distilled water. 3. Mount the slides using distilled water or Lugol's reagent itself.
Triple Staining for Starch Detection	This triple staining was developed to analyze structural tissue components and the starch grains concomitantly [16]. The application of safranin, astra blue and iodine-potassium iodide solution stains starch grains black, acidic substances (e.g., nucleic acids and lignin) brown, and non-lignified cell walls green (Figure 1e). 1. Stain the sections with 1% safranin for 1 min. 2. Rinse 3 times for few seconds in 50% ethanol to remove surplus stain. 3. Stain with 1% astra blue for 1 min. 4. Wash three times for few seconds in distilled water to remove surplus stain. 5. Apply the iodine-potassium iodide solution for 10 min. 6. Dip sections rapidly in distilled water. 7. Mount the slide with the smallest amount of water.
Carbohydrates	
PAS Reaction (Periodic Acid: Schiff's reagent)	This method is based on the reaction of periodic acid with carbohydrates, forming carbonyl groups revealed by Schiff's reagent [17]. Carbohydrates stain magenta (Figure 1f). 1. Apply 1% sodium tetraborate (freshly prepared) for 30 min. 2. Transfer sections to 1% periodic acid for 10 min. 3. Rinse briefly in distilled water. 4. Apply Schiff's reagent for 15 min in dark. 5. Wash the sections with sodium metabisulfite for 10 min. 6. Rinse in tap water for 10 min. 7. Mount the slides using glyceringelatin. 8. Control: Repeat the test excluding step 2 (periodic acid).
Aniline Blue Staining	This staining marks callose, which may be detected by a green fluorescence under UV light (Figure 1g) [18]. 1. Apply 0.05% aniline blue for 10 min. 2. Rinse briefly in distilled water. 3. Mount the slide in the same buffer used for staining.

Calcofluor White Staining	<p>This test is used to detect cellulose in cell walls, which fluoresces light blue under UV light (Figure 1h) [19].</p> <ol style="list-style-type: none"> 1. Place sections into 0.01% calcofluor white for 10 min. 2. Rinse briefly in distilled water. 3. Mount in distilled water. 	Sudan IV Staining	<p>Sudan IV also stains lipids, in general [14], which become red or red-orange (Figure 2b).</p> <ol style="list-style-type: none"> 1. Apply Sudan IV for 30 min. 2. Rinse briefly in 80% ethanol. 3. Wash in distilled water. 4. Mount in glyceringelatin. 5. Control: As with Sudan black B, the sections should be kept in the extraction solution for at least 6 h.
Proteins		Neutral Red Staining	<p>This fluorochrome emits different colors depending on the lipid composition [22]. Under blue light, the lipids of secretion fluoresce yellow or green (Figure 2f), cuticle fluoresces yellow and lignified cell walls fluoresce red.</p> <ol style="list-style-type: none"> 1. Stain with 0.1% neutral red for 20 min. 2. Rinse briefly in distilled water. 3. Mount in distilled water. 4. Control: As with Sudan black B, the sections should be kept in the extraction solution for at least 6 h.
Aniline Blue Black Staining	<p>This stain reveals proteins in blue (Figure 1i) [20], whether structural or acting in the primary or secondary metabolism.</p> <ol style="list-style-type: none"> 1. Dip sections into 1% aniline blue black for 1 min. 2. Wash twice in 0.5% acetic acid to remove excess stain. 3. Rinse briefly in distilled water. 4. Dehydrate sections passing quickly through 90%, 100% ethanol, then a mixture of 100% ethanol and xylene (1:1, v/v), and finally pure xylene. 5. Mount slides using synthetic resin. 6. Control: Put sections in a solution of acetic anhydride and pyridine (4:6, v/v) for 6 h prior to staining. 	Acidic and Neutral Lipids	
Coomassie Blue Staining	<p>This method stains proteins blue (Figure 1j) [20] and produces a similar result to aniline blue black.</p> <ol style="list-style-type: none"> 1. Stain in 0.25% Coomassie blue for 15 min. 2. Differentiate in 7% acetic acid. 3. Rinse briefly in distilled water. 4. Mount in glyceringelatin. 5. Control: Put sections in a solution of acetic anhydride and pyridine (4:6, v/v) for 6 h prior to staining. 	Nile Blue Staining	<p>Since lipids were detected in the material, Nile blue distinguishes acidic lipids, which stain blue, from neutral lipids, which stain pink (Figure 2c) [23].</p> <ol style="list-style-type: none"> 1. Stain with Nile blue solution for 5 min at 60°C. 2. Wash twice with 1% acetic acid at 60 °C. 3. Rinse in distilled water. 4. Mount in glyceringelatin. 5. Control: As with Sudan black B, the sections should be kept in the extraction solution for at least 6 h.
Detection of Lipophilic substances		Fatty Acids	
Lipids		Copper Acetate and Rubeanic Acid Staining	<p>This method for lipids is slightly more specific than the Sudan tests and identifies fatty acids through the reaction of copper acetate with these acidic lipids, which subsequently turn dark green when exposed to rubeanic acid (Figure 2d) [24,25]</p> <ol style="list-style-type: none"> 1. Treat sections with 0.05% copper acetate for 3 h. 2. Apply 0.1 M Na₂ EDTA (EDTA acid disodium salt solution) for 5 min. 3. Wash in distilled water for 5 min. 4. Transfer sections into 0.1% rubeanic acid (freshly prepared) for 20 min. 5. Wash in 70% ethanol for 5 min. 6. Rinse in distilled water. 7. Mount in glyceringelatin. 8. Control: As with Sudan black B, the sections should be kept in the extraction solution for at least 6 h.
Sudan Black Staining	<p>This is a general method which stains lipids dark blue to black (Figure 2a) [14].</p> <ol style="list-style-type: none"> 1. Stain with Sudan black B for 20 min. 2. Rinse briefly in 70% ethanol. 3. Wash in distilled water. 4. Mount in glyceringelatin. 5. Control: Sections should be kept in the extraction solution [21] for 6 h or more, depending on the composition of the secretion (determined empirically). After this time, the sections should be transferred to distilled water and washed in a period of 4 h (4 × 1 h). Then, the staining proceeds as described. 		

Terpenes	
NADI Reaction	<p>This reagent produces differential staining [26], with essential oils (Figure 2e; monoterpenes and sesquiterpenes) staining blue and resins (diterpenes, triterpenes, tetraterpenes and derivatives) staining red. Mixtures of essential oils and resins produce varied shades of violet to purple, depending on the prevalence of each compound.</p> <ol style="list-style-type: none"> 1. Apply NADI reagent for 1 h in the dark. 2. Wash in sodium phosphate buffer (0.1 M, pH 7.2) for 2 min. 3. Mount in the same buffer. 4. Control: As with Sudan black B, the sections should be kept in the extraction solution for at least 6 h.
Detection of Phenolic Compounds and Alkaloids	
Phenolic compounds	
Ferric Chloride Staining	<p>This method highlights phenolic compounds through iron precipitation, producing a dark color [12], usually black (Figure 3a), sometimes brown.</p> <ol style="list-style-type: none"> 1. Apply 10% ferric chloride for 30 min. 2. Wash twice in distilled water to remove surplus ferric chloride. 3. Mount in glyceringelatin.
Potassium Dichromate Staining	<p>This method also highlights phenolic compounds; in general [27], producing a brown or red-brown color (Figure 3b).</p> <ol style="list-style-type: none"> 1. Apply 10% potassium dichromate for 30 min. 2. Wash twice in distilled water to remove surplus reagent. 3. Mount in glyceringelatin.
Ferrous Sulfate-Formalin Fixation	<p>The best method to detect phenolic compounds is to introduce iron salts into the fixative since the iron compound fixes and stains the phenolic compounds (Figure 3c) [12].</p> <ol style="list-style-type: none"> 1. The samples should be fixed in the ferrous sulfate-formalin solution under vacuum for 48 h. 2. Wash 4 × 2 h (totaling 8 h) in distilled water. 3. Dehydrate the material in 30%, 50%, 70% ethanol for 12 h each. 4. Embed the material according to the chosen technique (Paraplast, Histo-resin, or PEG) and then section in a microtome
Vanillin-Hydrochloric Acid Staining for Tannins	<p>This test is more specific for some phenolic compounds, staining tannins red (Figure 3d) [28]. Use only sections of fresh material.</p> <ol style="list-style-type: none"> 1. Treat with 0.5% vanillin for 20 min. 2. Mount the slide using 9% hydrochloric acid.
Phloroglucinol-Hydrochloric Acid Staining for Lignin	<p>Phloroglucinol in an acidic medium stains lignin in cell walls pink to red (Figure 3e) [12]. It is possible to use either fresh or embedded material.</p> <ol style="list-style-type: none"> 1. Apply 10% phloroglucinol for 15 min. 2. Mount the slides carefully with 25% hydrochloric acid.

Acridine Orange	<p>This fluorescent dye is useful to identify several acidic compounds under blue light, such as nucleic acids and components of the cell wall [29], distinguishing lignified cell walls (yellow-green fluorescence) from non-lignified cell walls (red fluorescence; Figure 3g). It is possible to use fresh material as well as embedded material in this test.</p> <ol style="list-style-type: none"> 1. Apply 0.01% acridine orange for 20 min. 2. Mount the slides with distilled water.
Autofluorescence	<p>Plant tissues have several autofluorescent components which permit their analysis under UV radiation [30]. In relation to secondary metabolites, many phenolic compounds (including lignin) emit a blue or blue-green fluorescence (Figure 1g, 3f).</p> <p>However, it is necessary to be cautious in identifying compounds through autofluorescence because some alkaloids and terpenoids may also emit fluorescence in the blue band [32].</p>
Alkaloids	
Dragendorff's Reagent	<p>This reagent marks alkaloids in red-brown (Figure 3h) [31]. Fresh and fixed material may be used in this method, but fixed material shows a considerably loss of the alkaloids and the staining color when compared to fresh material.</p> <ol style="list-style-type: none"> 1. Treat with Dragendorff's reagent for 20 min. 2. Rinse briefly in 5% sodium nitrite. 3. Mount in distilled water. 4. Control: Treat sections with 5% tartaric acid in 95% ethanol for 72 h and repeat the staining procedure.
Wagner's Reagent	<p>This method also stains alkaloids red or red-brown (Figure 3i) [32]. It is recommended that fresh material be used for this test.</p> <ol style="list-style-type: none"> 1. Apply Wagner's reagent for 20 min. 2. Rinse briefly in distilled water. 3. Mount in distilled water. 4. Control: Treat sections with 5% tartaric acid in 95% ethanol for 72 h and repeat the staining procedure.

Table 1: Some stains that are commonly used in histochemical tests.

Applications of histochemistry in plant research
In detection and localization of secondary metabolites in certain medicinal plants

Histochemical study was carried out to localize polyphenolic terpenoid aldehydes and fixed oil in healthy seeds, stems, leaves and roots of *Gossypium barbadense* L. var. Giza 86. In all examined organs, polyphenolic terpenoid aldehydes and fixed oil were mainly detected inside lysigenous glands. In young leaves and roots, polyphenolic aldehydes were also observed as fine particles inside




Figure 1: Histochemical analysis of 1 structures. (a, c, g–i) Embedded material. (b, d–f, j) Fresh material. (a) Detection of acidic mucilage in the colleter of *Asclepias curassavica* L. (Apocynaceae) by ruthenium red. (b) Identification of acidic mucilage in the epidermis of *Rhododendron* sp. (Ericaceae) by Alcian Blue. (c) Positive result for mucilage in secretory idioblast of *Cattleya walkeriana* Gardner (Orchidaceae) using tannic acid and ferric chloride. (d) Starch grains in the nectary of *Inga edulis* Mart. (Fabaceae) detected by Lugol's reagent. (e) Observation of starch grains in laticifer of *Euphorbia milii* Des Moul. (Euphorbiaceae) and the tissue structure using the triple staining. (f) Detection of carbohydrates in the secretory idioblast of *Ceiba speciosa* [7].




Figure 2: Histochemical analysis of plant secretory structures. Fresh material. (a) Detection of lipids production by elaiophore of *Byrsonima intermedia* A. Juss. (Malpighiaceae) by Sudan black B. (b) Identification of lipids in the secretory duct of *Philodendron* sp. (Araceae) by Sudan IV. (c) Positive result for neutral lipids in the laticifer of *Sapium glandulatum* (Vell.) Pax (Euphorbiaceae) by Nile blue. (d) Observation of fatty acids in the glandular trichome of *Tetradeniariparia* (Hochst.) Codd (Lamiaceae) using copper acetate and rubeanic acid. (e) Detection of essential oils in the secretory idioblasts of *Peplonia axillaris* (Vell.) Fontella and Rapini (Apocynaceae) by NADI reagent. (f) Identification of lipids in secretory duct of *Kielmeyera apparicioniana* Saggi (Calophyllaceae) by neutral red under blue light. (g) Positive result for lipids in glandular trichome of *Tetradeniariparia* by Nile blue under blue light [7].




Figure 3: Histochemical analysis of plant secretory structures. (a, b, d, f-i) Fresh material. (c, e) Embedded material. (a-c) Detection of phenolic compounds in secretory idioblasts. (a) Ferric chloride. *Acalyphaamentacea* Roxb. (Euphorbiaceae). (b) Potassium dichromate. *Calliandrat weediei* Benth. (Fabaceae). (c) Ferrous sulfate in formalin [7].

the cytoplasm of some parenchymatous cells around glands. Lysigenous terpenoid-containing glands were noticed in all tap root regions except the apical 3 cm. The number of glands increased with increasing distance from the root tip. This may explain why the antimetabolic activity of gossypol does not affect the growing tip of the plant [33].

Three species used in folk medicine were chosen to determine their histochemical investigation: *Adhatod azeylanica*, *Ruta graveolens* and *Vitex negundo*. In general, these plants are used in folk-medicine in the treatment of gonorrhoea, antiperiodic, bronchitis, infected wounds, scrotal swelling, synovitis, arthritis pain and rheumatic arthritis. For histochemical studies the free hand sections of leaves and stem were taken and treated with the respective reagent to localize components, viz. starch, protein, tannin, saponin, fats, glucosides and alkaloids in the tissues [34].

A histochemical analysis of leaf and rhizome of *Curcuma neilgherrensis* was done. The study revealed the identification and location of the phytochemicals like alkaloids, saponins, tannins, oils, starch grains etc in various regions of leaf and rhizome of *C. neilgherrensis*. Free hand sections were taken and treated with respective reagents to localise the various cellular components. The observations could be of great use in chemotaxonomy and checking the drug adulteration [35].

Furthermore, the histochemical studies of leaves and wood of *Sesbania grandiflora*, *Sesbania spinosa* and *Sesbania cannabina* are medicinally important plants of Marathwada region in Maharashtra. For histochemical studies the free hand sections of leaves and wood were taken and treated with the respective reagent to localize components, viz. starch, protein, tannin, saponin, fat, glucosides and alkaloids in the tissues [36].

Solidago chilensis Meyen, belonging to the family *Asteraceae*, is a plant native to South America and the only representative of the genus in Brazil. This species is popularly known as "arnica" and is used to treat bruises, muscle pain and inflammation. Cross-sections were obtained, by freehand, for microscopic analysis of root, stem and leaf; for these parts of the plant maceration was also performed according to the method of Jeffrey. For the leaf were still made paradermal sections, scanning electron microscopy analysis, phytochemical and histochemical tests. Thus, it was determined anatomical features useful for diagnosis of the species which, together with identification of the chemical compounds and its histo localization, provides support to their quality control [37].

Carbohydrate storage in the form of starch grains has been examined in stems and roots of *Jatropha curcas*. The predominant starch-storing tissues were identified, and the cellular localization of the starch grains within these tissues was determined. In stem sections, starch was seen predominantly in parenchymatous cortex, medullary rays, pith while in the root sections, starch was seen

highly concentrated only in cortical tissues and to some extent as brownish black patches in the medullary rays [38].

Thymus quinqueco status, with more medical value, is a kind of wild plants. In order to exploit and utilize this plant, we studied the species and locations of alkaloids in its leaves. A histochemical study of leaves at different developing stages was taken to localize the alkaloids. Meanwhile, the kinds and content of alkaloids in leaves were identified using GC-MS technique. It was found that there were two kinds of glandular trichomes, namely, peltate trichomes and capitate trichomes, on the surface of leaves, and their secretory cells could secrete alkaloids. Results showed that trichomes could secrete alkaloids as soon as the first pair of leaves formed, and there were altogether 18 kinds of alkaloids identified by GC-MS. Nearly all of these alkaloids of leaves at different developing stages were distinct from each other, except one, 3-methoxy-a-methyl-benzeneethanamine, persists at different developing stages with high concentration [39].

Studies on the phytochemical profiling and histochemical localization in leaf and stem of *Trichosanthes cucumerina* (L) var. *cucumerina* with reference to the influence of plant age and geographical variations were carried out using Wagner's reagent concluded that alkaloids are mostly located in the parenchyma cells bordering the vascular bundles of stem and petiole. Flavonoids, localized with NaOH revealed that they were marked as a distinct yellow band in the sub hypodermal layer of stem. Tannin localization with FeCl₃ also suggested their storage in parenchyma cells [40].

Studies on the phytochemical profiling and histochemical localization in leaf and stem of *Trichosanthes cucumerina* (L) var. *cucumerina* with reference to the influence of plant age and geographical variations were carried out using Wagner's reagent concluded that alkaloids are mostly located in the parenchyma cells bordering the vascular bundles of stem and petiole. Flavonoids, localized with NaOH revealed that they were marked as a distinct yellow band in the sub hypodermal layer of stem. Tannin localization with FeCl₃ also suggested their storage in parenchyma cells [40].

An evaluation of the histochemistry of the leaf blade of *Byrsionimaverbascifolia* (L.) DC., *Malpighiaceae*, *Cam-pomanesiaadamantium* (Cambess.) O. Berg, *Myrtaceae*, *Roupalamontana* Aubl., *Proteaceae*, and *Solanumlycocarpum* A. St.-Hil., *Solanaceae*, species that have been reported as producers of secondary metabolites for pharmacological use was performed.

Tests were conducted for lipids, terpenoids, phenolic compounds, alkaloids, sugars and proteins. Alkaloids were observed only in *R. montana*, as well as the results for phenolic compounds. Flavonoids are present in *B. verbascifolia* and *R. montana*. The lipid composition was showed for the chemical compounds of *B. verbascifolia* and *C. adamantium*, which proved to be part of the es-

sential oils or resins oils in *C. Adamantium* idiolects. The chemical compounds of *B. verbascifolia*, *C. adamantium* and *R. montana* are present mainly in idioblasts among the parenchyma and epidermal cells. *C. adamantium* has secretory cavities, but only with lipid content [41].

The genus *Barleria lupulina* Lindl of the family *Acanthaceae* belongs to the sub tribe *Barlerieae* of the tribe *Justicieaesensu* Benth and Hook. An investigation mainly emphasized on the histochemical localization of phytochemicals like alkaloids, starch, tannins, reducing sugars, proteins, flavonoids, amino acids and lignins. The active compounds were identified prominently in different locations of the stem, leaf petiole and root of the medicinal plant *B lupulina* under study. It was found that presence of number of phytochemicals in xylem is higher than other tissues [42].

Histochemical localization of starch, protein, lipid and lignin in the callus, field-grown and *in vitro* raised plants

Scopariadulcis L. (*Scrophulariaceae*), commonly known as bitter broom or kallurukki, is much valued in tradition medicine to treat respiratory, gastric and hepatic disturbances, kidney stones, diabetes and inflammations. Histochemical localization of starch, protein, lipid and lignin was done in the calli and regenerated plants of *S. Dulcis* grown in MS medium supplemented with 0.1 mg/l IBA, and also in the field-grown plants. Starch grains were abundant in the cortex and pith of field-grown plants and also in the calli, while the *in vitro* raised plants had lesser starch content. Intense accumulation of proteins was observed in the calli and *in vitro* raised plants. Both the field-grown and *in vitro* raised plants showed accumulation of lipids in the cortex and pith, and heavy deposition of lignin in the xylem elements. The calli showed a greater deposition of starch, proteins, lipids and lignin, when compared to both the field-grown as well as *in vitro* raised plants. The study reveals the potential of utilizing calli in herbal formulations of the species, as this may yield better results including improved nutraceutical value [43].

In the supposed defense mechanism actions (Parasitic plant host - root interaction)

Roots of different hosts of the holoparasitic weed known as broomrape (*Orobancha* spp.) were examined histochemically for the occurrence of structural cellular barrier formation following wounding/penetration. Such barrier might function to impede the successful development of parasite haustorium interaction, i.e. as a self defence mechanism. In faba bean and white bean, brown deposits occurred in walls adjacent to the damaged cells of the epidermis, cortex and stele. Via stain reactions and colorations these deposits were detected as melanin. Additionally, walls bordering damaged site at the level of the endodermis and within the stele becomes suberized and lignified. In peas, which possesses a lignified hypodermis, the response was similar but lignin was also de-

posited in the walls of the endodermis and hypodermis adjacent to the wound. In sunflower, which possesses a suberized hypodermis, melanin was deposited in the hypodermis and lignin and suberin occurred within the stele. In all these broomrape host species melanization conferred the modified cell wall many of the properties associated with lignified and suberized structures such as impermeability and resistance to chemical degradation [44].

In detection of active substances in aromatic plant; citral accumulation in lemongrass

The sites of citral accumulation in lemongrass (*Cymbopogon flexuosus* Nees ex Steud) wats (cultivar OD-19) were located by Schiff's reagent, which upon its reaction with aldehydes (citral) gives a purple-red colouration. Using this technique, single oil-accumulating cells were detected in the adaxial side of leaf mesophyll commonly adjacent to non-photosynthetic tissue and between vascular bundles. In this respect, however, the citral lacking cultivar GRL-1 (geraniol rich) leaf sections, which also was subjected to Schiff's reagent could be compared to the cultivar OD-19 leaf sections. In lemongrass mutant GRL-1, those specialized cells, however, are not being stained due to lack of citral. Hence, it could be confirmed that the observed schiff's staining reaction is associated with the accumulation of citral substance in a given cell [45].

New applications

In localization of the site of monoterpene phenols accumulation in plant secretory structures

A new method is reported for the histochemical localization of monoterpene phenols in essential oil secretory structures. The method was adapted from a spot test originally devised for *in vitro* detection of phenolic compounds in organic analyses. Plant subjects were the *Lamiaceae* species *Thymus vulgaris* L., *Oreganum vulgare* L. and *Mentha x piperita* L., which accumulate essential oil in glandular trichomes. A reagent consisting of 4-nitrosophenol in conc. H_2SO_4 was applied to sample leaves of each species. A positive test for phenol was indicated by the production of coloured indo-phenols. Using this method, monoterpene phenols were identified in the trichomes of *T. vulgaris* (thymol) and *O. vulgare* (carvacrol), indicated by colour changes to red and green respectively [46].

Detection of cytokinins and auxin in plant tissues using histochemistry and immunocytochemistry

A new method for histochemical localization of cytokinins (CKs) in plant tissues based on bromophenol blue/silver nitrate staining was reported. The method was validated by immunohistochemistry using anti-trans-zeatin riboside antibody. Indole-3-acetic acid (auxin, IAA) was localized by anti-IAA antibody in plant tissues as a proof for IAA histolocalization. Root sections were used, because they are major sites of CKs synthesis, and insect galls of *Piptadeni-agonoacantha* that accumulate IAA. Immunostaining confirmed the presence of zeatin and sites of accumulation of IAA indicated

by histochemistry. The colors developed by histochemical reactions in free-hand sections of plant tissues were similar to those obtained by thin layer chromatography (TLC), which reinforced the reactive sites of zeatin. The histochemical method for detecting CKs is useful for galls and roots, whereas IAA detection is more efficient for gall tissues. Therefore, galls constitute a useful model for validating histochemical techniques due to their rapid cell cycles and relatively high accumulation of plant hormones [47].

Locating enzymes in plant tissue using nitrocellulose blotting

The blotting of fresh tissue sections onto nitrocellulose and the detection of enzyme activities on the blots for polyphenol oxidase, peroxidase, glycosidases, dehydrogenase and phosphatase activity has been shown successfully using a simple histochemical method. Two cm squares of nitrocellulose membrane filters BA 28, (Schleicher and Schüll, 0.45 μm pore size) were soaked in distilled- H_2O , placed on microscope slides and blotted dry with a tissue. A 2-3mm section through the plant tissue was placed on the membrane and then lightly pressed on it using an additional microscope slide. The section was removed carefully and the membrane thoroughly rinsed with distilled H_2O to remove non-proteinaceous material or soluble compounds and lightly pressed with a tissue to remove excess water. About 0.2 ml of the appropriate substrate or reagent was next spread over the surface of the membrane. The slides were incubated in a moist atmosphere for the reaction to proceed, the substrate was then rinsed off, and if necessary a developing reagent added. For fast reactions with immediate colour production the substrate was added and rinsed off as soon as colour development was optimal e.g. polyphenol oxidase or peroxidase. Alternatively the nitrocellulose blot can be placed (blot side upwards) on to a piece of filter paper moistened with substrate. This gives very good resolution and localization of enzyme activity [48].

In gene expression

The detection of plant transformation (using β -glucuronidase; GUS assay in *Lilium* are carried out via the application of histochemical methods. β -glucuronidase (GUS) assay is used to assess transient expression of the GUS gene using 5-bromo-4-chloro-3-indolyl β D-glucuronide (X-Gluc) as the substrate. Six days after co-cultivation, samples of 0.1g of callus collected from each treatment are subjected to transient histochemical GUS assay. The transformation efficiency of calli are evaluated by counting the number of blue spots, using stereo microscope, showing GUS enzyme activity on each callus sample [49]. The methods/technique are described by Azadi, *et al.* [50].

The use of the histochemical methods in iron and ferritin gene expression in transgenic indica rice (*Oryza sativa* L. Cv *Pusa basmati*) proved as efficient in such an investigation. Perl's Prussian blue staining of transgenic rice grain sections show distribution of iron accumulation (blue compound of ferricferrocyanide) through-

out the allure one and sub aleurone layers and in the central region of the starch endosperm. Whereas, in the nontransgenic grains, blue colour formation indicating iron accumulation was restricted to the aleurone layer and the intensity of color was also very low. Transverse section of the transgenic rice grains indicated the high iron accumulation in embryo as well as in the endosperm, in comparison to the non transgenic ones. In the latter, iron appeared restricted to the embryo and aleurone layer in which the intensity of color detected in the embryo was very low. This histochemical analysis of iron in rice specifically showed temporal and spatial deposition of storage iron [51].

Advances in histochemistry and cytochemistry made are possible to retrieve quantitative data from 2D and 3D microscopic images. In this way, valid quantitative results can be regenerated (e.g. gene expression data at the mRNA, protein and activity levels) from microscopic images in relation to structures in cells, tissues and organs in 2D and 3D. Volumes, areas, lengths and numbers of cells and tissues can be calculated and related to these gene expression data while preserving the 2D and 3D morphology [52].

In detection of heavy metals, i.e. pollution and contamination Heavy metals accumulation

Histochemical methods were employed in the detection of the heavy metals (Cd, Pb, Ni, and Zn) and strontium, their distribution, accumulation, and translocation within the tissues of higher plants. In this respect, detailed protocols of metal detection with metallochrome indicators dithizone (Cd, Pb), dimethylglyoxime (Ni), sodium rhodizonate (Sr), zincon (Zn), and fluorescent indicator Zinpyr_1 (Zn) by light and fluorescence microscopy were described [53].

The occurrence of heavy metals and their accumulation in water hyacinth [*Eichhornia crassipes* (Mart.) Solms] was investigated. The histochemical staining examinations indicated the accumulation in the epidermis and vascular bundles of the roots and petiole. In the leaf sections the palisade tissues were deeply stained, showing the high accumulation of the metals within the leaves [54].

In detection of H_2O_2 and O_2^- ; a result of cadmium contamination

The effect of cadmium on H_2O_2 and O_2^- production in leaves from pea plants grown for 2 weeks with 50 μm Cd, by histochemistry with diaminobenzidine (DAB) and nitro blue tetrazolium (NBT), respectively was studied. The subcellular localization of the reactive oxygen species (ROS) was studied by using CeCl_3 and Mn/DAB staining for H_2O_2 and O_2^- , respectively, followed by electron microscopy observation. In leaves from pea plants grown with 50 μm CdCl_2 , a rise of six times in the H_2O_2 content took place in comparison to control plants. The accumulation of H_2O_2 was localized mainly in the plasma membrane, mesophyll and epidermal cells, as

well as in the tonoplast of bundle sheath cells. In mesophyll cells, the accumulation of H_2O_2 was observed in mitochondria and peroxisomes. Localization of O_2^- production was demonstrated in the tonoplast of bundle sheath cells, and plasma membrane from mesophyll cells. The Cd-induced production of the ROS, H_2O_2 and O_2^- , could be attributed to the phytotoxic effect of Cd. In this connection, lower levels of ROS were assumed to function as signal molecules for the induction of defense genes against the injurious effects of the heavy metal [55].

However, Jin, *et al.* found that exposure to cadmium resulted in significant ultra structure changes in the root meristems and leaf mesophyll cells of *Sedum alfredii* Hance. Damages were more pronounced in NHE (non-hyperaccumulator ecotypes) even when Cd concentrations were one-tenth of those applied to HE (hyperaccumulator ecotypes). In the cadmium stress damaged chloroplasts resulted in imbalanced lamellae formation which is coupled with early leaf senescence. Histochemical results revealed that glutathione (GSH) biosynthesis inhibition led to overproduction of hydrogen peroxide (H_2O_2) and superoxide radical (O_2^-) in HE but not in NHE. The GSH biosynthesis induction in root and shoot exposed to elevated Cd conditions, however, might be involved in Cd tolerance and hyper accumulation in HE of *S. alfredii* H [56].

In the detection of aluminum accumulation, lipid peroxidation, callose production, and plasma membrane integrity

In peas (*Pisum sativum* L.) roots, staining were observed to distribute similarly on the entire surface of the root apex regarding aluminum accumulation, lipid peroxidation, and callose production. Meanwhile, the loss of plasma membrane integrity (detected by Evans blue uptake) was localized exclusively at the periphery of the cracks on the surface of root apex. The enhancement of four phenomena, i.e. aluminium accumulation, lipid peroxidation, callose production and root elongation inhibition displayed similar aluminum dose dependencies which occurred at 4hrs exposure. The loss of membrane integrity, however, was enhanced at lower aluminum concentrations and after a longer aluminum exposure of 8hrs. The addition of butylatedhydroxyanisole (a lipophilic antioxidant) during aluminum treatment was found to completely prevent only the lipid peroxidation and callose production by 40%. Thus, lipid peroxidation was suggested to represent relatively early symptom induced by the accumulation of aluminum and appear to cause, in part, callose production. Whereas, the loss of plasma membrane integrity represented a relatively late symptom caused by cracks in the root is due to the inhibition of root elongation [57].

Conclusion

Many plants contain medicinally important secondary products. Therefore, histochemical investigations of different plant parts dealing with the identification of chemical components of cells and

tissues. Furthermore, histochemical localization has numerous and unique economic and pharmaceutical implications. Based on the investigation, this review concludes that histochemical localization could be used in a rapid field survey to identify the existence of bioactive compounds in certain plants.

Bibliography

1. Kadam V B. "Histochemical investigations of different organs of three Endangered medicinal taxa of South Gujarat Forests". *Journal of Phytochemical Research* 12 (1999): 109-112.
2. Kuster E. "Die pflanzenzelle", 3rd ed, Jene Gustav Fischer verlag, Jena (1956): 232.
3. Kadam V B, *et al.* "Nutritional status of seeds of some tree species". *Journal of Environmental Biology* 5 (1996): 96-98.
4. Seifriz W. "Protoplasm Mac Graw Hill Book Company". Inc. New York (1936).
5. Dhar M L, *et al.* "Screening of Indian Plants for Biological Activity-Part I". *Indian Journal of Experimental Biology* 6.4 (1968): 232-247.
6. Wick M R. "Histochemistry as a tool in morphological analysis: A historical review". *Annals of Diagnostic Pathology* 16.1 (2012): 71-78.
7. Demarco D. "Histochemical Analysis of Plant Secretory Structures", chapter 24, Carlo, P. and Marco, B. (eds.), *Histochemistry of Single Molecules: Methods and Protocols, Methods in Molecular Biology*. 1560 (2017).
8. Coleman R. "The impact of histochemistry - a historical perspective". *Acta Histochemica* 102.1 (2000): 5-14.
9. Fahn A. "Secretory tissues in vascular plants". *New Phytology* 108.3 (1988): 229-257.
10. Evert R F. "Esau's plant anatomy". 3rd edn., John Wiley & Sons, Hoboken, NJ (2006).
11. Demarco D. "Micromorfología y histoquímica de los laticíferos de órganos vegetativos de especies de Asclepiadoideae (Apocynaceae)". *Acta Biológica Colombiana* 20.1 (2015): 57-65.
12. Johansen D A. "Plant microtechnique." McGraw-Hill, New York (1940).
13. Gregory M and Baas P. "A survey of mucilage cells in vegetative organs of the dicotyledons". *Israel Journal of Botany* 38 (1989): 125-174.
14. Pearse A G E. "Histochemistry: theoretical and applied". 2 4th ed., C. Livingstone Edinburgh (1985).

15. Pizzolato T D. "Staining of Tiliamucilages with Mayer's tannic acid-ferric chloride". *Bulletin of the Torrey Botanical Club* 104.3 (1977): 277-279.
16. Demarco D., et al. "Two laticifer systems in *Sapiumhaematospermum*- new records for Euphorbiaceae". *Botany* 91 (2013): 545-554.
17. McManus J F A. "Histological and histochemical uses of periodic acid". *Stain Technology* 23.3 (1948): 99-108.
18. Smith M M and McCully M E. "A critical evaluation of the specificity of aniline blue induce fluorescence". *Protoplasma* 95.3 (1978): 229-254.
19. Hughes J and McCully M E. "The use of an optical brightener in the study of plant structure". *Stain Technology* 50.5 (1975): 319-329.
20. Fisher D B. "Protein staining of ribbonedepon sections for light microscopy". *Histochemistry* 16.1 (1968): 92-96.
21. High O B. "Lipid histochemistry". Oxford University Press, New York, NY (1984).
22. Kirk P W. "Neutral red as a lipid fluorochrome". *Stain Technology* 45 (1970): 1-4.
23. Cain A J. "The use of Nile Blue in the examination of lipids". *Quarterly Journal of Microscopical Science* 88 (1947): 383-392.
24. Ganter P and Jollés G. "Histochimienormaleetpathologique, vol 1. Gauthier-Villars, Paris". (1969).
25. Ganter P and Jollés G. "Histochimienormaleetpathologique, vol 2. Gauthier-Villars, Paris (1970).
26. David R and Carde J P. "Coloration différentielle des inclusions lipidique et terpeniques des pseudophylles du Pin maritime au moyen du reactifNadi". *Comptes rendus de l'Académie des Sciences* 258 (1964): 1338-1340.
27. Gabe M. "Techniques histologiques". Masson & Cie, Paris (1968).
28. Mace M E and Howell C R. "Histochemistry and identification of condensed tannin precursor in roots of cotton seedlings". *Canadian Journal of Botany* 52.11 (1974): 2423-2426.
29. Ruzin S E. "Plant microtechnique and microscopy". Oxford University Press, New York, NY (1999).
30. Talamond P, et al. "Secondary metabolite localization by autofluorescence in living plant cells". *Molecules* 20.3 (2015): 5024-5037.
31. Svendsen A B and Verpoorte R. "Chromatography of alkaloids. Elsevier Scientific Publishing Company, New York, NY (1983).
32. Furr M and Mahlberg P G. "Histochemical analyses of laticifers and glandular trichomes in *Cannabis sativa*". *Journal of Natural Products* 44.2 (1981): 153-159.
33. Badria F A., et al. "Histochemical Localization Of Polyphenolic Aldehydes In *Gossypium Barbadense*". *International Journal of Research in Pharmaceutical and Nano Sciences* 7.5 (2018): 200-204.
34. Dhale D A. "Histochemical investigation of some medicinal plants". *ARPB* 1.2 (2011): 147-154.
35. Nissar A R., et al. "Histochemical Studies of *Curcuma Neilgherrensis*-An Antidiabetic Herb". *WJPR* 3.9 (2014): 731-741.
36. Momin R K and Kadam V B. "Histochemical investigation of different organce of genus *sesbania* of marathwada region in Maharashtra". *Journal of Phytology* 3.12 (2011): 31-34.
37. Douglas M F, et al. "Anatomical, phytochemical and histochemical study of *Solidagochilensis* Meyen". *Annals of the Brazilian Academy of Sciences* 90.2 (2018): 2107-2120.
38. Kumari V, et al. "Histochemical localization of starch in stem and root of *Jatropha curcas* (Euphorbiaceae)". *Bulletin of Environment, Pharmacology and Life Sciences* 7.4 (2018): 7-10.
39. Jing H., et al. "Histochemical Investigation and Kinds of Alkaloids in Leaves of Different Developmental Stages in *Thymus quinquecostatus*". *The Scientific World Journal* (2014): 1-6.
40. Aryakrishna U K., et al. "Phytochemical profiling and histochemical localization in leaf and stem of *Trichosanthes cucumerina* (L) var. *cucumerina* with reference to the influence of plant age and geographical variations". *Journal of Pharmacy Research* 10.10 (2016): 660-665.
41. Kuster V C and Vale F H. "Leaf histochemistry analysis of four medicinal species from Cerrado". *Revista Brasileira de Farmacognosia* 26.6 (2016): 673-678.
42. Sudipa M and Subrata M. "Studies on Screening And Histochemical Localisation Of Phytochemicals In The Medicinal Plant *BarleriaLupulina* Lindl". *IJPDT* 4 (2014): 95-97.
43. Thushara T and Devipriya V. "Histochemical localization of starch, protein, lipid and lignin in the callus, field-grown and in vitro raised plants of *Scopariadulcis* L". *International Journal of Scientific & Engineering Research* 8.9 (2017): 743-747.
44. Brammall R A and Hassan E A. "Histochemistry of wound healing response in roots of several hosts of *Orobanche* spp. The 61st Annual Meeting of the Canadian Phyopathological Society, Toronto, Ontario, Canada (1995): 43.

45. Luthra R, *et al.* "Histochemical localization of citral accumulating site in lemongrass (*Cymbopogon flexuosus* Ness Ex. Steud) wats cultivar OD-19". *Asian Journal of Plant Science* 6 (2007): 419-422.
46. Gersbach P V, *et al.* "A New Histochemical Method for Localization of the Site of Monoterpene Phenol Accumulation in Plant Secretory Structures". *Annals of Botany* 88.4 (2001): 521-525.
47. Bedetti C S. "Detection of cytokinins and auxin in plant tissues using histochemistry and immunocytochemistry". *Biotechnic & Histochemistry* 93.2 (2018): 1-6.
48. Spruce J, *et al.* "A Simple Histochemical Method for Locating Enzymes In Plant Tissue Using Nitrocellulose Blotting". *Phytochemistry* 26.11 (1987): 2901-2903.
49. Stomp A M. "Histochemical localization of β -glucuronidase. In: Gallagher SR (ed) GUSprotocols: using the GUS gene as a reporter of geneexpression". *Academic Press San Diego* (1992):103-113.
50. Azadi P, *et al.* "Macro elements inoculation and co-cultivation medium strongly affect the efficiency of Agrobacterium-mediated transformation in Liliium". *Plant Cell Tissue Organization Culture* 101.2 (2010): 201-209.
51. Sivaprakash K R, *et al.* "Tissue-specific histochemical localization of iron and ferritin gene expression in transgenic indica rice Pusa Basmati (*Oryza sativa* L.)". *Journal of Genetics* 85.2 (2006): 157-160.
52. Chieco P, *et al.* "Image Cytometry: Protocols for 2D and 3D Quantification in Microscopic Images". *Progress in Histochemistry and Cytochemistry* 47.4 (2013): 211-333.
53. Seregin I V and Kozhevnikova A D. "Histochemical methods for detection of heavy metals and strontium in the tissues of higher plants". *Russian Journal of Plant Physiology* 58.4 (2011): 721-727.
54. Warriar R R and Saroja S. "Histochemical studies on water hyacinth with particular references to water pollution". *International Journal of Integrative Biology* 3.2 (2008): 96-99.
55. Romero-Puertas M C, *et al.* "Cadmium-induced subcellular accumulation of O_2^- and H_2O_2 in pea leaves". *Plant Cell and Environment* 27.9 (2004): 1122-1134.
56. Jin X, *et al.* "Effects of cadmium on ultrastructure and antioxidant defense system in hyperaccumulator and non-hyperaccumulator ecotypes of *Sedum album*". *Journal of Hazardous Materials* 156 (2008): 387-397.
57. Yamamoto Y, *et al.* "Lipid Peroxidation is an early symptom triggered by aluminum, but not the primary cause of elongation inhibition in pea roots". *Plant Physiology* 125.1 (2001): 199-208.

Volume 3 Issue 7 July 2019

© All rights are reserved by Farid A Badria and Walaa S Aboelmaaty.